

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TILE HOMOTOPY GROUPS
Autor: Reid, Michael
Anhang: 7. Appendix: further examples
DOI: <https://doi.org/10.5169/seals-66684>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

shows that \bar{x} commutes with $\bar{x}\bar{y}\bar{x}^{-1}\bar{y}^{-1}$ in the tile path group. Then Theorem 6.1(a) shows that $\pi(\mathcal{T}) \cong \mathbf{Z}/9\mathbf{Z}$. This means that the tile homotopy group only detects area, modulo 9.

On the other hand, we can easily show that if \mathcal{T} tiles a rectangle, then both sides must be even. Consider the ways that a tile can touch the edge of a rectangle.




FIGURE 6.9

Tiles along an edge of a rectangle

We see that the first two possibilities cannot occur, so each tile that touches the edge does so along an even length. Therefore, each edge of the rectangle has even length. In fact, it is not much harder to show that if \mathcal{T} tiles an $m \times n$ rectangle, then both m and n are multiples of 6. A straightforward argument shows that every tiling of a quadrant by \mathcal{T} is a union of 6×6 squares, which implies the result.

7. APPENDIX: FURTHER EXAMPLES

Here we give some more tiling restrictions we have found using the tile homotopy technique. In each case, there are signed tilings that show that the result cannot be obtained by tile homology methods, and there are tilings that show that the result is non-vacuous. Further details will be published elsewhere.

THEOREM 7.1. Let $\mathcal{T} = \left\{ \begin{array}{c} \square \\ \square \end{array} \right\}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then either m or n is a multiple of 4.
- (b) A 1×6 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.2. Let $\mathcal{T} = \left\{ \begin{array}{c} \square \\ \square \end{array}, \begin{array}{c} \square \\ \square \end{array} \right\}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.
- (b) A 1×6 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.3. Let $\mathcal{T} = \{ \text{shape} \}$, where rotations are permitted, but reflections are not.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is even.
 (b) A 1×5 rectangle has a signed tiling by \mathcal{T} .

REMARK 7.4. It is easy to show that if \mathcal{T} tiles a rectangle, then both sides are multiples of 5. Also, Yuri Aksyonov [1] has given a clever geometric proof that one side must be a multiple of 10.

THEOREM 7.5. Let $\mathcal{T} = \{ \text{shape1}, \text{shape2}, \text{shape3}, \text{shape4} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.
 (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.6. Let

$$\mathcal{T} = \{ \text{shape1}, \text{shape2}, \text{shape3}, \text{shape4}, \text{shape5}, \text{shape6}, \text{shape7} \},$$

where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.
 (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.7. Let $\mathcal{T} = \{ \text{shape1}, \text{shape2}, \text{shape3}, \text{shape4}, \text{shape5}, \text{shape6} \}$,

$$\text{shape7}, \text{shape8}, \text{shape9}, \text{shape10}, \text{shape11} \}, \text{ where all orientations are allowed.}$$

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.
 (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.8. Let $\mathcal{T} = \{ \text{shape1}, \text{shape2} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.
 (b) A 2×2 square has a signed tiling by \mathcal{T} .

THEOREM 7.9. Let $\mathcal{T} = \left\{ \begin{array}{c} \square \\ \square \end{array}, \begin{array}{cc} \square & \square \\ \square & \square \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then either m is a multiple of 3 or n is a multiple of 6.

(b) A 1×1 square has a signed tiling by \mathcal{T} .

THEOREM 7.10. Let $\mathcal{T} = \left\{ \begin{array}{cc} \square & \square \\ \square & \square \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array}, \begin{array}{cccccc} \square & \square & \square & \square & \square & \square \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 8.

(b) A 1×1 square has a signed tiling by \mathcal{T} .

THEOREM 7.11. Let $\mathcal{T} = \left\{ \begin{array}{cccc} \square & \square & \square & \square \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 5.

(b) A 1×1 square has a signed tiling by \mathcal{T} .

THEOREM 7.12. Let $\mathcal{T} = \left\{ \begin{array}{cccc} \square & \square & \square & \square \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.

(b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.13. Let $\mathcal{T} = \left\{ \begin{array}{cccc} \square & \square & \square & \square \end{array}, \begin{array}{cc} \square & \square \\ \square & \square \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.

(b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.14. Let $\mathcal{T} = \left\{ \begin{array}{c} \square \\ \square \end{array}, \begin{array}{cccc} \square & \square & \square & \square \end{array}, \begin{array}{ccc} & \square & \\ \square & & \square \\ & \square & \end{array} \right\}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.

(b) A 1×1 square has a signed tiling by \mathcal{T} .

THEOREM 7.15. Let $\mathcal{T} = \{ \text{[rectangle with 6 vertical dashed lines]}, \text{[plus shape]}, \text{[L-shaped polyomino]} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.
- (b) A 2×3 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.16. Let $\mathcal{T} = \{ \text{[triangle with 3 internal dashed lines forming a smaller triangle]} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles a triangle of side n , then n is a multiple of 8.
- (b) A triangle of side 4 has a signed tiling by \mathcal{T} .

REMARK 7.17. That \mathcal{T} tiles any triangle is quite interesting. Karl Scherer [15, 2.6 D] has found a tiling of a side 32 triangle by \mathcal{T} .

ACKNOWLEDGMENT. I thank Torsten Sillke for some interesting discussions.

REFERENCES

- [1] AKSYONOV, YU. E-mail communication to Torsten Sillke. March 1999 (<http://www.mathematik.uni-bielefeld.de/~sillke/PENTA/qu5-y-right>).
- [2] BERLEKAMP, E. R., J. H. CONWAY and R. K. GUY. *Winning Ways for Your Mathematical Plays*, vol. 2. Academic Press, London, 1982.
- [3] BLACK, M. *Critical Thinking*. Prentice-Hall, New York, 1946.
- [4] CONWAY, J. H. and J. C. LAGARIAS. Tiling with polyominoes and combinatorial group theory. *J. Combin. Theory Ser. A*, 53 (1990), 183–208.
- [5] THE GAP GROUP. GAP – Groups, Algorithms and Programming, version 4.3, 2002 (<http://www.gap-system.org>).
- [6] GAREY, M. R. and D. S. JOHNSON. *Computers and Intractability*. Freeman, San Francisco, 1979.
- [7] GOLOMB, S. W. Checker boards and polyominoes. *Amer. Math. Monthly* 61 (1954), 675–682.
- [8] ———. Covering a rectangle with L -tetrominoes, Problem E1543. *American Mathematical Monthly* 69 (1962), 920.
Solution by D. A. KLARNER: *Amer. Math. Monthly* 70 (1963), 760–761.
- [9] HALL, M. JR. *The Theory of Groups*. Chelsea, New York, 1976.
- [10] KLARNER, D. A. Packing a rectangle with congruent N -ominoes. *J. Combin. Theory* 7 (1969), 107–115.
- [11] LANGMAN, H. *Play Mathematics*. Hafner, New York, 1962.