

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 49 (2003)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES  
**Autor:** Gautero, François  
**Kapitel:** 5.2 Straight telescopic paths  
**DOI:** <https://doi.org/10.5169/seals-66690>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 18.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**LEMMA 5.4.** *Let  $p$  be a horizontal geodesic which admits a decomposition in  $r$  subpaths  $p_i$  such that for some constant  $L \geq 0$ , for any  $i = 1, \dots, r$ , either  $|[p_i]_{r+nt_0}|_{r+nt_0} \leq |p_i|_r$  or  $L \geq |[p_i]_{r+nt_0}|_{r+nt_0} > |p_i|_r$ . Then there exists a constant  $C_{5.4}(n, r, L)$ , which is increasing in each variable, such that if  $p$  is dilated in the future after  $nt_0$ , then  $|p|_r \leq C_{5.4}(n, r, L)$ .*

*Proof.* We set  $n = 1$  in order to simplify the notation; the general case is treated in the same way. Up to permuting the indices,  $|[p_i]_{r+t_0}|_{r+t_0} > |p_i|_r$  for  $i = 1, \dots, j$ . Since  $p$  is dilated in the future after  $t_0$ ,

$$jL + \sum_{i=j+1}^r |p_i|_r \geq \lambda^{t_0} \sum_{i=1}^r |p_i|_r.$$

Therefore  $|p|_r \leq \frac{jL}{\lambda^{t_0} - 1}$ .  $\square$

## 5.2 STRAIGHT TELESCOPIC PATHS

**DEFINITION 5.5.** A *straight telescopic path* is a telescopic path  $S$  such that if  $x, y$  are any two points in  $S$  with  $x \in O^+(y) \cup O^-(y)$  then the subpath of  $S$  between  $x$  and  $y$  is equal to the orbit-segment of the semi-flow between  $x$  and  $y$ .

If  $S$  is a path containing a point  $x$ , let  $S_{x,t} \subset S$  be the maximal subpath of  $S$  containing  $x$ , whose pulled-tight projection  $[S_{x,t}]_{f(x)+t}$  on  $f^{-1}(f(x)+t)$  is well defined. The point  $\sigma_t(x)$  does not necessarily belong to  $[S_{x,t}]_{f(x)+t}$ . However there exists a unique point in  $[S_{x,t}]_{f(x)+t}$  which minimizes the horizontal distance between  $\sigma_t(x)$  and  $[S_{x,t}]_{f(x)+t}$ . This point is denoted by  $\bar{x}_t$ . Lemma 5.6 below gives an upper bound, depending on  $t$ , for the telescopic distance between  $x$  and  $\bar{x}_t$ .

**LEMMA 5.6.** *Let  $S$  be any straight telescopic path. If  $t$  is any non negative real number, there exists a constant  $C_{5.6}(t) \geq t$ , which increases with  $t$ , such that any point  $x \in S$  is at telescopic distance smaller than  $C_{5.6}(t)$  from the point  $\bar{x}_t$  (see above).*

*Proof.* If  $\sigma_t(x) \in [S_{x,t}]_{f(x)+t}$ , we set  $C_{5.6}(t) = t$ . Since  $S$  is straight, if  $\sigma_t(x) \notin [S_{x,t}]_{f(x)+t}$ ,  $x$  belongs to a cancellation  $c$  whose endpoints lie in the past orbits of  $\bar{x}_t$ . The bounded-cancellation property gives an upper bound on the horizontal length of  $c$ . This leads to the conclusion.  $\square$