

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 1. Lecture 1
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

proofs are postponed until Lecture 3). In Lecture 2, we explain the origin of the ring of quasi-invariants in the theory of integrable systems, and introduce some tools from integrable systems, such as the Baker-Akhieser function. Finally, in Lecture 3, we develop the theory of the rational Cherednik algebra, the representation-theoretic techniques due to Opdam and Rouquier, and finish the proofs of the geometric statements from Chapter 1.

1. LECTURE 1

1.1 DEFINITION OF QUASI-INVARIANTS

In this lecture we will define the ring of quasi-invariants Q_m and discuss its main properties.

We will work over the field \mathbf{C} of complex numbers. Let W be a finite Coxeter group, i.e. a finite group generated by reflections. Let us denote by \mathfrak{h} its reflection representation. A typical example is the Weyl group of a semisimple Lie algebra acting on a Cartan subalgebra \mathfrak{h} . In the case the Lie algebra is $\mathfrak{sl}(n)$, we have that W is the symmetric group S_n on n letters and \mathfrak{h} is the space of diagonal traceless $n \times n$ matrices.

Let $\Sigma \subset W$ denote the set of reflections. Clearly, W acts on Σ by conjugation. Let $m: \Sigma \rightarrow \mathbf{Z}_+$ be a function on Σ taking non negative integer values, which is W -invariant. The number of orbits of W on Σ is generally very small. For example, if W is the Weyl group of a simple Lie algebra of ADE type, then W acts transitively on Σ , so m is a constant function.

For each reflection $s \in \Sigma$, choose $\alpha_s \in \mathfrak{h}^* - \{0\}$ so that, for $x \in \mathfrak{h}$, $\alpha_s(sx) = -\alpha_s(x)$ (this means that the hyperplane given by the equation $\alpha_s = 0$ is the reflection hyperplane for s).

DEFINITION 1.1 ([CV1, CV2]). A polynomial $q \in \mathbf{C}[\mathfrak{h}]$ is said to be *m -quasi-invariant* with respect to W if, for any $s \in \Sigma$, the polynomial $q(x) - q(sx)$ is divisible by $\alpha_s(x)^{2m_s+1}$.

We will denote by Q_m the space of m -quasi-invariant polynomials with respect to W .

Notice that every element of $\mathbf{C}[\mathfrak{h}]$ is a 0-quasi-invariant, and that every W -invariant is an m -quasi-invariant for any m . Indeed if $q \in \mathbf{C}[\mathfrak{h}]^W$, then we have $q(x) - q(sx) = 0$ for all $s \in \Sigma$, and 0 is divisible by all powers of $\alpha_s(x)$. Thus in a way, $\mathbf{C}[\mathfrak{h}]^W$ can be viewed as the set of ∞ -quasi-invariants.

EXAMPLE 1.2. The group $W = \mathbf{Z}/2$ acts on $\mathfrak{h} = \mathbf{C}$ by $s(v) = -v$. In this case m is a non negative integer and $\Sigma = \{s\}$. So this definition says that q is in Q_m iff $q(x) - q(-x)$ is divisible by x^{2m+1} . It is very easy to write a basis of Q_m . It is given by the polynomials $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$.

1.2 ELEMENTARY PROPERTIES OF Q_m

Some elementary properties of Q_m are collected in the following proposition.

PROPOSITION 1.3 (see [FV] and references therein).

- 1) $\mathbf{C}[\mathfrak{h}]^W \subset Q_m \subseteq \mathbf{C}[\mathfrak{h}]$, $Q_0 = \mathbf{C}[\mathfrak{h}]$, $Q_m \subset Q_{m'}$ if $m \geq m'$, $\bigcap_m Q_m = \mathbf{C}[\mathfrak{h}]^W$.
- 2) Q_m is a graded subalgebra of $\mathbf{C}[\mathfrak{h}]$.
- 3) The fraction field of Q_m is equal to $\mathbf{C}(\mathfrak{h})$.
- 4) Q_m is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and a finitely generated algebra. $\mathbf{C}[\mathfrak{h}]$ is a finite Q_m -module.

Proof. 1) is immediate and has already been mentioned in 1.1.

2) Clearly Q_m is closed under addition. Let $p, q \in Q_m$. Let $s \in \Sigma$. Then

$$p(x)q(x) - p(sx)q(sx) = (p(x) - p(sx))q(x) + p(sx)(q(x) - q(sx)).$$

Since both $p(x) - p(sx)$ and $q(x) - q(sx)$ are divisible by $\alpha_s^{2m_s+1}$, we deduce that $p(x)q(x) - p(sx)q(sx)$ is also divisible by $\alpha_s^{2m_s+1}$, proving the claim.

3) Consider the polynomial

$$\delta_{2m+1}(x) = \prod_{s \in \Sigma} \alpha_s(x)^{2m_s+1}.$$

This polynomial is uniquely defined up to scaling. One has $\delta_{2m+1}(sx) = -\delta_{2m+1}(x)$ for each $s \in \Sigma$, hence $\delta_{2m+1} \in Q_m$. Take $f(x) \in \mathbf{C}[\mathfrak{h}]$. We claim that $f(x)\delta_{2m+1}(x) \in Q_m$. As a matter of fact,

$$f(x)\delta_{2m+1}(x) - f(sx)\delta_{2m+1}(sx) = (f(x) + f(sx))\delta_{2m+1}(x),$$

and by its definition $\delta_{2m+1}(x)$ is divisible by $\alpha_s(x)^{2m_s+1}$ for all $s \in \Sigma$. This implies 3).

4) By Hilbert's theorem on the finiteness of invariants, we get that $\mathbf{C}[\mathfrak{h}]^W$ is a finitely generated algebra over \mathbf{C} and $\mathbf{C}[\mathfrak{h}]$ is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and hence a finite Q_m -module, proving the second part of 4).

Now $Q_m \subset \mathbf{C}[\mathfrak{h}]$ is a submodule of the finite module $\mathbf{C}[\mathfrak{h}]$ over the Noetherian ring $\mathbf{C}[\mathfrak{h}]^W$. Hence it is finite. This immediately implies that Q_m is a finitely generated algebra over \mathbf{C} . \square

REMARK. In fact, since W is a finite Coxeter group, a celebrated result of Chevalley says that the algebra $\mathbf{C}[\mathfrak{h}]^W$ is not only a finitely generated \mathbf{C} -algebra but actually a free (=polynomial) algebra. Namely, it is of the form $\mathbf{C}[q_1, \dots, q_n]$, where the q_i are homogeneous polynomials of some degrees d_i . Furthermore, if we denote by H the subspace of $\mathbf{C}[\mathfrak{h}]$ of harmonic polynomials, i.e. of polynomials killed by W -invariant differential operators with constant coefficients without constant term, then the multiplication map

$$\mathbf{C}[\mathfrak{h}]^W \otimes H \rightarrow \mathbf{C}[\mathfrak{h}]$$

is an isomorphism of $\mathbf{C}[\mathfrak{h}]^W$ - and of W -modules. In particular, $\mathbf{C}[\mathfrak{h}]$ is a free $\mathbf{C}[\mathfrak{h}]^W$ -module of rank $|W|$.

1.3 THE VARIETY X_m AND ITS BIJECTIVE NORMALIZATION

Using Proposition 1.3, we can define the irreducible affine variety $X_m = \text{Spec}(Q_m)$. The inclusion $Q_m \subset \mathbf{C}[\mathfrak{h}]$ induces a morphism

$$\pi: \mathfrak{h} \rightarrow X_m,$$

which again by Proposition 1.3 is birational and surjective. (Notice that in particular this implies that X_m is singular for all $m \neq 0$.)

In fact, not only is π birational, but a stronger result is true.

PROPOSITION 1.4 (Berest, see [BEG]). *π is a bijection.*

Proof. By the above remarks, we only have to show that π is injective. In order to achieve this, we need to prove that quasi-invariants separate points of \mathfrak{h} , i.e. that if $z, y \in \mathfrak{h}$ and $z \neq y$, then there exists $p \in Q_m$ such that $p(z) \neq p(y)$. This is obtained in the following way. Let $W_z \subset W$ be the stabilizer of z and choose $f \in \mathbf{C}[\mathfrak{h}]$ such that $f(z) \neq 0$, $f(y) = 0$. Set

$$p(x) = \prod_{s \in \Sigma, sz \neq z} \alpha_s(x)^{2m_s+1} \prod_{w \in W_z} f(wx).$$

We claim that $p(x) \in Q_m$. Indeed, let $s \in \Sigma$ and assume that $s(z) \neq z$.

We have by definition $p(x) = \alpha_s(x)^{2m_s+1} \tilde{p}(x)$, with $\tilde{p}(x)$ a polynomial. So

$$p(x) - p(sx) = \alpha_s(x)^{2m_s+1} \tilde{p}(x) - \alpha_s(sx)^{2m_s+1} \tilde{p}(sx) = \alpha_s(x)^{2m_s+1} (\tilde{p}(x) + \tilde{p}(sx)).$$

If on the other hand, $sz = z$, i.e. $s \in W_z$, then s preserves the set $W \setminus W_z$, and hence preserves $\prod_{s \in \Sigma \cap (W \setminus W_z)} \alpha_s(x)^{2m_s+1}$ (as it acts by -1 on the products $\prod_{s \in \Sigma} \alpha_s(x)^{2m_s+1}$ and $\prod_{s \in \Sigma \cap W_z} \alpha_s(x)^{2m_s+1}$). Since $\prod_{w \in W_z} f(wx)$ is

W_z -invariant, we deduce that $p(x) - p(sx) = 0$, so that in this case $p(x) - p(sx)$ also is divisible by $\alpha_s(x)^{2m_s+1}$.

To conclude, notice that $p(z) \neq 0$. Indeed, for a reflection s , α_s vanishes exactly on the fixed points of s , so that $\prod_{s \in \Sigma, sz \neq z} \alpha_s(z)^{2m_s+1} \neq 0$. Also for all $w \in W_z$ $f(wz) = f(z) \neq 0$. On the other hand, it is clear that $p(y) = 0$. \square

EXAMPLE 1.5. Take $W = \mathbf{Z}/2$. As we have already seen, Q_m has a basis given by the monomials $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$. From this we deduce that setting $z = x^2$ and $y = x^{2m+1}$, $Q_m = \mathbf{C}[y, z]/(y^2 - z^{2m+1}) = \mathbf{C}[K]$, where K is the plane curve with a cusp at the origin, given by the equation $y^2 = z^{2m+1}$. The map $\pi: \mathbf{C} \rightarrow K$ is given by $\pi(t) = (t^{2m+1}, t^2)$, which is clearly bijective.

1.4 FURTHER PROPERTIES OF X_m

Let us get to some deeper properties of quasi-invariants. Let X be an irreducible affine variety over \mathbf{C} and $A = \mathbf{C}[X]$. Recall that, by the Noether Normalization Lemma, there exist $f_1, \dots, f_n \in \mathbf{C}[X]$ which are algebraically independent over \mathbf{C} and such that $\mathbf{C}[X]$ is a finite module over the polynomial ring $\mathbf{C}[f_1, \dots, f_n]$. This means that we have a finite morphism of X onto an affine space.

DEFINITION 1.6. A (and X) is said to be *Cohen-Macaulay* if there exist f_1, \dots, f_n as above, with the property that $\mathbf{C}[X]$ is a locally free module over $\mathbf{C}[f_1, \dots, f_n]$. (Notice that by the Quillen-Suslin theorem, this is equivalent to saying that A is a free module.)

REMARK. If A is Cohen-Macaulay, then for any f_1, \dots, f_n which are algebraically independent over \mathbf{C} and such that A is a finite module over the polynomial ring $\mathbf{C}[f_1, \dots, f_n]$, we have that A is a locally free $\mathbf{C}[f_1, \dots, f_n]$ -module, see [Eis], Corollary 18.17.

THEOREM 1.7 ([EG2], [BEG], conjectured in [FV]). Q_m is Cohen-Macaulay.

Notice that, using Chevalley's result that $\mathbf{C}[\mathfrak{h}]^W$ is a polynomial ring, it will suffice, in order to prove Theorem 1.7, to prove:

THEOREM 1.8 ([EG2, BEG], conjectured in [FV]). Q_m is a free $\mathbf{C}[\mathfrak{h}]^W$ -module.

We show how one can prove this Theorem in 3.10. This proof follows [BEG] (the original proof of [EG2] is shorter but somewhat less conceptual). The main idea of the proof is to show that the $\mathbf{C}[\hbar]^W$ -module Q_m can be extended to a module over a bigger (noncommutative) algebra, namely the spherical subalgebra of the rational Cherednik algebra. Furthermore, this module belongs to an appropriate category of representations of this algebra, called category \mathcal{O} . On the other hand, it can be shown that any module over the spherical subalgebra that belongs to this category is free when restricted to the commutative algebra $\mathbf{C}[\hbar]^W$.

1.5 THE POINCARÉ SERIES OF Q_m

Consider now the Poincaré series

$$h_{Q_m}(t) = \sum_{r \geq 0} \dim Q_m[r] t^r,$$

where $Q_m[r]$ denotes the graded component of Q_m of degree r . For every irreducible representation $\tau \in \widehat{W}$, define

$$\chi_\tau(t) = \sum_{r \geq 0} \dim \operatorname{Hom}_W(\tau, \mathbf{C}[\hbar][r]) t^r.$$

Consider the element in the group ring $\mathbf{Z}[W]$

$$\mu_m = \sum_{s \in \Sigma} m_s (1 - s).$$

The W -invariance of m implies that μ_m lies in the center of $\mathbf{Z}[W]$. Hence it is clear that μ_m acts as a scalar, $\xi_m(\tau)$, on τ . Let d_τ be the degree of τ .

LEMMA 1.9. *The scalar $\xi_m(\tau)$ is an integer.*

Proof. $\mathbf{Z}[W]$ and hence also its center, is a finite \mathbf{Z} -module. This clearly implies that $\xi_m(\tau)$ is an algebraic integer. Thus to prove that $\xi_m(\tau)$ is an integer, it suffices to see that $\xi_m(\tau)$ is a rational number. Let $d_{\tau,s}$ be the dimension of the space of s -invariants in τ . Taking traces we get

$$d_\tau \xi_m(\tau) = \sum_{s \in \Sigma} 2m_s (d_\tau - d_{\tau,s}),$$

which gives the rationality of $\xi_m(\tau)$. \square

THEOREM 1.10. *One has*

$$(1) \quad h_{Q_m}(t) = \sum_{\tau \in \widehat{W}} d_{\tau} t^{\xi_m(\tau)} \chi_{\tau}(t).$$

REMARK. This theorem was proved in [FeV] modulo Theorem 1.7 (conjectured in [FV]) using the so-called Matsuo-Cherednik correspondence (see [FeV] for details). Thus, Theorem 1.10 follows from [FeV] and [EG2]. Another proof of this theorem is given in [BEG]; this is the proof we will discuss below (in Lecture 3).

EXAMPLE 1.11. If $m = 0$, since $Q_0 = \mathbf{C}[\mathfrak{h}]$, the theorem says that

$$h_{Q_0}(t) = \frac{1}{(1-t)^n} = \sum_{\tau \in \widehat{W}} d_{\tau} \chi_{\tau}(t).$$

Indeed, as a W -module one has

$$\mathbf{C}[\mathfrak{h}] = \bigoplus_{\tau} \tau \otimes \operatorname{Hom}_W(\tau, \mathbf{C}[\mathfrak{h}]).$$

EXAMPLE 1.12. If $W = \mathbf{Z}/2$, then $\widehat{W} = \{+, -\}$, where $+$ (respectively $-$) denotes the trivial (respectively the sign) representation. One has

$$\mathbf{C}[x] = \mathbf{C}[x^2] \oplus \mathbf{C}[x^2]x,$$

where $\mathbf{C}[x^2] = \mathbf{C}[x]^W$ and $\mathbf{C}[x^2]x$ is the isotypic component of the sign representation. Thus

$$\chi_+(t) = \frac{1}{1-t^2}, \quad \chi_-(t) = \frac{t}{1-t^2},$$

$\mu_m = m(1-s)$. Thus $\xi_m(+) = 0$, $\xi_m(-) = 2m$. We deduce that

$$h_{Q_m}(t) = \frac{1}{1-t^2} + \frac{t^{2m+1}}{1-t^2},$$

as we already know.

Recall now that as a graded W -module $\mathbf{C}[\mathfrak{h}]$ is isomorphic to $\mathbf{C}[\mathfrak{h}]^W \otimes H$, H being the space of harmonic polynomials. We deduce that the τ -isotypic component in $\mathbf{C}[\mathfrak{h}]$ is isomorphic to $\mathbf{C}[\mathfrak{h}]^W \otimes H_{\tau}$.

Set $K_\tau(t) = \sum_{r \geq 0} \dim \operatorname{Hom}_W(\tau, H[r])t^r$. This is a polynomial, called the Kostka polynomial relative to τ . We deduce that

$$(2) \quad \chi_\tau(t) = \frac{K_\tau(t)}{\prod_{i=1}^n (1 - t^{d_i})}.$$

Also, if $\tau' = \tau \otimes \varepsilon$, ε being the sign representation, one has

$$K_{\tau'}(t) = K_\tau(t^{-1})t^{|\Sigma|}.$$

Set now

$$P_m(t) = \sum_{\tau \in \widehat{W}} d_\tau t^{\xi_m(\tau)} K_\tau(t).$$

We have

PROPOSITION 1.13 ([FeV]).

$$h_{Q_m}(t) = \frac{P_m(t)}{\prod_{i=1}^n (1 - t^{d_i})}.$$

Furthermore $P_m(t) = t^{\xi_m(\varepsilon) + |\Sigma|} P_m(t^{-1})$.

Proof. Substituting the expression (2) for $\chi_\tau(t)$ in (1.10) and using the definition of $P_m(t)$, we get

$$h_{Q_m}(t) = \sum_{\tau \in \widehat{W}} d_\tau t^{\xi_m(\tau)} \frac{K_\tau(t)}{\prod_{i=1}^n (1 - t^{d_i})} = \frac{P_m(t)}{\prod_{i=1}^n (1 - t^{d_i})},$$

as desired.

Now notice that

$$\xi_m(\tau) + \xi_m(\tau') = \sum_{s \in \Sigma} 2m_s = \xi_m(\varepsilon).$$

Using this we get

$$\begin{aligned} t^{\xi_m(\varepsilon) + |\Sigma|} P_m(t^{-1}) &= \sum_{\tau \in \widehat{W}} d_\tau t^{\xi_m(\varepsilon) - \xi_m(\tau)} t^{|\Sigma|} K_\tau(t^{-1}) \\ &= \sum_{\tau' \in \widehat{W}} d_{\tau'} t^{\xi_m(\tau')} K_{\tau'}(t) = P_m(t), \end{aligned}$$

as desired. \square

From this we deduce

THEOREM 1.14 ([EG2, BEG, FeV], conjectured in [FV]). *The ring Q_m of m -quasi-invariants is Gorenstein.*

Proof. By Stanley's theorem (see [Eis]), a positively graded Cohen-Macaulay domain A is Gorenstein iff its Poincaré series is a rational function $h(t)$ satisfying the equation $h(t^{-1}) = (-1)^n t^l h(t)$, where l is an integer and n is the dimension of the spectrum of A . Thus the result follows immediately from Proposition 1.13. \square

1.6 THE RING OF DIFFERENTIAL OPERATORS ON X_m

Finally, let us introduce the ring $\mathcal{D}(X_m)$ of differential operators on X_m , that is the ring of differential operators with coefficients in $\mathbb{C}(\mathfrak{h})$ mapping Q_m to Q_m . It is clear that this definition coincides with Grothendieck's well-known definition ([Bj]).

THEOREM 1.15 ([BEG]). *$\mathcal{D}(X_m)$ is a simple algebra.*

REMARK 1.16. a) The ring of differential operators on a smooth affine algebraic variety is always simple (see [Bj], Chapter 3).

b) By a result of M. van den Bergh [VdB], for a non-smooth variety, the simplicity of the ring of differential operators implies the Cohen-Macaulay property of this variety.

2. LECTURE 2

We will now see how the ring Q_m appears in the theory of completely integrable systems.

2.1 HAMILTONIAN MECHANICS AND INTEGRABLE SYSTEMS

Recall the basic setup of Hamiltonian mechanics [Ar]. Consider a mechanical system with configuration space X (a smooth manifold). Then the phase space of this system is T^*X , the cotangent bundle on X . The space T^*X is naturally a symplectic manifold, and in particular we have an operation of Poisson bracket on functions on T^*X . A point of T^*X is a pair (x, p) , where $x \in X$ is the position and $p \in T_x^*X$ is the momentum. Such pairs are