

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 49 (2003)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA  
**Autor:** Etingof, Pavel / Strickland, Elisabetta  
**Kapitel:** 2.2 The classical Calogero-Moser System  
**DOI:** <https://doi.org/10.5169/seals-66677>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 18.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

called states of the system. The dynamics of the system  $x = x(t)$ ,  $p = p(t)$  depends on the Hamiltonian, or energy function,  $E(x, p)$  on  $T^*X$ . Given  $E$  and the initial state  $x(0)$ ,  $p(0)$ , one can recover the dynamics  $x = x(t)$ ,  $p = p(t)$  from Hamilton's differential equations  $\frac{df(x, p)}{dt} = \{f, E\}$ . If  $X$  is locally identified with  $\mathbf{R}^n$  by choosing coordinates  $x_1, \dots, x_n$ , then  $T^*X$  is locally identified with  $\mathbf{R}^{2n}$  with coordinates  $x_1, \dots, x_n, p_1, \dots, p_n$ . In these coordinates, Hamilton's equations may be written in their standard form

$$\dot{x}_i = \frac{\partial E}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial E}{\partial x_i}.$$

A function  $I(x, p)$  is called an integral of motion for our system if  $\{I, E\} = 0$ . Integrals of motion are useful, since for any such integral  $I$  the function  $I(x(t), p(t))$  is constant, which allows one to reduce the number of variables by 2. Thus, if we are given  $n$  functionally independent integrals of motion  $I_1, \dots, I_n$  with  $\{I_l, I_k\} = 0$  for all  $1 \leq l, k \leq n$ , then all  $2n$  variables  $x_i, p_i$  can be excluded, and the system can be completely solved by quadratures. Such a situation is called complete (or Liouville) integrability.

## 2.2 THE CLASSICAL CALOGERO-MOSER SYSTEM

Quasi-invariants are related to many-particle systems. Consider a system of  $n$  particles on the real line  $\mathbf{R}$ . A potential is an even function

$$U(x) = U(-x), \quad x \in \mathbf{R}.$$

Two particles at points  $a, b$  have energy of interaction  $U(a - b)$ . The total energy of our system of particles is

$$E = \sum_{i=1}^n \frac{p_i^2}{2} + \sum_{i < j} U(x_i - x_j).$$

Here,  $x_i$  are the coordinates of the particles,  $p_i$  their momenta. The dynamics of the particles  $x_i = x_i(t)$ ,  $p_i = p_i(t)$  is governed by the Hamilton equations with energy function  $E$ .

This is a system of nonlinear differential equations, which in general can be difficult to solve explicitly. However, for special potentials this system might be completely integrable. For instance, we will see that this is the case for the Calogero-Moser potential,

$$U(x) = \frac{\gamma}{x^2},$$

$\gamma$  being a constant.

The Calogero-Moser system has a generalization to arbitrary Coxeter groups. Namely, consider a finite group  $W$  generated by reflections acting on the space  $\mathfrak{h}$ , and keep the notation of the previous section. Fix a  $W$ -invariant nondegenerate scalar product  $(-, -)$  on  $\mathfrak{h}$ . It determines a scalar product on  $\mathfrak{h}^*$ . Define the “energy function”

$$E(x, p) = \frac{(p, p)}{2} + \frac{1}{2} \sum_{s \in \Sigma} \frac{\gamma_s(\alpha_s, \alpha_s)}{\alpha_s(x)^2}, \quad x \in \mathfrak{h}, \quad p \in \mathfrak{h}^*$$

on  $T^*\mathfrak{h} = \mathfrak{h} \times \mathfrak{h}^*$ , where  $\gamma: \Sigma \rightarrow \mathbf{C}$  is a  $W$ -invariant function. Notice that although  $\alpha_s$  is defined up to a non zero constant, by homogeneity,  $E$  is independent of the choice of  $\alpha_s$ . We will call the system defined by  $E$  the Calogero-Moser system for  $W$ .

If  $W$  is the symmetric group  $S_n$ ,  $\mathfrak{h} = \mathbf{C}^n$ , then  $\Sigma$  is the set of transpositions  $s_{i,j}$ ,  $i < j$ , and we can take  $\alpha_s = e_i - e_j$ . Then we clearly obtain the usual Calogero-Moser system.

Below we will see that the Calogero-Moser system for  $W$  is completely integrable.

### 2.3 THE QUANTUM CALOGERO-MOSER SYSTEM

Let us now discuss quantization of the Calogero-Moser system. We start by quantizing the energy  $E$  by formally making the substitution

$$p_j \Rightarrow -i\hbar \frac{\partial}{\partial x_j},$$

where  $\hbar$  is a parameter (Planck's constant). This yields the Schrödinger operator

$$\widehat{E} := -\frac{\hbar^2}{2} \Delta + \frac{1}{2} \sum_{s \in \Sigma} \frac{\gamma_s(\alpha_s, \alpha_s)}{\alpha_s^2},$$

where  $\Delta$  denotes the Laplacian.

In particular, in the case of  $W = S_n$  we have

$$\widehat{E} = -\frac{\hbar^2}{2} \Delta + \sum_{i < j} \frac{c}{(x_i - x_j)^2},$$

where  $\Delta = \sum_i \frac{\partial^2}{\partial x_i^2}$ . Setting  $\beta_s = \frac{\gamma_s}{2\hbar^2}$ , we will from now on consider the operator

$$H := -\frac{2}{\hbar^2} \widehat{E} = \Delta - \sum_{s \in \Sigma} \frac{\beta_s(\alpha_s, \alpha_s)}{\alpha_s^2(x)},$$

called the Calogero-Moser operator.