

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 3.6 Category O
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

DEFINITION 3.13. The *spherical subalgebra* of H_c is the algebra $eH_c e$.

Notice that $1 \notin eH_c e$. On the other hand, since $ex = xe = e$ for $x \in eH_c e$, e is the unit for the spherical subalgebra. We can embed both $\mathbf{C}[\mathfrak{h}^*]^W$ and $\mathbf{C}[\mathfrak{h}]^W$ in the spherical subalgebra as follows. Take $f \in \mathbf{C}[\mathfrak{h}^*]^W$ (the other case is identical) and set $m_e(f) = fe$. Since f is invariant, we have $efe = fe^2 = fe = m_e(f)$, so that m_e actually maps $\mathbf{C}[\mathfrak{h}^*]^W$ to $eH_c e$. The injectivity is clear from the PBW-theorem. As for the fact that m_e is a homomorphism, we have $m_e(fg) = fge = fge^2 = fege = m_e(f)m_e(g)$. From now on, we will consider both $\mathbf{C}[\mathfrak{h}^*]^W$ and $\mathbf{C}[\mathfrak{h}]^W$ as subalgebras of the spherical subalgebra.

3.6 CATEGORY \mathcal{O}

We are now going to study representations of the algebras H_c and $eH_c e$.

DEFINITION 3.14. The category $\mathcal{O}(H_c)$ (resp. $\mathcal{O}(eH_c e)$) is the full subcategory of the category of H_c -modules (resp. $eH_c e$ -modules) whose objects are the modules M such that

- 1) M is finitely generated.
- 2) For all $v \in M$, the subspace $\mathbf{C}[\mathfrak{h}^*]^W v \subset M$ is finite dimensional.

We can define a functor

$$F: \mathcal{O}(H_c) \rightarrow \mathcal{O}(eH_c e)$$

by setting $F(M) = eM$. It is easy to show that $F(M)$ is an object of $\mathcal{O}(eH_c e)$.

We are now going to explain how to construct some modules in $\mathcal{O}(H_c)$ which, by analogy with the case of enveloping algebras of semisimple Lie algebras, we will call Whittaker and Verma modules. First, take $\lambda \in \mathfrak{h}^*$. Denote by $W_\lambda \subset W$ the stabilizer of λ . Take an irreducible W_λ -module τ . We define a structure of $\mathbf{C}[\mathfrak{h}^*] \rtimes \mathbf{C}[W_\lambda]$ -module on τ by

$$(fw)v = f(\lambda)(wv) \quad \forall v \in \tau, w \in W_\lambda, f \in \mathbf{C}[\mathfrak{h}^*].$$

It is easy to see that this action is well defined and we denote this module by $\lambda \# \tau$. We can then consider the H_c -module

$$M(\lambda, \tau) = H_c \otimes_{\mathbf{C}[\mathfrak{h}^*] \rtimes \mathbf{C}[W_\lambda]} \lambda \# \tau.$$

This is called a Whittaker module. In the special case $\lambda = 0$ (and hence $W_\lambda = W$), the module $M(0, \tau)$ is called a Verma module. It is clear that these are objects of \mathcal{O} . Notice that as $\mathbf{C}[\mathfrak{h}] \rtimes \mathbf{C}[W]$ -module, $M(\lambda, \tau) = \mathbf{C}[\mathfrak{h}] \otimes_{\mathbf{C}} \mathbf{C}[W] \otimes_{\mathbf{C}[W_\lambda]} \tau$.

EXAMPLE 3.15. If $\lambda = 0$ and $\tau = \mathbf{1}$ is the trivial representation of W , the Verma module $M(0, \mathbf{1}) = \mathbf{C}[\mathfrak{h}]$. The action of $\mathbf{C}[\mathfrak{h}]$ is given by multiplication, that of $\mathbf{C}[\mathfrak{h}^*]$ is generated by the Dunkl operators and W acts in the usual way.

3.7 GENERIC c

Opdam and Rouquier have recently studied the structure of the categories $\mathcal{O}(H_c)$, $\mathcal{O}(eH_c e)$, and found that it is especially simple if c is “generic” in a certain sense. Namely, recall that for a W -invariant function $q: \Sigma \rightarrow \mathbf{C}^*$ one can define the *Hecke algebra* $\text{He}_q(W)$ to be the quotient of the group algebra of the fundamental group of U/W by the relations $(T_s - 1)(T_s + q_s) = 0$, where T_s is the image in U/W of a small half-circle around the hyperplane of s in the counterclockwise direction. It is well known that $\text{He}_q(W)$ is an algebra of dimension $|W|$, which coincides with $\mathbf{C}[W]$ if $q = 1$. It is also known that $\text{He}_q(W)$ is semisimple (and isomorphic to $\mathbf{C}[W]$ as an algebra) unless q_s belongs for some s to a finite set of roots of unity depending on W (see [Hu]).

DEFINITION 3.16. The function c is said to be *generic* if for $q = e^{2\pi ic}$, the Hecke algebra $\text{He}_q(W)$ is semisimple.

In particular, any irrational c is generic, and (more important for us) an integer valued c is generic (since in this case $q = 1$). We can now state the following central result:

THEOREM 3.17 (Opdam-Rouquier [OR]; see also [BEG] for an exposition). *If c is generic (in particular, if c takes non negative integer values), then the irreducible objects in \mathcal{O} are exactly the modules $M(\lambda, \tau)$. Moreover, the category \mathcal{O} is semisimple.*

We also have

THEOREM 3.18 ([OR]). *If c is generic then the functor F is an equivalence of categories.*

From Theorem 3.17 we can deduce

THEOREM 3.19 ([BEG]). *If c is generic, then H_c is a simple algebra.*

In the case $c = 0$, we get the simplicity of $\mathbf{C}[\mathfrak{h} \oplus \mathfrak{h}^*] \rtimes \mathbf{C}[W]$, which is well known.