

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 3.11 Proof of Theorem 1.15
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The following important theorem shows that this action extends to Q_m .

THEOREM 3.22 ([BEG]). *There exists a unique representation of the algebra eH_me on Q_m in which an element $q \in \mathbf{C}[\mathfrak{h}]^W$ acts by multiplication and an element $q \in \mathbf{C}[\mathfrak{h}^*]^W$ by L_q .*

Proof. Since by Proposition 3.5, L_q preserves Q_m , we get a uniquely defined representation of the subalgebra of eH_me generated by $\mathbf{C}[\mathfrak{h}]^W$ and $\mathbf{C}[\mathfrak{h}^*]^W$ on Q_m . The result now follows from Theorem 3.21. \square

3.10 PROOF OF THEOREM 1.8

Finally we can prove Theorem 1.8.

To do this, observe that as an eH_me -module, Q_m is in the category $\mathcal{O}(eH_me)$, and $\mathbf{C}[\mathfrak{h}^*]^W$ acts locally nilpotently in Q_m (by degree arguments). We can now apply Theorem 3.18 and Theorem 3.17 and deduce that Q_m is a direct sum of modules of the form $eM(0, \tau)$. As a $\mathbf{C}[\mathfrak{h}] \rtimes \mathbf{C}[W]$ -module, $M(0, \tau) = \mathbf{C}[\mathfrak{h}] \otimes \tau$. On the other hand, by Chevalley's theorem, there is an isomorphism $\mathbf{C}[\mathfrak{h}] \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \mathbf{C}[W]$, commuting with the action of W and $\mathbf{C}[\mathfrak{h}]^W$. Thus we get an isomorphisms of $\mathbf{C}[\mathfrak{h}]^W$ -modules

$$eM(0, \tau) \simeq (M(0, \tau))^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes (\mathbf{C}[W] \otimes \tau)^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \tau,$$

proving that $eM(0, \tau)$ and hence Q_m is a free $\mathbf{C}[\mathfrak{h}]^W$ -module. \square

EXAMPLE 3.23. For $W = \mathbf{Z}/2$ and $\mathfrak{h} = \mathbf{C}$, take the polynomials $1, x^{2m+1}$. Notice that $L(1) = L(x^{2m+1}) = 0$ while $s(1) = 1, s(x^{2m+1}) = -x^{2m+1}, s \in \mathbf{Z}/2$ being the element of order two. It follows that Q_m as a eH_me -module is the direct sum of $\mathbf{C}[x^2] \oplus x^{2m+1}\mathbf{C}[x^2]$. These modules are irreducible. Moreover, $\mathbf{C}[x^2] \simeq eM(0, \mathbf{1}), x^{2m+1}\mathbf{C}[x^2] \simeq eM(0, \varepsilon), \varepsilon$ being the sign representation.

3.11 PROOF OF THEOREM 1.15

Let I be a nonzero two-sided ideal in $\mathcal{D}(X_m)$. First we claim that I nontrivially intersects Q_m . Indeed, otherwise let $K \in I$ be a lowest order nonzero element in I . Since the order of K is positive, there exists $f \in Q_m$ such that $[K, f] \neq 0$. Then $[K, f] \in I$ is of smaller order than K , a contradiction.

Now let $f \in Q_m$ be an element of I . Then $g = \prod_{w \in W} {}^w f \in I$. But g is W -invariant. This shows that the intersection J of I with the subalgebra H_m in $\mathcal{D}(X_m)$ is nonzero. But H_m is simple by Theorem 3.19, so $J = H_m$. Hence, $1 \in J \subset I$, and $I = \mathcal{D}(X_m)$. \square