

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ADDITIVE NUMBER THEORY SHEDS EXTRA LIGHT ON THE HOPF-STIEFEL o FUNCTION
Autor: Plagne, Alain
Kurzfassung
DOI: <https://doi.org/10.5169/seals-66682>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 13.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ADDITIVE NUMBER THEORY SHEDS EXTRA LIGHT ON THE HOPF-STIEFEL \circ FUNCTION

by Alain PLAGNE

ABSTRACT. The famous Hopf-Stiefel \circ function appears in several places in mathematics (linear and bilinear algebra, topology, intercalate matrices, . . .). However, although the object of much study, this function kept a part of mystery since no simple formula was known for it. We shall derive a simple and practical explicit formula for \circ and more generally for β_p (p arbitrary prime), a generalized function due to Eliahou and Kervaire. The proof relies on a new result in combinatorial group theory which follows from additive number theoretical arguments. It is shown that this last result generalizes earlier ones by Eliahou and Kervaire and by Yuzvinsky.

1. INTRODUCTION

A *composition formula* of size $[r, s, n]$ over some field \mathbf{F} (that we assume to be of characteristic different from 2) is an identity of the form

$$(x_1^2 + \cdots + x_r^2)(y_1^2 + \cdots + y_s^2) = (z_1^2 + \cdots + z_n^2),$$

where z_1, z_2, \dots, z_n are n bilinear forms in the variables (x_1, \dots, x_r) and (y_1, \dots, y_s) , with coefficients in \mathbf{F} . For example, the law of moduli for complex numbers provides the identity

$$(x_1^2 + x_2^2)(y_1^2 + y_2^2) = (x_1y_1 - x_2y_2)^2 + (x_2y_1 + x_1y_2)^2,$$

which is a composition formula of size $[2, 2, 2]$. The law of moduli for quaternions (respectively for octonions) provides a composition formula of size $[4, 4, 4]$ (respectively $[8, 8, 8]$) in a similar way. Conversely, Hurwitz's theorem [7] (see also [8]) states that the only possible values of n for which a composition formula of size $[n, n, n]$ exists are 1, 2, 4 and 8.