

Zeitschrift: L'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Isomorphism conjectures for the mapping class group
Autor: Davis, James F.
DOI: <https://doi.org/10.5169/seals-109895>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 07.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

26

ISOMORPHISM CONJECTURES FOR THE MAPPING CLASS GROUP

by James F. DAVIS

I enjoyed Guido's lecture in Münster where he showed that Teichmüller space is a classifying space for actions of the mapping class group whose isotropy groups are all finite. I will pose a question and then recall one I posed at that time.

Let Σ_g be a closed surface of genus g , let Γ_g be its mapping class group, and $\tau_g \cong \mathbf{R}^{6g-6}$ its Teichmüller space.

QUESTION 26.1 (A special case of the Borel Conjecture). *Let Γ be a torsion free subgroup of the mapping class group, for example the Torelli group. Is the action of Γ on τ_g topologically rigid? That is, is any proper homotopy equivalence of manifolds $h: M \rightarrow \tau_g/\Gamma$, which is a homeomorphism outside of a compact set, homotopic to a homeomorphism, via a homotopy which is constant outside of a compact set?*

QUESTION 26.2 (Isomorphism conjecture injectivity). *Is there a contractible compactification of Teichmüller space which is small at infinity and equivariant with respect to the action of the mapping class group?*

To say a compactification $\bar{\tau}_g$ of τ_g is *small at infinity* means that for every compact subset K of Teichmüller space, for every boundary point $y \in \bar{\tau}_g - \tau_g$, and for every neighborhood U of y in the compactification $\bar{\tau}_g$, there is a smaller neighborhood V so that for every $\gamma \in \Gamma_g$, if $\gamma K \cap V$ is nonempty, then $\gamma K \subset U$.

DISCUSSION. A solution to Question 26.1 would likely involve carrying out the program of Farrell–Jones [1] in the mapping class group case. A positive solution to Question 26.2 would likely lead to a proof of the injectivity map

of the assembly map in K - and L -theory with respect to the family of finite subgroups (by modifying Rosenthal's thesis [3]), and thereby a new proof of the Novikov conjecture in this case. (It seems that the Novikov conjecture in the mapping class group case has been recently proved by Ursula Hamenstädt [2].)

REFERENCES

- [1] FARRELL, F. T. and L. E. JONES. Isomorphism conjectures in algebraic K -theory. *J. Amer. Math. Soc.* 6 (1993), 249–297.
- [2] HAMENSTÄDT, U. Geometry of the mapping class groups I: Boundary amenability. Preprint arXiv: math.GR/0510116 (2005–2008).
- [3] ROSENTHAL, D. Splitting with continuous control in algebraic K -theory. *K-Theory* 32 (2004), 139–166.

James F. Davis

Department of Mathematics
Indiana University
Bloomington, Indiana 47405
USA
e-mail: jfdavis@indiana.edu