

Zeitschrift: L'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Exotic classes in the mod 2 cohomology of $GL_n(\mathbb{Z}[\frac{1}{2}])$
Autor: Henn, Hans-Werner / Lannes, Jean
DOI: <https://doi.org/10.5169/seals-109907>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 07.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

38

EXOTIC CLASSES IN THE MOD 2 COHOMOLOGY OF $\mathrm{GL}_n(\mathbf{Z}[\frac{1}{2}])$

by Hans-Werner HENN and Jean LANNES

Let D_n denote the subgroup of the algebraic group GL_n consisting of diagonal matrices; let ι_n denote the inclusion of discrete groups $D_n(\mathbf{Z}[\frac{1}{2}]) \hookrightarrow \mathrm{GL}_n(\mathbf{Z}[\frac{1}{2}])$.

S. Mitchell [5] has shown that the image of the restriction map

$$\iota_n^*: H^*(\mathrm{GL}_n(\mathbf{Z}[\frac{1}{2}]); \mathbf{F}_2) \rightarrow H^*(D_n(\mathbf{Z}[\frac{1}{2}]); \mathbf{F}_2)$$

is isomorphic to $\mathbf{F}_2[w_1, w_2, \dots, w_n] \otimes \Lambda(e_1, e_3, \dots, e_{2n-1})$. Here the w_i are the *Stiefel–Whitney classes* of the tautological representation $\mathrm{GL}_n(\mathbf{Z}[\frac{1}{2}]) \rightarrow \mathrm{GL}_n(\mathbf{R})$ and the e_{2i-1} are closely related to Quillen’s odd-dimensional *modular characteristic classes* in the mod 2 cohomology of $\mathrm{GL}_n(\mathbf{F}_3)$.

By explicit calculation the map ι_n^* is known to be injective for $n \leq 3$ ([5] for $n = 2$ and [3] for $n = 3$). Work of Dwyer [1] shows that ι_n^* fails to be injective if $n \geq 32$ and still unpublished work of ours [4] — which will hopefully see the light of the day sometime soon — shows that it already fails if $n \geq 14$. However, the reasoning in [1] as well as in [4] is very indirect and does not produce any explicit elements in the kernel of ι_n^* . On the other hand one of us has shown [2] that the kernel of ι_n^* becomes very large as n grows.

PROBLEM 38.1. Construct explicit elements in the kernel of ι_n^* .

COMMENTS. (1) Injectivity of ι_n^* amounts to the validity of an unstable Lichtenbaum–Quillen conjecture for $\mathbf{Z}[\frac{1}{2}]$ at the prime 2 (cf. [1]) so any (even partial) progress on this problem would shed more light on why such an unstable conjecture fails.

(2) The solution of the Milnor conjecture by Voevodsky led to the proof that the map ι_∞^* (ι_∞ being the colimit of the ι_n) is injective: in other words, the stable Lichtenbaum-Quillen conjecture for $\mathbf{Z}[\frac{1}{2}]$ at the prime 2 holds.

(3) Now let us come back to the ghost reference [4]. Let O_n be the subgroup of GL_n consisting of orthogonal matrices (orthogonal for the euclidian metric) and ρ_n be the homomorphism of discrete groups $O_n(\mathbf{Z}[\frac{1}{2}]) \rightarrow O_n(\mathbf{F}_3)$ induced by mod 3 reduction. The reasoning in [4] is as follows: first we show that if ι_n^* is injective then ρ_n^* is bijective and afterwards we prove that ρ_n^* is not bijective for $n \geq 15$. (We use a variation of this argument to show that even ι_{14}^* is not injective.)

QUESTION 38.2. Is ρ_∞^* bijective?

A positive answer to this question amounts to the validity of an “orthogonal Lichtenbaum–Quillen conjecture for $\mathbf{Z}[\frac{1}{2}]$ at the prime 2”.

REFERENCES

- [1] DWYER, W. G. Exotic cohomology for $GL_n(\mathbf{Z}[1/2])$. *Proc. Amer. Math. Soc.* 126 (1998), 2159–2167.
- [2] HENN, H.-W. Commutative algebra of unstable K -modules, Lannes’ T -functor and equivariant mod- p cohomology. *J. Reine Angew. Math.* 478 (1996), 189–215.
- [3] ——— The cohomology of $SL(3, \mathbf{Z}[1/2])$. *K-Theory* 16 (1999), 299–359.
- [4] HENN, H.-W. and J. LANNES. (In preparation.)
- [5] MITCHELL, S. A. On the plus construction for $BGL(\mathbf{Z}[\frac{1}{2}])$. *Math. Z.* 209 (1992), 205–222.

Hans-Werner Henn

Département de Mathématique et
Inst. de Rech. Mathématique Avancée
Université Louis Pasteur
7, rue René Descartes
F-67084 Strasbourg Cedex
France
e-mail: henn@math.u-strasbg.fr

Jean Lannes

Centre de Mathématiques
Laurent Schwartz
École Polytechnique
F-91128 Palaiseau Cedex
France
e-mail: lannes@math.polytechnique.fr