

The Dehn function of $SL_n(\mathbb{Z})$

Autor(en): **Riley, Tim**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **23.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109924>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

55

THE DEHN FUNCTION OF $\mathrm{SL}_n(\mathbf{Z})$

by Tim RILEY

For a word w on $a_1^{\pm 1}, \dots, a_m^{\pm 1}$ representing 1 in a finite presentation $\mathcal{P} = \langle a_1, \dots, a_m \mid \mathcal{R} \rangle$ of a group Γ , define $\mathrm{Area}(w)$ to be the minimal $A \in \mathbf{N}$ such that there is an equality $w = \prod_{i=1}^A u_i^{-1} r_i^{\varepsilon_i} u_i$ in the free group $F(a_1, \dots, a_m)$ for some $\varepsilon_i = \pm 1$, some words u_i , and some $r_i \in \mathcal{R}$. Equivalently, $\mathrm{Area}(w)$ is the minimal A such that there is a van Kampen diagram for w over \mathcal{P} with at most A 2-cells. Defining $\mathrm{Area}(n)$ to be the maximum of $\mathrm{Area}(w)$ over all w that have length at most n and represent 1 in Γ , gives the *Dehn function* $\mathrm{Area}: \mathbf{N} \rightarrow \mathbf{N}$ of \mathcal{P} . Whilst $\mathrm{Area}: \mathbf{N} \rightarrow \mathbf{N}$ is defined for \mathcal{P} , a different finite presentation \mathcal{P}' for Γ will yield a Dehn function $\mathrm{Area}': \mathbf{N} \rightarrow \mathbf{N}$ that is qualitatively the same — for example, $(\exists C > 1, \forall n, (1/C)n^2 \leq \mathrm{Area}'(n) \leq Cn^2)$ if and only if the same is true for $\mathrm{Area}: \mathbf{N} \rightarrow \mathbf{N}$. (The C may differ.)

QUESTION 55.1. *Is the Dehn function of $\mathrm{SL}_n(\mathbf{Z})$ quadratic when $n \geq 4$?*

Presenting this as a question, rather than a claim, conjecture, or the like, may be unduly conservative. In his 1993 survey article [5], Gersten describes the quadratic Dehn function as an *assertion* of W.P. Thurston.

I am not even aware of a proof that the Dehn function of $\mathrm{SL}_n(\mathbf{Z})$ is bounded above by a polynomial when $n \geq 4$. By contrast, the Dehn function of $\mathrm{SL}_2(\mathbf{Z})$ is known to grow linearly — $\mathrm{SL}_2(\mathbf{Z})$ is hyperbolic — and that of $\mathrm{SL}_3(\mathbf{Z})$ grows like $n \mapsto \exp(n)$: Epstein–Thurston [4] proved the lower bound and a result sketched by Gromov [6] §5A₇ gives the upper bound. (An elementary proof might be a step towards 55.1.)

Of course, 55.1 presupposes $\mathrm{SL}_n(\mathbf{Z})$ is finite presentable, but that has been long known. The $n^2 - n$ matrices e_{ij} with 1's on the diagonal, the off-diagonal ij -entry 1, and all others 0, generate $\mathrm{SL}_n(\mathbf{Z})$. Milnor [11],

following J.R. Sylvester and in turn Nielsen and Magnus, explains that the Steinberg relations $\{[e_{ij}, e_{kl}] = 1\}_{i \neq l, j \neq k}$ and $\{[e_{jk}, e_{kl}] = e_{jl}\}_{j \neq l}$ together with $\{(e_{ij}e_{ji}^{-1}e_{ij})^4 = 1\}_{i \neq j}$ are defining relations. A proof of 55.1 would be an exacting quantitative proof of finite presentability.

One can regard 55.1 as a higher dimensional version of the Lubotzky–Mozes–Raghunathan Theorem ([9], [10]) establishing the existence of efficient words representing elements g of $\mathrm{SL}_n(\mathbf{Z})$ for $n \geq 3$, that is, words of length like the log of the maximum of the absolute values of the matrix entries. As a word representing g amounts to a path in the Cayley graph from 1 to g , the L.–M.–R. Theorem can be thought of as saying that 0-spheres admit efficient fillings by 1-discs. A word w representing 1 in a finite presentation \mathcal{P} corresponds to a loop ρ_w in the Cayley graph; a van Kampen diagram for w can be regarded as a combinatorial homotopy disc for ρ_w in the Cayley 2-complex of \mathcal{P} . So 55.1 is, roughly speaking, the claim that 1-spheres admit efficient fillings by 2-discs in $\mathrm{SL}_n(\mathbf{Z})$ for $n \geq 4$.

Gromov [6], §5D(5)(c), cf. §2B₁, takes this further and suggests that in $\mathrm{SL}_n(\mathbf{Z})$, Euclidean isoperimetric inequalities concerning filling k -spheres by $(k+1)$ -discs persist up to $k = n - 3$. (For $k = n - 2$, the exponential lower bound of [4] applies.)

One attack on 55.1 is that whilst $\mathrm{SL}_n(\mathbf{Z})$ is not a *cocompact* lattice in the symmetric space $X := \mathrm{SL}_n(\mathbf{R})/\mathrm{SO}(n)$, and so the quadratic isoperimetric inequality enjoyed by X does not immediately pass to $\mathrm{SL}_n(\mathbf{Z})$, open horoballs can be removed from X to give a space X_0 on which $\mathrm{SL}_n(\mathbf{Z})$ acts cocompactly. Druţu [2] and Leuzinger–Pittet [8] have made progress in this direction, including a quadratic isoperimetric inequality for the boundary horosphere of each removed horoball.

Chatterji has asked whether for $n \geq 4$, $\mathrm{SL}_n(\mathbf{Z})$ enjoys her property L_δ for some $\delta \geq 0$, which would imply a sub-cubic Dehn function [3].

The author's efforts towards 55.1 have, to date, yielded [12] a version of L.–M.–R. giving explicit efficient words. This may aid the construction of van Kampen diagrams, but that remains to be seen. However it has led to progress elsewhere [7].

Finally, we mention that for $n > 3$, the Dehn functions of the cousins $\mathrm{Aut}(F_n)$ and $\mathrm{Out}(F_n)$ of $\mathrm{SL}_n(\mathbf{Z})$ are also unknown [1].

REFERENCES

- [1] BRIDSON, M.R. and K. VOGTMANN. Automorphism groups of free groups, surface groups and free abelian groups. In: *Problems on Mapping Class Groups and Related Topics*, (edited by B. Farb), 301–316. Proc. Sympos. Pure Math. 74. Amer. Math. Soc., 2006.
- [2] DRUȚU, C. Filling in solvable groups and in lattices in semisimple groups. *Topology* 43 (2004), 983–1033.
- [3] ELDER, M. L_δ groups are almost convex and have a sub-cubic Dehn function. *Algebr. Geom. Topol.* 4 (2004), 23–29.
- [4] EPSTEIN, D.B.A., J.W. CANNON, D.F. HOLT, S.V.F. LEVY, M.S. PATERSON and W.P. THURSTON. *Word Processing in Groups*. Jones and Bartlett Publishers, Boston, London, 1992.
- [5] GERSTEN, S.M. Isoperimetric and isodiametric functions of finite presentations. In: *Geometric Group Theory, Vol. 1 (Sussex, 1991)*, 79–96. London Math. Soc. Lecture Note Ser. 181. Cambridge Univ. Press, Cambridge, 1993.
- [6] GROMOV, M. Asymptotic invariants of infinite groups. In: *Geometric Group Theory, Vol. 2 (Sussex, 1991)*, 1–295. London Math. Soc. Lecture Note Ser. 182. Cambridge Univ. Press, Cambridge, 1993.
- [7] KASSABOV, M. and T.R. RILEY. Diameters of Cayley graphs of Chevalley groups. *European J. Combin.* 28 (2007), 791–800.
- [8] LEUZINGER, E. and CH. PITTET. On quadratic Dehn functions. *Math. Z.* 248 (2004), 725–755.
- [9] LUBOTZKY, A., S. MOZES and M.S. RAGHUNATHAN. Cyclic subgroups of exponential growth and metrics on discrete groups. *C. R. Acad. Sci. Paris Sér. I Math.* 317 (1993), 735–740.
- [10] LUBOTZKY, A., S. MOZES and M.S. RAGHUNATHAN. The word and Riemannian metrics on lattices of semisimple groups. *Publ. Math. Inst. Hautes Études Sci.* 91 (2000), 5–53.
- [11] MILNOR, J. *Introduction to Algebraic K-Theory*. Annals of Mathematics Studies 72. Princeton University Press, Princeton, 1971.
- [12] RILEY, T.R. Navigating in the Cayley graphs of $SL_N(\mathbf{Z})$ and $SL_N(\mathbf{F}_p)$. *Geom. Dedicata* 113 (2005), 215–229.

T. Riley

Department of Mathematics
 University Walk
 Bristol
 BS8 1TW
 United Kingdom
e-mail: tim.riley@bris.ac.uk