

Zeitschrift: L'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: The regularity conjecture in the cohomology of groups
Autor: Benson, Dave
DOI: <https://doi.org/10.5169/seals-109879>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 07.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

10

THE REGULARITY CONJECTURE IN THE COHOMOLOGY OF GROUPS

by Dave BENSON

Let k be a field of characteristic p and let G be a finite group.

CONJECTURE 10.1. *The Castelnuovo–Mumford regularity of the cohomology ring is equal to zero:*

$$\text{Reg } H^*(G, k) = 0.$$

This conjecture was first announced at the opening workshop of the MSRI commutative algebra year ([1]), as a refinement of a conjecture of Benson and Carlson ([4]). Subsequent work on the conjecture was reported in [2] and [3].

We begin with the definitions. Let H be a finitely generated graded commutative k -algebra, with $H^0 = k$ and $H^i = 0$ for $i < 0$ (e.g., $H = H^*(G, k)$). Write \mathfrak{m} for the maximal ideal generated by the elements of positive degree. If M is a graded H -module then the local cohomology is doubly graded: $H_{\mathfrak{m}}^{i,j} M$ denotes the part in local cohomological degree i and internal degree j . Local cohomology can either be regarded as the right derived functors of the \mathfrak{m} -torsion functor $\Gamma_{\mathfrak{m}}(M) = \{x \in M \mid \exists n \geq 0, \mathfrak{m}^n \cdot x = 0\}$, or as the cohomology of the stable Koszul complex (see for example Theorem 3.5.6 of Bruns and Herzog [6]).

The a -invariants of M are defined to be

$$a_{\mathfrak{m}}^i(M) = \max\{j \in \mathbf{Z} \mid H_{\mathfrak{m}}^{i,j} M \neq 0\}$$

(or $a_{\mathfrak{m}}^i(M) = -\infty$ if $H_{\mathfrak{m}}^i M = 0$).

The *Castelnuovo–Mumford regularity* of M is then defined as

$$\text{Reg } M = \max_{i \geq 0} \{a_{\mathfrak{m}}^i(M) + i\}.$$

Of particular interest is the regularity of the ring itself, $\text{Reg } H$.

The reason for the interest in local cohomology of group cohomology comes from the Greenlees version ([7]) of Benson–Carlson duality ([4]), in the form of a spectral sequence

$$H_m^{i,j} H^*(G, k) \Rightarrow H_{-i-j}(G, k).$$

In particular, the existence of the “last survivor” of [4] shows the following ([2]):

THEOREM 10.2. $\text{Reg } H^*(G, k) \geq 0$.

The regularity conjecture is known to hold in the following situations:

- $H^*(G, k)$ is Cohen–Macaulay; e.g., groups with abelian Sylow p -subgroups; groups with extraspecial Sylow 2-subgroups with $p = 2$; groups of Lie type with characteristic coprime to p ([1]).
- Krull dimension minus depth at most two; e.g., 2-groups of order ≤ 64 ([2]).
- Symmetric and alternating groups in any characteristic; these are examples where Krull dimension minus depth is arbitrarily large ([3]).

There is also a corresponding conjecture for compact Lie groups. Let G be a compact Lie group of dimension d , and suppose that the adjoint action of G on $\text{Lie}(G)$ preserves orientation. Then there is a spectral sequence (Benson–Greenlees [5])

$$H_m^{i,j} H^*(BG; k) \Rightarrow H_{-i-j-d}(BG; k).$$

CONJECTURE 10.3. $\text{Reg } H^*(BG; k) = -d$.

To explain the orientation condition, let $G = T^3 \rtimes \mathbf{Z}/2$, a semidirect product of a 3-torus by an involution acting through inversion, and k be a field of characteristic $\neq 2$. Then

$$H^*(BG; k) = H^*(BT; k)^{\mathbf{Z}/2}$$

is Cohen–Macaulay but not Gorenstein, and $\text{Reg } H^*(BG; k) = -5$. The appropriate modification in this situation is that if ε denotes the orientation character, then $\text{Reg } H^*(BG; \varepsilon) = -d$.

ADDED IN PROOF. David Green has verified the regularity conjecture for all groups of order 128. See D. J. GREEN. ‘Testing Benson’s regularity conjecture’. Preprint arXiv : math.GR/0710.2311 (2007).

REFERENCES

- [1] BENSON, D. J. Commutative algebra in the cohomology of groups. In: *Trends in Commutative Algebra*. MSRI Publications, vol. 51, Cambridge Univ. Press, 2004, 1–50.
- [2] ——— Dickson invariants, regularity and computation in group cohomology. *Illinois J. Math.* 48 (2004), 171–197.
- [3] ——— On the regularity conjecture for the cohomology of finite groups. (Preprint, 2005.) To appear in *Proc. Edinb. Math. Soc.*
- [4] BENSON, D. J. and J. F. CARLSON. Projective resolutions and Poincaré duality complexes. *Trans. Amer. Math. Soc.* 342 (1994), 447–488.
- [5] BENSON, D. J. and J. P. C. GREENLEES. Commutative algebra for cohomology rings of classifying spaces of compact Lie groups. *J. Pure Appl. Algebra* 122 (1997), 41–53.
- [6] BRUNS, W. and J. HERZOG. *Cohen–Macaulay Rings*. Cambridge Studies in Advanced Mathematics 39, Cambridge Univ. Press, Cambridge, 1993.
- [7] GREENLEES, J. P. C. Commutative algebra in group cohomology. *J. Pure Appl. Algebra* 98 (1995), 151–162.

Dave Benson

King’s College
Aberdeen AB24 3UE
Scotland
United Kingdom
e-mail: bensondj@maths.abdn.ac.uk