

Zeitschrift: L'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Periodic p-torsion in the Farrell cohomology of some symplectic groups
Autor: Busch, Cornelia Minette
DOI: <https://doi.org/10.5169/seals-109888>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 07.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

19

PERIODIC p -TORSION IN THE FARRELL COHOMOLOGY OF SOME SYMPLECTIC GROUPS

by Cornelia Minette BUSCH

For an odd prime p and a nonzero integer $0 \neq n \in \mathbf{Z}$ we consider $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$, the group of symplectic matrices with coefficients in $\mathbf{Z}[1/n]$. It is defined to be

$$\mathrm{Sp}(p-1, \mathbf{Z}[1/n]) := \left\{ Y \in \mathrm{GL}(p-1, \mathbf{Z}[1/n]) \mid Y^t J Y = J := \begin{pmatrix} 0 & \mathbf{1} \\ -\mathbf{1} & 0 \end{pmatrix} \right\},$$

where $\mathbf{1}$ denotes the identity on $\mathbf{Z}[1/n]^{(p-1)/2}$. This symplectic group has finite virtual cohomological dimension and contains elements of order p . Moreover $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$ has p -periodic Farrell cohomology since each of its elementary abelian p -subgroups has rank ≤ 1 . Let

$$\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})_{(p)}$$

denote the p -primary part of the Farrell cohomology of $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$ with coefficients in \mathbf{Z} .

QUESTION 19.1. *What is the p -period of the Farrell cohomology ring*

$$\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})?$$

Here p is an odd prime and $0 \neq n \in \mathbf{Z}$ is any nonzero integer.

Since the group we are considering has the properties given above we get, by a result of K. S. Brown [1], the isomorphism

$$\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})_{(p)} \cong \prod_{P \in \mathfrak{P}} \widehat{H}^*(N(P), \mathbf{Z})_{(p)}.$$

Here \mathfrak{P} is a set of representatives of conjugacy classes of subgroups P of order p in $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$ and $N(P)$ denotes the normalizer of P .

In order to use this isomorphism, we analyze the structure of the subgroups of order p in $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$. We see in [3] that the conjugacy classes of elements of order p in $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$ are related to some ideal classes in $\mathbf{Z}[1/n][\xi]$, where ξ is a primitive p th root of unity. Therefore, if $n = 1$, the number of conjugacy classes of elements of order p in $\mathrm{Sp}(p-1, \mathbf{Z})$ depends on h^- , the relative class number of p . In [2] we get the following result.

THEOREM 19.2. *Let p be an odd prime for which the relative class number h^- is odd and let y be such that $p-1 = 2^r y$ with y odd. Then the period of $\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}), \mathbf{Z})_{(p)}$ equals $2y$.*

The smallest prime p for which h^- is even is $p = 29$. In fact it is not known if the statement of Theorem 19.2 is true for primes with even relative class number. Since the number of ideal classes in $\mathbf{Z}[\xi]$ is finite, it is possible to choose $0 \neq n \in \mathbf{Z}$ such that $\mathbf{Z}[1/n][\xi + \xi^{-1}]$ and $\mathbf{Z}[1/n][\xi]$ are principal ideal domains and moreover the odd prime p divides n . With these assumptions, we get the following result [4].

THEOREM 19.3. *Choose p and n as above. Let y be the greatest odd divisor of $p-1$. Then $2y$ is the p -period of the Farrell cohomology ring $\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})$ and, moreover, for any $i \in \mathbf{Z}$ we get an isomorphism*

$$\widehat{H}^i(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z}) \cong \widehat{H}^{i+d}(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})$$

with $d = y$ if and only if for each $j \mid y$ a prime $q \mid n$ exists with inertia degree f_q such that $j \mid \frac{p-1}{2f_q}$, and with $d = 2y$ if for some j no such q exists.

Here f_q is the inertia degree of the prime q in $\mathbf{Z}[\xi]$. It is the order of q in the group of units of \mathbf{F}_p . It is a consequence of Dirichlet's theorem on primes in arithmetic progression that for every $f_q \mid p-1$ an infinite number of primes q exist with inertia degree f_q . Let us guess the answer to Question 19.1.

CONJECTURE 19.4. *Let p be an odd prime. Then the p -period of the Farrell cohomology ring $\widehat{H}^*(\mathrm{Sp}(p-1, \mathbf{Z}[1/n]), \mathbf{Z})$ equals $2y$ for all $n \neq 0$, where y is the greatest odd divisor of $p-1$.*

REFERENCES

- [1] BROWN, K.S. *Cohomology of Groups*. Graduate Texts in Mathematics 87. Springer, 1982.
- [2] BUSCH, C.M. The Farrell cohomology of $\mathrm{Sp}(p-1, \mathbf{Z})$. *Doc. Math.* 7 (2002), 239–254.
- [3] ——— Conjugacy classes of p -torsion in symplectic groups over S -integers. *New York J. Math.* 12 (2006), 169–182.
- [4] ——— Isomorphisms in the Farrell cohomology of $\mathrm{Sp}(p-1, \mathbf{Z}[1/n])$. To appear in *New York J. Math.* 14 (2008).

Cornelia Minette Busch

Katholische Universität Eichstätt-Ingolstadt
Mathematisch-Geographische Fakultät
D-85071 Eichstätt
Germany
e-mail: cornelia.busch@ku-eichstaett.de