

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes Rübel in Zürich
Herausgeber: Geobotanisches Institut Rübel (Zürich)
Band: 8 (1932)

Artikel: The beech forests of Czechoslovakia
Autor: Domin, Karel
Kapitel: VIII: Seasonal aspect of the ground vegetation : phenological spectrum
DOI: <https://doi.org/10.5169/seals-307034>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 12.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

VIII. Seasonal aspect of the ground vegetation (phenological spectrum).

The aspect of the undergrowth of the beech forest changes rather considerably during every vegetative period. Some of the early spring species as *Isopyrum thalictroides*, *Galanthus nivalis*, *Corydalis*, *Adoxa*, *Scilla bifolia* soon disappear completely while others are developed fully later in the summer. *Epipogon aphyllus* has also only a short life-duration.

As an example, I quote the three aspects distinguished by MIKYŠKA (1) in the undergrowth of the beech forests and of the oak forests mixed with beech in the Štiavnické Středohoří (Slovakia).

Early spring aspect (species flowering to about the middle of May): *Anemone ranunculoides*, *Corydalis cava*, *C. digitata*, *Daphne mezereum*, *Chrysosplenium alternifolium*, *Isopyrum thalictroides*, *Petasites albus* and *Scilla bifolia*. The optimal floral development falls into the period before the new leaf-canopy closes.

Late spring aspect (lasting to the latter half or eventually to the end of June): *Actaea spicata*, *Alliaria officinalis*, *Asperula odorata*, *Carex pilosa*, *Dentaria bulbifera*, *Euphorbia amygdaloides*, *Glechoma hirsuta*, *Melica nutans*, *M. uniflora*, *Melitis melisophyllum*, *Oxalis acetosella*, *Poa nemoralis*, *Polygonatum officinale*, *P. multiflorum*, *Primula elatior*, *Ranunculus auricomus*, *R. lanuginosus*, *Stellaria holostea*, *Symphytum tuberosum*, *Veronica chamaedrys*, *Viola Riviniana* and *V. silvatica*.

Summer aspect (less distinctive): to the species with a longer flowering period belong the following: *Ajuga reptans*, *Asperula odorata*, *Galium Schultesii*, *Geranium Robertianum*, *Hieracium murorum*, *Myosotis silvatica*, *Viola Riviniana* also *silvatica*. Species flowering only during the summer are not numerous: *Astrantia major*, *Campanula trachelium*, *Epilobium montanum*, *Galeopsis pubescens*, *Hypericum hirsutum*, *Chrysanthemum corymbosum*, *Impatiens noli tangere*, *Knautia silvatica*, *Lactuca muralis*, *Lilium martagon*, *Melampyrum nemorosum*, *Phyteuma spicatum*, *Prenanthes purpurea*, *Sanicula europaea*, *Scrophularia nodosa*, *Senecio Fuchsii*, *Stachys silvatica* and *Valeriana sambucifolia*.

I restrict myself to this one example, because it is impossible to

present phenological spectra of every sociation and every type, not to mention the fact that the spectrum varies according to the exposure, altitude, as well as to climatic regions, etc.

IX. Shrub vegetation below canopy.

There are not, always, many strata in beech forests. The moss covering is nearly always lacking, the herbaceous undergrowth (in one or two layers) is more or less well developed but can also be suppressed (see *Fagetum nudum*), and the shrubby growth (young trees and true shrubs) is very unequally developed. As the most characteristic shrubs accompanying the beech, the following can be mentioned:

<i>Cornus sanguinea</i>	<i>Rosa pendulina</i>
<i>Corylus avellana</i>	<i>Rubus idaeus</i>
<i>Daphne mezereum</i>	<i>Sambucus racemosa</i>
<i>Lonicera nigra</i>	<i>Sorbus aria</i> (chiefly on limestone)
<i>Lonicera xylosteum</i>	<i>Sorbus torminalis</i>
<i>Ribes alpinum</i>	<i>Spiraea media</i> (only in Subcarpa-
<i>Ribes grossularia</i>	thian Russia)

Rather characteristic for some beech forests are also *Evonymus verrucosa*, *Ligustrum vulgare* (ab.), *Stachylea pinnata* and *Viburnum opulus*. Besides, a good many other shrubs (for instance *Cornus mas*, *Crataegus*, *Cotoneaster tomentosa* (Carpathians only), *Berberis vulgaris*, *Rhamnus cathartica*, *Viburnum lantana* and some *Rosa* and *Rubus* species are sometimes present in some beech forest sociations.

X. Ground vegetation.

The ground vegetation is the most reliable basis for a sociological classification of beech forests, because the general tree stratum is uniform and the small number of accompanying trees cannot be depended upon for establishing definite sociations. Since a sociological classification of beech forests is exceedingly difficult, many authors avoid a definite evaluation and distinguish simply «types», often characterised also ecologically. These types, however, are not identical with the well-known C a j a n d e r's forest types, because these authors interpret the beech forest, including its tree