

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübel, in Zürich
Herausgeber: Geobotanisches Institut, Stiftung Rübel (Zürich)
Band: 95 (1987)

Artikel: Biosystematic investigations in the family of duckweeds ("Lemnaceae"). Vol. 4 : The family of "Lemnaceae" : a monographic study. Volume 2
Autor: Landolt, Elias / Kandeler, Riklef
Register: List of the figures
DOI: <https://doi.org/10.5169/seals-308870>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 30.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LIST OF THE FIGURES

Fig. 1. 1. The N and P content of <u>Lemna gibba</u> in relation to the N and P content of the nutrient solution	14
Fig. 1. 2. The Cd, Cr, Cu, Pb, and Zn content of <u>Spirodela polyrrhiza</u> in relation to the concentration of the metals in the nutrient solution	15
Fig. 1. 3. Banding patterns for 6 different enzyme systems of 11 <u>Lemnaceae</u> species	54
Fig. 2. 1. Light response curves for photosynthetic rates of <u>Lemna gibba</u> measured in air and in CO ₂ enriched air at 5000 ppm CO ₂	72
Fig. 2. 2. Growth rate of <u>Lemna gibba</u> and <u>Wolffia arrhiza</u> in relation to the concentration of Hutner solution	80
Fig. 2. 3. The effect of the concentrations of potassium, calcium, magnesium, nitrate, phosphate, and sulfate on the growth rate of <u>Spirodela polyrrhiza</u>	88
Fig. 2. 4. The effect of the concentrations of iron, manganese, zinc, and chloride on the growth rate of <u>Spirodela polyrrhiza</u>	89
Fig. 2. 5. Requirement and tolerance of <u>Spirodela intermedia</u> for Ca and Mg	94
Fig. 2. 6. Requirement and tolerance of <u>Wolffia angusta</u> for Ca and Mg	95
Fig. 2. 7. Relation between growth rate and nitrogen supply in <u>Lemna gibba</u>	100
Fig. 2. 8. Depletion of ammonium and nitrate in the medium by <u>Lemna gibba</u> adapted to a relative nitrogen addition rate of 0.25 per day	107
Fig. 2. 9. The relative distribution of dissolved inorganic carbon among its various chemical forms as a function of pH	119
Fig. 2.10. Growth rate of <u>Spirodela polyrrhiza</u> in relation to sucrose concentration	136
Fig. 2.11. Growth rate of <u>Spirodela polyrrhiza</u> in the dark with 1% sucrose, at 1000 lux without sucrose and at 1000 lux with 1% sucrose	137
Fig. 2.12. The effect of EDTA on the growth of <u>Spirodela polyrrhiza</u>	143
Fig. 2.13. Growth rates of <u>Lemna minor</u> at different temperatures and different light intensities	176
Fig. 2.14. Growth rate of different <u>Lemnaceae</u> species in relation to temperature	177
Fig. 2.15. Mean dry weight per frond of <u>Lemna minor</u> at different temperatures and different light intensities	178
Fig. 2.16. Mean net assimilation rates of <u>Lemna minor</u> plotted against temperature	179
Fig. 2.17. Relation between growth rate and assimilation rate of <u>Lemna minor</u> at different temperatures	180
Fig. 2.18. Turion formation of <u>Spirodela polyrrhiza</u> and <u>Lemna turionifera</u> in relation to temperature	181
Fig. 2.19. Germination of turions in relation to temperature	182
Fig. 2.20. Relation between light intensity and growth rate in <u>Wolffia arrhiza</u>	189
Fig. 2.21. Growth rates of <u>Lemna minor</u> at different light intensities and different temperatures	190
Fig. 2.22. Growth rate of different species in relation to light intensity	191

Fig. 2.23. Mean dry weight per frond of <u>Lemna minor</u> at different light intensities and different temperatures	192
Fig. 2.24. Dry weight per frond in relation to light intensity	193
Fig. 2.25. Growth rate of <u>Lemna gibba</u> at three light intensities in solutions with and without sucrose in relation to temperature	195
Fig. 2.26. Growth rate of <u>Lemna gibba</u> and in relation to the length of day in solution with sucrose and without sucrose	198
Fig. 2.27. Growth rate of <u>Wolffia arrhiza</u> in relation to the length of day in solution with sucrose and without sucrose	199
Fig. 2.28. The effect of red light and kinetin on the development of intercellular air spaces in <u>Lemna minor</u>	213
Fig. 2.29. The vascular system in fronds of <u>Lemna minor</u> as influenced by four concentrations of triiodobenzoic acid	214
Fig. 2.30. The length of life of <u>Lemna minor</u> fronds, the number of daughter fronds produced during the life-span of one mother frond, and the length of daughter fronds as influenced by the N- and P- content of nutrient medium	221
Fig. 2.31. Senescence and frond development in <u>Lemna aequinoctialis</u>	227
Fig. 2.32. Arrangement of chloroplasts in the same cells induced by polarized light	233
Fig. 2.33. Effect of abscisic acid on growth and turion formation of <u>Spirodela polyrrhiza</u>	237
Fig. 2.34. The contrary influence of mother frond senescence on flower induction in daughter fronds of <u>Lemna gibba</u> Gl and <u>L. aequinoctialis</u> 6746	249
Fig. 2.35. Movability of stomata in <u>Lemnaceae</u>	271
Fig. 2.36. Scheme of the xanthophyll cycle	277
Fig. 2.37. Photorespiration in <u>Lemna minor</u> as shown by the course of CO ₂ -evolution during repeated light-dark changes (20:20 min) under pure oxygen	280
Fig. 2.38. ATP content of intact <u>Lemna aequinoctialis</u> plants as a function of respiration rate	281
Fig. 2.39. Transmembrane potential changes in single subepidermal cells of <u>Lemna aequinoctialis</u> fronds induced by short red and far-red light pulses	287
Fig. 2.40. Velocity of K ⁺ -absorption plotted against K ⁺ -concentration (<u>Lemna minor</u>)	288
Fig. 2.41. Accumulation of released ABA in the medium of a <u>Spirodela polyrrhiza</u> culture	296
Fig. 2.42. The effects of continuous light, DCMU, and sucrose on malate accumulation in <u>Lemna aequinoctialis</u> 6746	305
Fig. 2.43. Biosynthesis of myo-inositol, apiose, and xylose in <u>Lemna</u>	309
Fig. 2.44. Location of phosphatase in control and phosphorus-deficient <u>Spirodela punctata</u> plants	313
Fig. 2.45. The glutamine pathway of ammonium assimilation	317
Fig. 2.46. Summary of the free and bound pathways of sulfate assimilation in plants	321
Fig. 2.47. Methionine biosynthesis in <u>Lemna</u>	325
Fig. 2.48. Biosynthetic pathway of branched-chain amino acids	329
Fig. 2.49. Methionine metabolism	330
Fig. 2.50. Possible secondary structure of <u>Lemna minor</u> chloroplast 5 S rRNA	341
Fig. 2.51. Restriction fragment map of <u>Spirodela punctata</u> chloroplast DNA	345
Fig. 2.52. (4R)-4-hydroxyisophytol, a novel diterpene isolated from <u>Lemna minor</u>	350

Fig. 2.53. Flavonoid nucleus, numbering system, and substitutions of the flavone, glycoflavone, and flavonols and of the anthocyanidins detected in <u>Lemnaceae</u>	354
Fig. 2.54. Metabolic pathways leading to several subgroups of flavonoids in <u>Lemnaceae</u>	355
Fig. 2.55. Intermediate steps in the light action on anthocyanin biosynthesis in <u>Spirodela punctata</u>	358
Fig. 2.56. Hypothetical scheme for a self-sustaining and light- entrainable rhythm working in <u>Lemna gibba</u> G3	364
Fig. 3. 1. Activity of some enzymes of <u>Wolffia arrhiza</u>	404