

Zeitschrift: Veröffentlichungen des Geobotanischen Institutes der Eidg. Tech. Hochschule, Stiftung Rübél, in Zürich

Herausgeber: Geobotanisches Institut, Stiftung Rübél (Zürich)

Band: 95 (1987)

Artikel: Biosystematic investigations in the family of duckweeds ("Lemnaceae"). Vol. 4 : The family of "Lemnaceae" : a monographic study. Volume 2

Autor: Landolt, Elias / Kandeler, Riklef

Register: List of the tables

DOI: <https://doi.org/10.5169/seals-308870>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 06.02.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LIST OF THE TABLES

| | |
|---|-----|
| Table 1. 1. Variation of content of elements in % of the dry weight | 10 |
| Table 1. 2. Mineral ratios in <u>Lemnaceae</u> , according to many authors | 11 |
| Table 1. 3. Variation of chemical composition of <u>Wolffia globosa</u> from Japan between summer and winter | 17 |
| Table 1. 4. Differences in mineral content between <u>Spirodela polyrrhiza</u> , <u>Lemna minor</u> , and <u>L. trisulca</u> | 17 |
| Table 1.5a. Mineral content of three <u>Lemna</u> species cultivated under identical conditions | 17 |
| Table 1.5b. Differences of minimum and maximum content of N and P between different species | 18 |
| Table 1. 6. Mineral content of <u>Lemnaceae</u> species grown at concentrations of minerals to accomplish half of the maximum growth rate | 18 |
| Table 1. 7. Variations of organic components in <u>Lemnaceae</u> | 20 |
| Table 1. 8. Amino acid composition of the proteins of <u>Lemnaceae</u> | 24 |
| Table 1. 9. Carbohydrate content in <u>Lemnaceae</u> | 26 |
| Table 1.10. Percentage of sugars in <u>Lemnaceae</u> | 26 |
| Table 1.11. Content of different sugars within the 3 carbohydrate groups of pectins, hemicelluloses and celluloses in <u>Lemna gibba</u> | 27 |
| Table 1.12. Chlorophyll and carotinoid content in <u>Lemnaceae</u> | 32 |
| Table 1.13. Flavonoids of <u>Lemnaceae</u> | 37 |
| Table 1.14. Content of lignin components in <u>Lemnaceae</u> | 41 |
| Table 1.15. Relative chemical composition of raphide idioblasts in <u>Lemna minor</u> | 43 |
| Table 1.16. Enzymes determined in members of <u>Lemnaceae</u> | 48 |
| Table 1.17. Enzyme reactions to 7 different stainings of 11 <u>Lemnaceae</u> species | 53 |
| Table 1.18. Polypeptid composition of large and small subunits of Fraction-1-protein for different species of <u>Lemnaceae</u> | 55 |
| Table 1.19. Relative chemical composition of tannin idioblasts <u>Lemna minor</u> | 56 |
| Table 2. 1. Composition of some nutrient solutions used for <u>Lemnaceae</u> growth | 60 |
| Table 2. 2. Comparison of the molar nutrient content of 7 different nutrient solutions | 62 |
| Table 2. 3. CO ₂ uptake of <u>Lemna minor</u> at different light intensities and 5% CO ₂ | 73 |
| Table 2. 4. CO ₂ uptake of <u>Spirodela punctata</u> , <u>Lemna minor</u> and <u>L. valdiviana</u> at a light intensity of 240 E m ² and second and at different CO ₂ concentrations | 74 |
| Table 2. 5. Carbon uptake from water in <u>Spirodela polyrrhiza</u> in mg C per g dry weight and hour | 75 |
| Table 2. 6. Carbon uptake from water in <u>Spirodela polyrrhiza</u> | 76 |
| Table 2. 7. Comparison of growth rates of 15 species of <u>Lemnaceae</u> grown in Hutner solution 1/3 and 1/100 | 80 |
| Table 2. 8. Lower pH limits for growth of <u>Lemnaceae</u> | 84 |
| Table 2. 9. Nutrient requirements of <u>Spirodela polyrrhiza</u> | 86 |
| Table 2.10. The effect of low potassium content on characteristics of <u>Lemnaceae</u> | 88 |
| Table 2.11. Maximum and minimum requirement of Ca and Mg for long-lasting growth of <u>Lemnaceae</u> | 96 |
| Table 2.12. Nitrogen requirement and nitrogen tolerance of different species of <u>Lemnaceae</u> | 101 |

| | |
|---|-----|
| Table 2.13. The effect of low nitrogen supply on characteristics of <u>Lemnaceae</u> | 103 |
| Table 2.14. pH values of culture solutions with nitrate, ammonium-nitrate, and ammonium after cultivation of 14 and 40 days | 110 |
| Table 2.15. Phosphorus requirement and phosphorus tolerance for growth of some <u>Lemnaceae</u> species | 113 |
| Table 2.16. The effect of low phosphorus supply on characteristics of <u>Lemnaceae</u> | 115 |
| Table 2.17. Tolerance of <u>Lemna minor</u> towards different micro-nutrients | 123 |
| Table 2.18. Influence of cytokinins on growth rates of <u>Lemnaceae</u> | 151 |
| Table 2.19. Effects of some herbicides on <u>Lemnaceae</u> | 162 |
| Table 2.20. Effects of some algicides, fungicides, insecticides, and other toxic substances on <u>Lemnaceae</u> | 169 |
| Table 2.21. Growth rate of different species of <u>Lemnaceae</u> at two light intensities | 196 |
| Table 2.22. Grouping of different species and clones of <u>Lemnaceae</u> according to their behaviour towards yeast extracts and casamino acids in darkness | 207 |
| Table 2.23. Meaning of the categories A), B), C), and D) in table 2.22. | 208 |
| Table 2.24. Radio sensitivity of different <u>Lemnaceae</u> species | 209 |
| Table 2.25. Substances influencing flower initiation in <u>Lemnaceae</u> | 244 |
| Table 2.26. Effects of blue, red, and far-red irradiation on flowering in <u>Lemna gibba</u> (LDP) and <u>L. aequinoctialis</u> (SDP) | 262 |
| Table 2.27. Concentration of some ions in cell compartments of <u>Spirodela polyrrhiza</u> | 299 |
| Table 3. 1. Productivity of <u>Lemnaceae</u> in different regions of the world | 372 |
| Table 3. 2. Number of fronds of <u>Lemnaceae</u> per m ² water surface in nature | 373 |
| Table 3. 3. Energy content of different <u>Lemnaceae</u> | 374 |
| Table 3. 4. Content of essential amino acids of <u>Lemnaceae</u> , rice, soybean, <u>Chlorella</u> , and egg compared to FAO reference pattern | 376 |
| Table 3. 5. Content of protein, carbohydrate and fat in different crop plants from northern Thailand | 377 |
| Table 3. 6. Daily removal of N, P, and K by <u>Lemnaceae</u> during warm seasons | 391 |
| Table 3. 7. Accumulation factors for different elements in <u>Lemnaceae</u> | 396 |