

The low energy proton polarimeter for the polarized H⁻ beam source at the Brookhaven AGS

Autor(en): **Schüler, K.P. / Liu, C.J. / Gay, T.J.**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **59 (1986)**

Heft 4

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-115740>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THE LOW ENERGY PROTON POLARIMETER
 FOR THE POLARIZED H^- BEAM SOURCE AT THE BROOKHAVEN AGS*

K.P. Schüler, C.J. Liu and T.J. Gay†

Yale University, Physics Department
 New Haven, Connecticut 06511, USA

ABSTRACT

A proton polarimeter suitable for spin polarized H^+ and H^- ion beams of low energy (10 keV to 100 keV) has been developed and tested with the 20 keV polarized H^- beam source at the Brookhaven AGS. Following the production of neutral hydrogen atoms in the 2P state by collision with a thin carbon foil, the beam polarization is measured by detecting the circular polarization of the resultant Lyman- α decay photons.

A new type of proton polarimeter has been developed and tested with the 20 keV polarized H^- beam source [1] at the Brookhaven AGS. The instrument, which is depicted schematically in Fig. 1, is based on beam foil spectroscopy and involves the measurement of circular polarization of Lyman- α light. The method has previously been applied only to heavier nuclei with photon detection at longer wavelengths [2][3].

The principle of the method is the following. The polarized hydrogen ion beam passes at low energy (10 keV to 100 keV) through a thin carbon foil ($5\mu\text{g}/\text{cm}^2$). Any electrons associated with the projectiles are stripped upon entering the foil. Therefore it does not matter whether H^+ or H^- beams are being used. The foil is traversed within 10^{-14} s, which is short compared to all hyperfine periods. The proton spin orientation is therefore not affected by the passage through the foil. When the protons exit from the foil, a fraction of them will pick up an electron and emerge as neutral hydrogen atoms in the 2P state. As the excited atoms move away from the foil, a transfer of angular momentum (orientation) occurs via hyperfine interaction from the proton spin to the electron orbit, which will manifest itself through circular polarization of the Lyman- α photon decay. Maximum circular polarization will be




Fig. 1. Beam-Foil Proton Polarimeter observed for photons emitted along the

*Work supported in part by U.S. Department of Energy
 †Present address: Univ. of Missouri, Rolla, Physics Dept.,
 Rolla, Missouri 65401, USA

spin quantization axis, and the extent of the circular polarization will be a direct measure of the original proton polarization.

The circular polarization Stokes parameter can be expressed in the form $S/I = P \cdot A(t)$, where P is the beam polarization, and $A(z)$ or $A(t)$ is the analyzing power of the process at the distance z from the foil (or time of flight $t = z/v$), which is the degree of circular polarization obtained in the case of complete ($P=1$) beam polarization. Using the formalism developed by Ellis [4], we reach the following result

$$A(t) = (5/24) \cdot (1 - \cos \omega_{3/2} t) + (1/6) \cdot (1 - \cos \omega_{1/2} t) \quad (1)$$

for the Lyman- α decay of 2P states originating at the foil at $t=0$, where $\omega_{1/2} = 2\pi \cdot 59\text{MHz}$ and $\omega_{3/2} = 2\pi \cdot 24\text{MHz}$ are the hyperfine splittings of the $J=1/2$ and $J=3/2$ levels. A more exact version § of eq.(1), allowing for alignment of the 2P state [6], is only slightly different. Much more significant, however, is the effect of cascading from higher excited states. In this regard the 3D state, with a 3D \rightarrow 2P lifetime of 15.6 nsec, is of primary concern. We obtain for the total intensity $I = I_{2P} + I_{3D}$ and for the cascade modified analyzing power A the following expressions

$$I(t) = C \exp(-\gamma_2 t) \{1 + [B\gamma_3 / (\gamma_2 - \gamma_3)] \cdot [\exp(\gamma_2 - \gamma_3)t - 1]\} \quad (2)$$

$$A(t) = (C/I(t)) \cdot \{ \tilde{A}(t) \exp(-\gamma_2 t) + B\gamma_3 \int_0^t \tilde{A}(t-\alpha) \exp(-\gamma_3 \alpha) \exp(-\gamma_2(t-\alpha)) d\alpha \} \quad (3)$$

where C is a constant, $1/\gamma_2$ is the 2P lifetime (1.6ns), $1/\gamma_3$ is the 3D lifetime (15.6ns), the cascading parameter B is the initial 3D/2P population ratio at the foil ($t=0$), and the original analyzing power from eq. (1) appears now as $\tilde{A}(t)$ in eq.(3).

The theoretical Lyman- α photon intensity and analyzing power according to eq. (2) and eq. (3) are plotted in Fig. 2 for a cascading parameter $B=0.15$ that follows from measurements of Bukow et al.[7]. For a realistic comparison with measurements, the energy loss of the projectiles in the foil, the time resolution corresponding to the slit width and the geometrical acceptance of the optical polarimeter have to be considered.

The principal components of the instrument are shown in Fig.1. The carbon foil target can be moved along the beam axis. The UV-polarimeter has a 3 mm slit that samples a slice of the beam, followed by a rotatable MgF_2 waveplate [8] and a Brewster angle type four-mirror polarizer [8][9], both obtained from NASA, and a photomultiplier with CsI cathode and MgF_2 window. A second photomultiplier serves as intensity monitor. The circular analyzing power of the waveplate now used is $\sin \delta = 1.00 \pm 0.04$. This is a considerable improvement over the earliest operation of the instrument ($\sin \delta = 0.5 \pm 0.1$) reported at the Osaka Conference [10].

The beam-foil polarimeter has been tested with the 20 keV polarized H^- source PONI-1 at BNL [1][11], which delivered a pulsed beam of up to $5 \times 10^{10} \text{H}^-$ per 300 μs pulse at ~ 1 pps. Photons were counted in two sets of scalars, flipping either the waveplate ($\pm 45^\circ$) or the beam polarization (\uparrow, \downarrow).

$$\S \quad A(t) = [(10 + 2A_0^{\text{col}})(1 - \cos \omega_{3/2} t) + 8(1 - \cos \omega_{1/2} t)] / [48 - A_0^{\text{col}}(5 + 3 \cos \omega_{3/2} t)],$$

where A_0^{col} is the Fano-Macek alignment parameter (Ref.[5]); for H(2P) alignment data see Ref.[6].




Fig.2. Theoretical analyzing power $A(t)$ and Ly- α photon intensity $I(t)$. The curves include modifications due to cascading from the 3D state(15%) according to eq.(2) and eq.(3).

The observed dependence of the Lyman- α photon intensity and circular polarization on foil position (Fig. 3) are well described by the cascade modified theory. The measured asymmetries are now larger by more than a factor of two compared to earlier data reported at the Osaka Conference [10]. This is mainly due to the better optics in the polarimeter. There has also been some work on the rf transition regions in the polarized H^0 -stage of the source [11] that has resulted in a much improved weak field polarization. The magnitude of the polarization of the H^- beam, as determined from the size of our measured asymmetries, was:

(Weak Field Pol. :) $P \dagger = 0.81$
 (Strong Field Pol. :) $P \dagger = 0.67$

The statistical error is about ± 0.01 . The systematic error is estimated to be around ± 0.05 , which includes the uncertainties of the cascading factor, the speed of particles after the foil and the analyzing power of the optics.

Fig.4 and Fig.5 show some examples of the measured asymmetries as a function of various parameter settings in the source. It indicates that this polarimeter can be used as a tuning device to maximize beam polarization.





Fig.3. Measured circular polarization of Ly- α photons and intensity vs. foil position. The curves are fits to the functional forms given in Fig.2, with instrumental acceptance and resolution included, from which an average proton polarization of $P=0.74$ was deduced.

The analyzing power of this polarimeter will not depend on beam energy, because the cascading factor is believed to be reasonably constant in the energy range of interest [7]. This has been verified in a limited range between 10 keV and 20 keV. Counting rate considerations favor the energy range from 10 keV to 100 keV. At higher energies the 2P production cross section drops rapidly [12]. The beam-foil polarimeter is intrinsically simple and has the great advantage that it can operate at typical source energies without further acceleration.




Fig.4. Tuning of the high frequency (1.49 GHz) rf transition region in the polarized H^0 -stage of the source for maximum $P\uparrow$ polarization of the H^- beam. Correspondingly, there is a low-frequency (19.5 MHz) transition region for the $P\uparrow$ polarization.




Fig.5. Tuning of solenoids in the source for maximum proton polarization. The magnetic field in the interaction region decouples the spin of the electron and proton. The spin rotation solenoid rotates the proton spin into an up-right orientation.

REFERENCES

- [1] A. Kponou, J. Alessi, Th. Sluyters, Proceedings of the Particle Accelerator Conference in Vancouver (Canada), 1985, IEEE Transactions NS-32 (1985) 1764.
- [2] H.J. Andrä, H.J. Plöhn, A. Gaupp, R. Fröhling, Nuclear Spin Polarization Produced by Ion-Beam-Surface-Interaction, its Optical Detection and Use in an Atomic Quantum-Beat Experiment, Zeitschrift für Physik A281 (1977) 15.
- [3] W. Dreves, P. Egelhof, K.H. Möbius, E. Steffens, G. Tungate, R. Böttger, D. Fick, Production of a Polarized ^{23}Na Beam, Zeitschrift für Physik A288 (1978) 413.
- [4] D.G. Ellis, Optical Emission by Spin-Polarized Atoms, Journal of Physics B10 (1977) 2301.
- [5] U. Fano and J.H. Macek, Impact Excitation and Polarization of the Emitted Light, Reviews of Modern Physics 45 (1973) 553.
- [6] H. Winter: Alignment Studies in the Hydrogen 2P-State after Beam-Foil Excitation, Journal de Physique Colloque C1 (1979) C1-307.
- [7] H.H. Bukow, H.v. Buttlar, D. Haas, P.H. Heckmann, M. Holt, W. Schlagheck, D. Schürmann, R. Tielert, R. Woodruff, Lifetime and Initial Populations of Foil-Excited Hydrogen and Lithium States, Nuclear Instruments and Methods 110 (1973) 89.
- [8] J. Calvert, D. Griner, J. Montenegro, F. Nola, F. Rutledge, E. Tandberg-Hanssen, C.L. Wyman, J.M. Beckers, An Ultraviolet Polarimeter for the Solar Maximum Mission, Optical Engineering 18 (1979) 287;
B.E. Woodgate, E.A. Tandberg-Hanssen, E.C. Bruner, J.M. Beckers, J.C. Brandt, W. Henze, C.L. Hyder, M.W. Kalet, P.J. Kenny, E.D. Knox, A.G. Michalitsianos, R. Rehse, R.A. Shine, H.D. Tinsley, The Ultraviolet Spectrometer and Polarimeter on the Solar Maximum Mission, Solar Physics 65 (1980) 73;
M.S. Miller, A.J. Caruso, B.E. Woodgate, A.A. Sterk, Ultraviolet Spectrometer and Polarimeter for the Solar Maximum Mission, Applied Optics 20 (1981) 3805.
- [9] G. Hass and W.R. Hunter, Reflection Polarizers for the Vacuum Ultraviolet Using Al + MgF₂ Mirrors and an MgF₂ Plate, Applied Optics 17 (1978) 76.
- [10] K.P. Schüller, C.J. Liu and T.J. Gay, The Low Energy Proton Polarimeter for the Polarized H⁻ Beam Source at the Brookhaven AGS, Proceedings of the 6th Int. Symposium on Polarization Phenomena in Nuclear Physics, Osaka (Japan) 1985, to be published.
- [11] J. Alessi, A. Kponou and Th. Sluyters, Contribution to this Conference.
- [12] Y. Baudinet-Robinet and P.D. Dumont, Populations of 2p and 3p Terms in Hydrogen Excited by H⁺, H₂⁺ and H₃⁺ Ions Passing Through Thin Carbon Foils, Physical Review A29 (1984) 1825.