

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 62 (1989)  
**Heft:** 6-7

**Artikel:** Measurements of metal atom and cluster polarizabilities  
**Autor:** Milani, P. / Chatelain, A. / Heer, W. de  
**DOI:** <https://doi.org/10.5169/seals-116123>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 15.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**MEASUREMENTS OF METAL ATOM AND CLUSTER POLARIZABILITIES**

P. Milani, A. Chatelain, W. de Heer

Institut de Physique Expérimentale, EPF-Lausanne, 1015 Lausanne, Switzerland

**Abstract**

A new technique to measure the polarizabilities of atoms and small clusters of refractory metals in a molecular beam is described. Clusters are produced in a laser vaporization source and the collimated cluster beam is deflected in an inhomogeneous electric field. The clusters are then photoionized and detected by a recently developed position sensitive time of flight mass analyzer. The polarizabilities are calculated from the measured deflections. We have measured for the first time the polarizabilities of the aluminum atom and the aluminum dimer for which we find  $6.8 \pm 0.3 \cdot 10^{-24} \text{ cm}^3$  and  $19 \pm 2 \cdot 10^{-24} \text{ cm}^3$  respectively.

**Introduction**

Experimental values for the polarizabilities of clusters are of fundamental importance. For small sodium clusters the polarizability is approximately  $(R + \delta)^3$  (1), where  $R$  is the classical radius and  $\delta$  is the electronic spill out factor. For these clusters it is found that  $\delta$  is approximately constant for all sizes from the atom to the bulk. Since the static polarizability is a ground state property, it can be accurately calculated and is a good test of electronic structure calculations. However, because of difficulties in producing molecular beams of refractory materials, the atomic polarizabilities of relatively few elements have been measured.

**Experimental Method**

Clusters are produced in a laser vaporization source (1). The inert gas carried cluster beam is collimated and passes between the two electric field plates which produce a field gradient of about 1150 kV/cm<sup>2</sup> at a field of 220 kV/cm. Due to

this field the clusters are deflected on the order of 0.5 mm at the detector which is located 1m from the field plates. The total distance from source to detector is 2.4 m. The deflections are given by  $d = \alpha g V^2/mv^2$ , where  $\alpha$  is the polarizability,  $V$  the plate voltage,  $m$  the cluster mass,  $v$  the cluster velocity, and  $g$  is a geometrical factor.

In the detector the clusters are ionized with a UV light pulse from an eximer laser. The clusters ions are accelerated perpendicular to the molecular beam by uniform electric fields. These fields are adjusted so that the flight time of a cluster depends on its mass as well as on its position at the time of ionization. A shift of 1mm in the position of the cluster causes a shift of about 0.1% in its flight time. One can therefore determine the deflections of the clusters caused by the inhomogeneous electric field. The ratios of the deflections of the clusters are then directly proportional to the ratios of the polarizabilities.

## Results

From the deflections we find that the ratio of the polarizability of the aluminum atom and the lithium atom is 0.284 and the ratio of the aluminum dimer and the aluminum atom is 2.77. Using the measured value for the lithium atom we then find for the polarizabilities of the aluminum atom and the aluminum dimer  $6.8 \pm 0.3 \cdot 10^{-24} \text{ cm}^3$  and  $19 \pm 2 \cdot 10^{-24} \text{ cm}^3$  respectively. These values agree well with recent calculations by I.Moulet who finds 6.67 and  $17.39 \cdot 10^{-24} \text{ cm}^3$  (2). Measurements for larger clusters up to about Al 50, and of atoms of refractory metals are currently being performed.

## References

- 1) W.A.de Heer, W.D.Knight, M.Y.Chou, M.L.Cohen "Electronic Shell Structure and Metal Clusters" in Solid State Physics 40, Ehrenreich and Turnbull Eds. (Academic Press, 1987), and references therein.
- 2) I. Moulet, private communication (1989).