

Zeitschrift: Helvetica Physica Acta
Band: 62 (1989)
Heft: 6-7

Artikel: Binding energies of p-type acceptor states in GaAs/Ga_{1-x}Al_xAs quantum wells
Autor: Pasquarello, A. / Andreani, L.C. / Buczko, R.
DOI: <https://doi.org/10.5169/seals-116139>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

BINDING ENERGIES OF *p*-TYPE ACCEPTOR STATES IN GaAs/Ga_{1-x}Al_xAs QUANTUM WELLS

A. Pasquarello^(a,c), L. C. Andreani^(a,c), R. Buczko^(b,c)

(a) Institut de Physique Théorique, EPFL, Lausanne, Switzerland.

(b) Institute of Physics, Academy of Sciences, Warsaw, Poland.

(c) Scuola Normale Superiore, Pisa, Italia.

Abstract: Binding energies of *p*-type shallow acceptor states in GaAs/Ga_{1-x}Al_xAs quantum wells are calculated. The complex valence band structure has been taken into account in a four band effective mass theory. The acceptor envelope function is expanded in valence envelope functions in the two-dimensional *k*-space.

1. Theory and results

In this paper we develop an effective mass theory to describe excited shallow acceptor states in GaAs/Ga_{1-x}Al_xAs quantum wells. The acceptor Hamiltonian contains a kinetic part which is given by the Luttinger-Kohn Hamiltonian [1], a square well potential due to the valence band discontinuity, and the Coulomb potentials of the impurity charge and of all its image charges.

We expand the acceptor envelope function in the basis of valence envelope functions [2], and obtain

$$F^s(\rho, \theta, z) = \sum_n \int d\mathbf{k} G_n(\mathbf{k}) v_{n\mathbf{k}}^s(z) e^{i\mathbf{k} \cdot \boldsymbol{\rho}}, \quad (1)$$

where $\boldsymbol{\rho} = (\rho, \theta)$ is the in-plane coordinate and $\mathbf{k} = (k, \alpha)$ is the Bloch vector of the subbands. The acceptor envelope function of Eq. (1) satisfies by construction the current conserving boundary conditions. In expansion (1), *n* runs over all the discrete subbands in the quantum well.

In our calculations we neglect the in-plane anisotropy of the subbands. Thus the acceptor Hamiltonian is invariant for rotations around the growth axis (*z*-axis) and the eigenstates have a definite *z*-component of angular momentum, which we call *m*. Because of time-reversal symmetry each eigenstate of angular momentum *m* is degenerate with a state of opposite *m*. For on-center acceptors, parity with respect to inversion is another good quantum number. By axial symmetry $G_n(\mathbf{k}) = e^{im\alpha} g_n(k)$: the radial function $g_n(k)$ is expanded in a variational basis of hydrogenic wavefunctions. The evaluation of the




Figure 1 - Energy levels of p -type states for on-center acceptors in GaAs/Ga_{0.6}Al_{0.4}As quantum wells as a function of the well width. The acceptor energies are given as binding energies with respect to the first heavy hole subband. The symmetry of the states is indicated by the absolute value of the angular momentum m and the parity with respect to inversion.

Coulomb potential between acceptor states is considerably simplified by the fact that the valence envelope functions are analytically known [2].

This method is particularly suited to find binding energies of excited states, which are extended in r -space; it is less suited to find the binding energy of the ground state, due to the neglect of the subband continuum. In Fig. 1, we present binding energies for excited acceptor states, which in the limit of wide wells go over into the bulk $2P_{3/2}[\Gamma_8]$, $2P_{5/2}[\Gamma_7 + \Gamma_8]$, and $3P_{3/2}[\Gamma_8]$ states. All these states have negative parity with respect to inversion. Comparison of the results of the present theory with far infrared absorption experiments [3] will be presented elsewhere.

2. References

- [1] J.M.Luttinger and W.Kohn, Phys. Rev. **97**, 869 (1955).
- [2] L.C.Andreani, A.Pasquarello and F.Bassani, Phys. Rev. **36**, 5887 (1987).
- [3] A.A.Reeder, B.D.McCombe, F.A.Chambers and G.P.Devane, Phys. Rev. B **38**, 4318 (1988).