Zeitschrift:	Helvetica Physica Acta
Band:	62 (1989)
Heft:	6-7
Artikel:	Theoretical calculation of the resistivity of the high-T_c superconductors
Autor:	Jorand, M. / Houghton, A.
DOI:	https://doi.org/10.5169/seals-116145

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

**Download PDF:** 15.05.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Theoretical calculation of the resistivity of the High-Tc superconductors

M. Jorand, Université de Lausanne, Institut de Physique Expérimentale, 1015 Dorigny, Switzerland

A. Houghton, Department of Physics, Brown University, Providence, RI 02906, USA

<u>Abstract</u> : Working within the Anderson lattice model, we calculate the first correction to the mean field theory and evaluate the self-energy  $\Sigma$  of the carriers. The resistivity is then given by Drude's formula  $\rho = m^*/ne^2\tau$  where  $\tau^{-1} = 2 \text{ Im } \Sigma$ .

1. Introduction

As shown by the observations, the interaction between the oxygen and copper atoms, within their two-dimensional layers, seems to play the leading role for a new microscopic mechanism leading to the pairing of the holes. We consider then the typical Anderson lattice model, used in heavy fermion systems [1], which can describe all electronic structure effects.

2. <u>Description of the model and some ideas about the calculations</u> The Anderson Hamiltonian, in the slave boson representation, writes [2, 3] :

$$H = \sum_{k\sigma} \varepsilon_{k\sigma} c_{k\sigma} c_{k\sigma} + \sum_{i} E_{o} b^{+}_{i} b_{i} + \sum_{i\sigma} E_{1} d^{+}_{i\sigma} d_{i\sigma} + \sum_{i} E_{2} a^{+}_{i} a_{i}$$
$$+ \sum_{ik\sigma} V_{k} [b_{i} d^{+}_{i\sigma} c_{k\sigma} e^{ikr_{i}} + a_{i} d^{+}_{i\sigma} c^{+}_{k-\sigma} e^{-ikr_{i}} sgn\sigma + h.c.]$$
(1)

Here the  $c_{k\sigma}$  are fermion operators for holes of spin  $\sigma$  in the o-band;  $E_1$ ,  $E_0$  and  $E_2$  are the energies of the d<sup>9</sup>, d<sup>10</sup> and d<sup>8</sup> Cu-states respectively represented by the localised fermion d<sub>i\sigma</sub> and by the bosons b<sub>i</sub> and a<sub>i</sub> respectively. The matrix elements  $V_k = \langle od | V | k\alpha \rangle$  define the hopping amplitude from an atomic d-state at the origin to an itinerant o-state in the band  $\alpha$ . Newns and Rasolt [2] worked out a mean field theory of this model. They could estimate the critical temperature and its dependence in the doping. Performing a 1/N expansion, where N = 2\sigma+1 is the spin degeneracy, we consider the first order correction and calculate the self-energy of the holes.

On a microscopic level, the resistivity is due to the scattering of the holes by many different processes but in our model, the most important one appears to be the scattering by the a-boson[4]. The main contribution to the self-energy writes :

$$\Sigma(p,\omega_n) = -V^2/\beta \sum_{\substack{i\omega_v \ q}} \Sigma D(q,i\omega_v) G(p+q,i\omega_n+i\omega_v)$$
(2)  
where G is the quasi-hole propagator (G<sup>-1</sup>(p,\omega) =  $\omega$  -  $\xi(p)$  where  $\xi(p) = p^2/2m^* - \mu$ )

and D is the a-boson propagator, given by

$$D^{-1}(q,i\omega_{V}) = -i\omega_{V} + U_{eff} - NV^{2}/\beta \sum_{i\omega_{n}} \sum_{k} \left[ G_{dd}(q-k, i\omega_{V}-i\omega_{n})G_{cc}(k,i\omega_{n}) + G_{cd}(q-k,i\omega_{V}-i\omega_{n})G_{cc}(k,i\omega_{n}) + G_{cd}(k,i\omega_{V}-i\omega_{N})G_{cc}(k,i\omega_{N}) + G_{cd}(k,i\omega_{N})G_{cc}(k,i\omega_{N}) + G_{cd}(k,i\omega_{N})G_{c$$

Performing the integration over the momentum and frequencies of the localised dand itinerant c-electrons, we find [4], in the approximation  $\omega_V$ ,  $v_F^*q$  << 4 $\pi$ KT :

$$D^{-1}(q,i\omega_{V}) = A \left[ B \ln(T/Tc) - i\omega_{V} + C q^{2} \right]$$
(4)  
where  $A = \frac{s_{o}^{2}}{s_{o}^{2} + \varepsilon_{d}^{2}} \frac{NV^{2}N_{s}\rho_{o}}{8\pi KT} \qquad B = \frac{4\pi KT}{\pi^{3}} \qquad C = \frac{7\zeta(3)}{\pi^{2}} \frac{v_{F}^{*}}{4\pi KT}$ 

The parameters  $\epsilon_d$  and  $s_o$  were calculated self-consistently in the mean field theory [2] while V ~2eV and  $\rho_o \approx 0.25 eV^{-1}$  are given by band structure calculations [2] . The effective Fermi velocity is  $v^*_F = m/m^* \ v_F \approx 0.2 \ v_F$  and  $N_s$  is the number of unit cells. Introducing (3) in (2) and performing the usual integrals over momentum and energy , one finds :

$$\rho = \frac{m^{*}}{ne^{2}\tau_{a}} = \frac{m^{*}}{ne^{2}} \frac{2}{h} \ln \Sigma = \frac{m^{*}}{hne^{2}} \frac{4V^{2}}{(hv^{*}_{F})^{2}A} E(\bar{x})$$
  
where  $E(\bar{x}) = \int_{-\mu}^{\infty} dz \frac{e^{z}}{(1+e^{z})^{2}} \int_{0}^{\bar{x}} dx \int_{-x-z}^{x-z} dy \frac{xy}{[(0.5lnT/Tc + x^{2}/4\pi)^{2} + y^{2}]} \frac{[n_{F}(y+z) + n_{B}(y)]}{\sqrt{x^{2} - (y+z)^{2}}}$ 

with  $z = \omega / KT$ ,  $y = \omega' / KT$ ,  $x = v_F^*q / KT$  and  $\overline{x} = v_F^*\overline{q} / KT$ 

# 3. Results and discussion

In the above expression, there is one single free parameter  $\overline{x} = v_F^* \overline{q} / KT$  which means that nothing in the theory fixes the value of the cut-off  $\overline{q}$  of the a-boson momentum. The temperature dependence of the resistivity is very much sensitive to the value of this parameter, but if we choose the value  $\overline{x} = 1$ , which satisfies all the approximations done in the calculations, the resistivity is almost linear in the temperature (see graph) as shown in the experiments [5]. Moreover, for n=3\*10<sup>27</sup> m<sup>-3</sup> [5], its

value is very close to the experimental one [5]  $\rho^{exp}(T=300) \approx 4 \mu\Omega m$ . We should of course understand the meaning of the relation  $\overline{q}$ = KT/v\*<sub>F</sub> if it has any and also analyse carefully Hall experiments to know more about the density n, taken here as a constant. Besides the results, the above calculation is interesting since it considers one of the many theories on the market of the High-Tc and we have to understand the normal state in order to know more about the superconducting state.



<u>Acknowledgements</u> : One of us (M.J.) is grateful to the Fonds National Suisse for its financial support during his stay at Brown University, where this work has been initiated.

4. <u>References</u>

[1] A.J. Millis and P.A. Lee : Phys. Rev. B, vol.35, no7, 3394 (1 March 1987)

- [2] D.M. Newns and M. Rasolt : IBM preprint
- [3] A. Houghton and A. Sudbo : to be published
- [4] M. Jorand and A. Houghton : to be published
- [5] S.W. Tozer, A.W. Kleinsasser, T. Penney, D. Kaiser and F. Holtzberg: Phys.Rev.Lett., vol.59, no15,1768 (12 october 1987)