

Zeitschrift: Helvetica Physica Acta
Band: 64 (1991)
Heft: 7

Artikel: A remark on long-range Stark scattering
Autor: Graf, Gian Michele
DOI: <https://doi.org/10.5169/seals-116335>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 16.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A remark on long-range Stark scattering

Gian Michele Graf*

Division of Physics, Mathematics, and Astronomy
California Institute of Technology
Pasadena, CA 91125

(26. IV. 1991)

Abstract. We propose a modified wave operator for long-range scattering in presence of an accelerating force. Its relevance in connection with classical scattering is addressed.

*Work supported in part by the U.S. NSF grant DMS-9101715

Introduction

The wave operators for the pair (H, H_0) of Stark Hamiltonians with $H = H_0 + V$ and

$$H_0 = \frac{p^2}{2} - g \cdot x, \quad 0 \neq g \in \mathbb{R}^\nu$$

on $\mathcal{H} = L^2(\mathbb{R}^\nu)$ do not exist [6] if $V = V(x)$ behaves as $|x|^{-\varepsilon}$, $\varepsilon \leq 1/2$ as $x \rightarrow \infty$. Recently, several proposals [3, 5, 8] for modified wave operators dealing with slowly decaying potentials were made. Here we add one more, based on a simple Dollard type argument: The classical free Stark trajectories $x(t) = gt^2/2 + p_0t + x_0$ suggest that $e^{-itH_0}e^{-i\Phi(t)}$ with

$$\Phi(t) := \int_0^t V(gs^2/2) ds$$

is a candidate for a comparison dynamics.

Theorem 1. *Let $V \in C^1(\mathbb{R}^\nu)$ be real-valued with*

$$|\nabla V(x)| \leq C(1 + |x|)^{-(1+\varepsilon)} \quad (1)$$

for some $C, \varepsilon > 0$. Then the wave operator

$$\Omega := s\text{-}\lim_{t \rightarrow \infty} e^{itH} e^{-itH_0} e^{-i\Phi(t)} \quad (2)$$

exists and is unitary.

The reason for stating this result comes from [6]. There a discrepancy between the quantum and the classical scattering problem was discovered, as for in the latter one the wave operators exist and are complete without modification basically as long as $V(x) = O(|x|^{-\varepsilon})$ for some $\varepsilon > 0$. We believe our result heals this discrepancy for the class of potentials we consider, since the comparison dynamics differs multiplicatively from the free Stark one only by a phase, which is physically irrelevant. Actually, no modification at all is needed if the asymptotic condition is formulated in the Heisenberg picture: The above result then implies that

$$\mu(A) := s\text{-}\lim_{t \rightarrow \infty} e^{itH} e^{-itH_0} A e^{itH_0} e^{-itH}$$

exists and is an automorphism on $\mathcal{L}(\mathcal{H})$. By contrast, this limit typically exists in long-range scattering only for $A \in \{H_0\}'$, the von Neumann algebra of operators commuting with bounded functions of H_0 [1]. In this respect (1) behaves as a short-range potential.

It follows from (1) that $\lim_{x \rightarrow \infty} V(x)$ exists. We may assume this limit to be zero, since (2) is not affected if the potential is shifted by a constant. Then $|V(x)| \leq C(1 + |x|)^{-\varepsilon}$, which is assumed also by [3, 5, 8]. However there stronger oscillations of the potential are allowed, since (1) is replaced by a weaker decay.

Remark. The hypothesis (1) can be weakened logarithmically [6].

Proofs

Let us state some kinematical remarks beforehand. The following propagation estimate for the free dynamics without electric field is well-known (see e.g. [4], Lemma 6.3): Let $f \in C_0^\infty(\mathbf{R}^v)$ with $f(y) = 0$ for $|y| \geq 1$, and let $\alpha > 1$. Then for $R > 0, t \geq 0$ and any $N > 0$

$$\|F(|x| > \alpha(R+t))e^{-itp^2/2}f(p)F(|x| < R)\| \leq C_N(R+t)^{-N},$$

with C_N independent of R, t . States supported in momentum space in $\{|y| \leq v\}$, $v > 0$ are accounted for through scaling by $v > 0$: We apply the unitary dilation U satisfying $UpU^{-1} = p/v$, $UxU^{-1} = vx$ to the operator in the estimate above, replace t by v^2t and R by vR , the result being

$$\|F(|x| > \alpha(R+vt))e^{-itp^2/2}f(p/v)F(|x| < R)\| \leq C_N[v(R+vt)]^{-N}.$$

An estimate for the free Stark dynamics is readily obtained by means of the Avron-Herbst formula

$$e^{-itH_0} = T(t)e^{-itp^2/2}, \quad T(t) := e^{-ig^2t^3/6}e^{itg \cdot x}e^{-it^2g \cdot p/2}$$

and of $T(t)x = (x - gt^2/2)T(t)$, namely

$$\|F(|x - gt^2/2| > \alpha(R+vt))e^{-itH_0}f(p/v)F(|x| < R)\| \leq C_N[v(R+vt)]^{-N}. \quad (3)$$

Proof of Theorem 1 (existence). By Cook's theorem we have to show that

$$\int_0^\infty \|U(x,t)e^{-itH_0}\varphi\| dt < \infty$$

for φ in a dense set D , where $U(x,t) := V(x) - V(gt^2/2)$. Choosing

$$D := \{f(p/v)F(|x| < R)\psi \mid f \in C_0^\infty(|y| < 1), v, R > 0, \psi \in \mathcal{H}\},$$

this follows from

$$\begin{aligned} & \|U(x,t)e^{-itH_0}f(p/v)F(|x| < R)\psi\| \\ & \leq \|U(x,t)F(|x - gt^2/2| \leq \alpha(R+vt))\|\|\psi\| + C'_N[v(R+vt)]^{-N}\|\psi\| = O(t^{-1-2\epsilon})\|\psi\|. \end{aligned} \quad (4)$$

Here we estimated the first term on the second line as follows: If $|x - gt^2/2| \leq \alpha(R+vt)$, then $U(x,t) = \nabla V(\tilde{x}) \cdot (x - gt^2/2)$ for some \tilde{x} with $|\tilde{x} - gt^2/2| \leq \alpha(R+vt)$. Then $|\tilde{x}| > |g|t^2/4$ for t large enough, and hence $\nabla V(\tilde{x}) = O(t^{-2(1+\epsilon)})$, proving (4). \square

Immediate consequences of the existence of the wave operator are that it is an isometry and that

$$\Omega = s\text{-}\lim_{s \rightarrow \infty} e^{isH}e^{-isH_0}e^{-i\Phi(s+t)} \quad (5)$$

for any $t \in \mathbf{R}$, since $\Phi(s+t) - \Phi(s) = \int_s^{s+t} V(g\tau^2/2) d\tau \rightarrow 0$ as $s \rightarrow \infty$. By comparing (5) with the definition (2) we get the intertwining relation $e^{itH}\Omega = \Omega e^{itH_0}$.

Another formulation of (3) is obtained using

$$e^{-itH_0}p = (p - gt)e^{-itH_0}, \quad e^{-itH_0}x = (x - pt + gt^2/2)e^{-itH_0},$$

which is

$$\|F(|x - gt^2/2| > \alpha(R+vt))f((p-gt)/v)F(|x - pt + gt^2/2| < R)\| \leq C_N[v(R+vt)]^{-N}. \quad (6)$$

The same remark also leads to

Corollary 2.

$$\lim_{t \rightarrow \infty} \|(\Omega - e^{-i\Phi(t)})f((p-gt)/v)F(|x - pt + gt^2/2| < R)\| = 0. \quad (7)$$

Proof. We apply Cook's estimate to (5) yielding

$$\|(\Omega - e^{-i\Phi(t)})e^{-itH_0}\varphi\| \leq \int_0^\infty \|U(x, s+t)e^{-i(s+t)H_0}\varphi\| ds = \int_t^\infty \|U(x, s)e^{-isH_0}\varphi\| ds.$$

We then set $\varphi = f(p/v)F(|x| < R)e^{itH_0}\psi$ and use (4). □

Let us now address the completeness question. We recall [2] that any $V \in L^\infty(\mathbf{R}^\nu)$ with $\lim_{x \rightarrow \infty} V(x) = 0$ is relatively compact with respect to H_0 and to H . This follows by density if it holds for V with compact support. In this case all terms in

$$\begin{aligned} V(H_0 + i)^{-1} - V(p^2/2 + i)^{-1} &= V(p^2/2 + i)^{-1}g \cdot x(H_0 + i)^{-1} \\ &= Vg \cdot x(p^2/2 + i)^{-1}(H_0 + i)^{-1} \\ &\quad + iV(p^2/2 + i)^{-1}g \cdot p(p^2/2 + i)^{-1}(H_0 + i)^{-1} \end{aligned}$$

are compact. We will also use that $\mathcal{H}_{cont} = \mathcal{H}$ for the spectral decomposition with respect to H [2]. Without using this result, one could prove that $\text{Ran } \Omega = \mathcal{H}_{cont}$ instead of unitarity of Ω by suitably modifying the proofs below.

Lemma 3. For $f \in C_\infty(\mathbf{R}^\nu)$

$$s\text{-}\lim_{t \rightarrow \infty} e^{itH}f\left(\frac{p-gt}{t}\right)e^{-itH} = f(0), \quad (8)$$

$$s\text{-}\lim_{t \rightarrow \infty} e^{itH}f\left(\frac{x-pt+gt^2/2}{t^2}\right)e^{-itH} = f(0). \quad (9)$$

Proof. We first remark that $D(p) \cap D(x)$ is invariant under e^{-itH} and that

$$\begin{aligned} e^{itH}(p - gt)e^{-itH}\psi &= p\psi - \int_0^t e^{isH} \nabla V(x) e^{-isH} \psi \, ds, \\ e^{itH}(x - pt + gt^2/2)e^{-itH}\psi &= x\psi + \int_0^t s e^{isH} \nabla V(x) e^{-isH} \psi \, ds \end{aligned} \quad (10)$$

for $\psi \in D(p) \cap D(x)$.

It suffices to prove the lemma for f depending on a single coordinate of its argument, since linear combinations of products of such functions are dense in $C_\infty(\mathbf{R}^\nu)$. By [7], Theorem VIII.20 (b) it is enough to show that (8, 9) hold for $f(y) = (y_i - z)^{-1}$, $z \notin \mathbf{R}$ and moreover strong and weak convergence are equivalent for resolvents. We thus set $p(t) := e^{itH} p e^{-itH}$, $x(t) := e^{itH} x e^{-itH}$ and have to show that for $\varphi \in \mathcal{H}$, $\psi \in D(p) \cap D(x)$

$$(\varphi, [(\frac{p_i(t) - g_i t}{t} - z)^{-1} - (-z)^{-1}] \psi) = z^{-1} ((\frac{p_i(t) - g_i t}{t} - \bar{z})^{-1} \varphi, \frac{p_i(t) - g_i t}{t} \psi),$$

vanishes as $t \rightarrow \infty$. Indeed, by (10) this is bounded by a constant times

$$\frac{1}{t} \|p_i \psi\| + \frac{1}{t} \int_0^t \|\partial_i V(x) e^{-isH} \psi\| \, ds \xrightarrow[t \rightarrow \infty]{} 0.$$

The limit just stated holds by the RAGE theorem ([7], Theorem XI.115), because $\partial_i V$ is relatively compact with respect to H . The other observable is treated analogously. \square

Lemma 3 is still a rather weak result, since it roughly says that $p(t) - gt = o(t)$, $x(t) - p(t)t + gt^2/2 = o(t^2)$, while these quantities are constants of motion for the free Stark problem. The idea to improve this is the following: Let

$$S_t := \{(x, p) \mid |p - gt| < \delta t_0, |x - pt + gt^2/2| < \delta t_0^2\}$$

for some $\delta, t_0 > 0$. Classically S_{t_0} at time t_0 is mapped into S_t at time $t \geq t_0$ under the free Stark evolution. Moreover

$$\begin{aligned} |x - gt^2/2| &\leq |x - pt + gt^2/2| + t|p - gt| < \delta t_0(t + t_0) \leq 2\delta t^2, \\ |x| &> (|g|/2 - 2\delta)t^2 \end{aligned}$$

i.e. the particle is far from the scatterer if $\delta < |g|/4$. Therefore S_{t_0} should be mapped approximately in S_t also under the full dynamics.

Let $f \in C_0^\infty(\mathbf{R}^\nu)$ with $f(y) = 0$ for $|y| \geq \delta$ and set

$$\begin{aligned} f_1(t, t_0) &:= f\left(\frac{x - pt + gt^2/2}{t_0^2}\right), \quad f_2(t, t_0) := f\left(\frac{p - gt}{t_0}\right), \\ \Theta(t, t_0) &:= f_1(t, t_0)^* f_2(t, t_0)^* f_2(t, t_0) f_1(t, t_0). \end{aligned} \quad (11)$$

Lemma 4. *If $\delta < |g|/4$, then*

$$\lim_{t_0 \rightarrow \infty} \sup_{t \geq t_0} \|e^{itH} \Theta(t, t_0) e^{-itH} - e^{it_0 H} \Theta(t_0, t_0) e^{-it_0 H}\| = 0. \quad (12)$$

Proof. The supremum in (12) is bounded by

$$\int_{t_0}^{\infty} \|D\Theta(t, t_0)\| dt, \quad (13)$$

where $D \cdot = i[H, \cdot] + \partial \cdot / \partial t$, i.e.

$$D\Theta = (Df_1)^* f_2^* f_2 f_1 + f_1^* (Df_2)^* f_2 f_1 + f_1^* f_2^* (Df_2) f_1 + f_1^* f_2^* f_2 (Df_1).$$

We start by computing

$$Df_1 = \int \hat{f}(s) i[V, e^{i(x-pt+gt^2/2)t_0^{-2} \cdot s}] ds = i \int \hat{f}(s) e^{i(x-pt+gt^2/2)t_0^{-2} \cdot s} (V(x+t_0^{-2}ts) - V(x)) ds$$

and split the integral into $|s| > t$ and $|s| \leq t$. The contribution to (13) of the first part is bounded by $O(t_0^{-N})$ for all $N > 0$, due to the decay of \hat{f} . For the other part, (6) implies that for $1 < \alpha < |g|(4\delta)^{-1}$

$$f_2^* f_2 f_1 = F(|x - gt^2/2| < \alpha\delta t_0(t + t_0)) f_2^* f_2 f_1 + O([t_0^2(t_0 + t)]^{-N}),$$

where the remainder contributes to (13) as little as $O(t_0^{-N})$. One is then left with estimating

$$\int_{|s| \leq t} |\hat{f}(s)| \| (V(x+t_0^{-2}ts) - V(x)) F(|x - gt^2/2| < \alpha\delta t_0(t + t_0)) \| ds \leq \text{const} \|s\hat{f}\|_1 t_0^{-2} t^{-(1+2\epsilon)},$$

contributing $O(t_0^{-2(1+\epsilon)})$ to (13). Here we used that if $|x - gt^2/2| < \alpha\delta t_0(t + t_0) \leq 2\alpha\delta t^2$ then $|V(x + t_0^{-2}ts) - V(x)| \leq t_0^{-2}t|s| |\nabla V(\tilde{x})|$ for some \tilde{x} with

$$|\tilde{x}| \geq |g|t^2/2 - |x - gt^2/2| - |x - \tilde{x}| \geq t^2(|g|/2 - 2\alpha\delta - t_0^{-2}),$$

where the bracket is positive for large t_0 . Terms arising from

$$Df_2 = i \int \hat{f}(s) e^{i(p-gt)t_0^{-1} \cdot s} (V(x - t_0^{-1}s) - V(x)) ds$$

are dealt with similarly. The integral restricted to $|s| \leq t$ is estimated up to a constant by $\|s\hat{f}\|_1 t_0^{-1} t^{-2(1+\epsilon)}$, its contribution to (13) thus by $O(t_0^{-2(1+\epsilon)})$. \square

Lemma 5. *Let f be as in the previous lemma, and real-valued with $f \leq 1$, $f(0) = 1$ in addition. Then*

$$\lim_{t_0 \rightarrow \infty} \sup_{t \geq t_0} \|(1 - f_2(t, t_0)f_1(t, t_0))e^{-itH}\psi\| = 0 \quad (14)$$

for any $\psi \in \mathcal{H}$.

Proof. Equations (8) and (9) imply $(e^{-it_0H}\psi, \Theta(t_0, t_0)e^{-it_0H}\psi) \rightarrow (\psi, \psi)$ as $t_0 \rightarrow \infty$. Then

$$\sup_{t \geq t_0} (\psi, \psi) - (e^{-itH}\psi, \Theta(t, t_0)e^{-itH}\psi) \xrightarrow{t_0 \rightarrow \infty} 0 \quad (15)$$

follows from (12). Suppressing arguments (t, t_0) we have $1 - f_2f_1 = (1 - f_1) + (1 - f_2)f_1$ and

$$\begin{aligned} \|(1 - f_1)e^{-itH}\psi\|^2 + \|(1 - f_2)f_1e^{-itH}\psi\|^2 \\ \leq (\|\psi\|^2 - \|f_1e^{-itH}\psi\|^2) + (\|f_1e^{-itH}\psi\|^2 - \|f_2f_1e^{-itH}\psi\|^2) \\ = (\psi, \psi) - (e^{-itH}\psi, \Theta(t, t_0)e^{-itH}\psi), \end{aligned}$$

since $f_i^2 \leq f_i$, $i = 1, 2$. Hence (14) is immediate from (15). \square

Proof of Theorem 1 (completeness). Given $\psi \in \mathcal{H}$,

$$\begin{aligned} \|(\Omega - e^{-i\Phi(t)})e^{-itH}\psi\| \\ \leq \|(\Omega - e^{-i\Phi(t)})f_2(t, t_0)f_1(t, t_0)e^{-itH}\psi\| + 2\|(1 - f_2(t, t_0)f_1(t, t_0))e^{-itH}\psi\| \end{aligned}$$

can be made arbitrarily small due to (7, 14) by first choosing t_0 and then t large enough. Thus

$$\psi = e^{itH}\Omega e^{i\Phi(t)}e^{-itH}\psi + o(1) = \Omega e^{itH_0}e^{i\Phi(t)}e^{-itH}\psi + o(1)$$

as $t \rightarrow \infty$, implying $\psi \in \overline{\text{Ran } \Omega} = \text{Ran } \Omega$. \square

Acknowledgements. We thank B. Simon for pointing out to us Ref. [1], and A. Jensen for useful discussions.

References

- [1] Amrein, W.O., Martin, Ph.A., Misra, B.: On the asymptotic condition in scattering theory. *Helv. Phys. Acta* **43**, 313-344 (1970).
- [2] Avron, J.E., Herbst, I.W.: Spectral and scattering theory for the Schrödinger operators related to the Stark effect. *Commun. Math. Phys.* **52**, 239-254 (1977).
- [3] Buslaeva, M.V.: On the asymptotic dynamics and on the spectral analysis for Schrödinger operator with accelerating potential. Preprint, Leningrad University.
- [4] Enss, V.: "Geometric methods in spectral and scattering theory of Schrödinger operators", in *Rigorous Atomic and Molecular Physics* ed. by G. Velo, A.S. Wightman, pp. 61-92, Plenum, New York (1981)
- [5] Jensen, A., Yajima, K.: On the long range scattering for Stark Hamiltonians. Preprint, Univ. of Aalborg (1990)
- [6] Jensen, A., Ozawa, T.: Classical and quantum scattering theory for Stark Hamiltonians with slowly decaying potentials. Preprint, Univ. of Aalborg (1990).
- [7] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics*, Vol. I-IV, Academic Press, (1972-79)
- [8] White, D.: The Stark effect and long range scattering in two Hilbert spaces. *Indiana Univ. Math. J.* **39**, 517-546 (1990).