

Zeitschrift: Mémoires de la Société des Sciences Naturelles de Neuchâtel
Herausgeber: Société des Sciences Naturelles de Neuchâtel
Band: 5 (1914)

Artikel: Terrestrial Isopods of Colombia
Autor: Richardson, Harriet
DOI: <https://doi.org/10.5169/seals-100116>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 08.02.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Terrestrial Isopods of Colombia

BY

HARRIET RICHARDSON

The following report is of the Isopods collected in Colombia by Dr Otto Fuhrmann during his recent scientific expedition to that country.

Family Oniscidæ

1. *Porcellio scaber* (Latr.)

Porcellio scaber LATREILLE, Hist. Crust. Ins., VII, 1804, p. 45. — Leach, Edinb. Encycl., VII, 1814, p. 406.

Oniscus granulatus LAMARCK, Hist. nat. des animaux sans vertèbres, V, 1818, p. 154.

Porcellio nigra SAY, Journ. Phil. Acad. Nat. Sc., I, 1818, p. 432.

Porcellio scaber DESMAREST, Consid. Crust., 1825, p. 321. — Brandt and Ratzeburg, Med. Zool., II, 1830-1834, p. 77, pl. XII, figs. 1-4 and A-B. — Brandt, Bull. Soc. Imp. de Naturalistes de Moscou, VI, 1833, p. 14.

Porcellio brandtii MILNE EDWARDS, Hist. Nat. Crust., III, 1840, p. 168.

Porcellio granulatus MILNE EDWARDS, Hist. Nat. Crust., III, 1840, p. 169, pl. XXXII, fig. 21.

Porcellio scaber MILNE EDWARDS, Cuvier, Rg. An., 1849, pl. LXXI-LXXI bis. — Koch, Deutschlands Crust., 1835-1844, p. 34.

Porcellio dubius KOCH, Deutschlands Crust., 1835-1844, p. 34.

Porcellio gemmulatus DANA, Crust., U. S. Expl. Exp., XIV, 1853, p. 725, pl. XLVII.

Porcellio scaber BATE AND WESTWOOD, Brit. Sess-eyed Crust., II, 1868, p. 475. — Budde-Lund, Crust. Isop. Terrestria, 1885, pp. 129-131. — Sars, Crust. Norway, II, 1899, pp. 176-177, pl. LXXVII. — Chilton, Trans. Linn. Soc. London (2), VIII, 1901, Pt. 4, p. 139.

Localities. Environs de Bogota (2600 m.); dans une Quebrada au-dessus de Bogota (C. or. 2750 m.); près de la chute du Tequendama (C. or. 2210 m.); Zipaquira (C. or. 2600 m.); entre Boca del Monte et Tambo (C. or. 1800 m.).

Porcellio scaber var. *marmoratus* BRANDT and RATZEBURG and *Porcellio scaber* var. *marginatus* Brandt and Ratzeburg were also collected from the above localities.

2. *Procellionides pruinosus* (Brandt).

Porcellio pruinosus BRANDT, Bull. Soc. Imp. Naturalistes de Moscou, VI, 1833, p. 19.

Porcellio truncatus MILNE EDWARDS, Hist. Nat. Crust., III, 1840, p. 171.

Porcellio maculicornis KOCH, Deutschl. Crust., 1835-44, p. 34.

Metoponorthus pruinosus BUDDE-LUND, Crust. Isop. Terrestria, 1885, pp. 169-171; Entom. Meddelel., IV, 1893-94, p. 118. — Sars, Crust. Norway, II, 1899, pp. 184-185, pl. LXXX, fig. 2. — Chilton, Trans. Linn. Soc. Lond. (2), VIII, 1901, p. 141.

Procellionides pruinosus Stebbing, Records of the Indian Museum, VI, Pt. 4, 1911, p. 189.

Localities. — Sur le Porce (rivière) dans un tronc de saule avec des fourmis près Medellin (C. c. 1547 m.); Cafetal Camelia près Angelopolis (C. c. 1820 m.); près du Ruiz (C. c. 3700 m.).

3. *Philoscia variegata* Dollfus.

Philoscia variegata DOLLFUS, Ann. Soc. Entomologique de France, 1893, LXII, p. 343, pl. X, fig. 10.

Localities. — Bellio près Medellin vallée du Porce (C. c. 1547 m.); plantation de café Camelia près d'Angelopolis (C. c. 1820 m.); Alto San Miguel (C. c. 2200 m.); près du Ruiz (C. c. 3700 m.); entre Guaduas et Sensitiva (C. or. 1600 m.); entre Honda et Guaduas (C. or. 1200 m.); environs de Bogota (C. or. 2700 m.); près de la chute du Tequendama (C. or. 2210 m.); Zipaquira (2600 m.); entre Boca del Monte et Tambo (C. or. 1800 m.); Paramo Cruz Verde (C. or. 3600 m.).

In the male the first three pairs of legs have the propodus dilated as shown in the figure by DOLLFUS. In the female the propodus of the first three pairs of legs is not dilated and the first article of the flagellum of the second antennæ is not as long as in the male.

4. *Alloniscus papillosus* Budde-Lund.

Alloniscus papillosus BUDDE-LUND, Entomologiske Meddelelser, IV, 1893-94, pp. 123-124.

Locality. — Sur le Porce (rivière), dans un tronç de saule avec des fourmis près Medellin (C. c. 1547 m.).

The specimens are the same size as those of BUDDE-LUND's, being 3 1/2 mm. in length. Instead of being marked with black, they are marked with a reddish brown color.

The specimens referred to this species by DOLLFUS¹ are much larger and probably represent a different species. About twenty specimens were collected.

Family Armadillididæ

5. *Cubaris grenadensis* (Budde-Lund).

Armadillo grenadensis BUDDE-LUND, Entomologiske Meddelelser, IV, 1893-94, pp. 115-116. — Dollfus, Proc. Zool. Soc. London, 1896, pp. 392-393. — Richardson, Bull. U. S. Nat. Museum, No. 54, 1905, p. 651.

Locality. — Calamar sur le Magdalena.

6. *Sphæroniscus flavomaculatus* Gerstäcker.

Sphæroniscus flavomaculatus GERSTÄCKER, Entomolog. Zeit., 1854, pp. 314-315, fig. 2. — Budde-Lund, Crust. Isop. Terrestria, 1885, p. 45.

Locality. — Entre Boca del Monte et Tambo (C. or. 1800 m.).

7. *Sphæroniscus granulatus* Dollfus.

Sphæroniscus granulatus DOLLFUS, Ann. Soc. Entomologique de France, LXII, 1893, p. 341.

Locality. — Sur le Cauca, Puerto de los Pobres (C. c. 800 m.).

Two specimens measuring 5 mm. in length. The inner branch of the uropoda does not quite reach the tip of the outer branch.

8. *Sphæroniscus frontalis* sp. nov.

This species is very close to *Sphæroniscus flavomaculatus*, GERSTÄCKER, but differs in not having the epistome produced a great distance beyond the frontal margin of the

¹ Ann. Soc. Entomologique de France, 62, 1893, p. 342, pl. 10, fig. 9.

head ; in having the first segment of the thorax with the lateral margin not rolled back so much and not so wide as in *S. flavomaculatus* and in not having the lateral margins of the segments lighter in color.

Six specimens were collected at the plantation of café, Argelia (1800 m.) near Viota (C. or.) a uniform orange brown color, with light wavy lines of yellow on either side of the median line.

One specimen from the plantation of the café Buenavista (1020 m.) near Viota (C. or.), is dark gray, with wavy lines of yellow on either side of the median line.

LIST OF REFERENCES

- BUDDE-LUND, G. Crustacea Isopoda Terrestria, 1885, Hauniæ.
— Landisopoder fra Venezuela, indsamlede af Dr. Fr. Meinert. Entomologiske Meddelelser, IV, 1893-94, pp. 111-129, Kjobenhavn.
- DOLLFUS, ADRIEN. Sur quelques Isopodes du Musée de Leyde. Notes from the Leyden Museum, XI, 1889, pp. 91-94, pl. v. fig. 2, Leyden.
— Voyage de M. E. Simon au Venezuela (Décembre 1887-Avril 1888). Isopodes terrestres. Ann. Soc. Entomologique de France, LXII, 1893, pp. 339-345, pls. 9-10, Paris.
— On West Indian terrestrial Isopod Crustaceans. Proc. Zool. Soc. London, 1896, pp. 338-400, London.
- GERSTÆCKER, A. Ueber eine neue Myriapoden-und Isopoden-Gattung. Entomolog. Zeitung, XV, 1854, pp. 314-315, fig. 2, Stettin.
- SARS, G.-O. An account of the Crustacea of Norway, II, 1899, pp. 169-192, pls. LXXV-LXXXIII, Bergen.
-