

Toxins from Cyanobacteria : chemical and biological studies addressing their mode of action

Autor(en): **Grundler, Verena / Gademann, Karl**

Objektyp: **Article**

Zeitschrift: **Mitteilungen der Naturforschenden Gesellschaften beider Basel**

Band (Jahr): **15 (2014)**

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-676620>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Toxins from Cyanobacteria – Chemical and Biological Studies Addressing their Mode of Action

VERENA GRUNDLER AND KARL GADEMANN

Zusammenfassung: Steigende Temperaturen, extensive Landwirtschaft sowie die Globalisierung begünstigen den unkontrollierbaren Wuchs von Cyanobakterien (alte Bezeichnung: Blaualgen). Da sie zu den ältesten Lebensformen auf unserem Planeten zählen, besiedeln sie viele Habitate, wobei die Mehrheit in Gewässern zu finden ist. Unglücklicherweise produzieren diese Organismen eine Reihe von verschiedenen Toxinen, weswegen von einer Algenblüte unter Umständen große Gefahren für Mensch und Tier ausgehen können. Ein kurzer Überblick über diese interessante, jedoch teilweise gefährliche Spezies wird im Rahmen dieser Arbeit gegeben.

Abstract: Rising temperatures, extensive agriculture together with globalization favor the uncontrolled growth of cyanobacteria, formerly called blue-green algae. These organisms belong among the oldest life forms on our planet populate many habitats, in particular marine and freshwater environments. Unfortunately, those organisms are known to produce various toxic compounds, wherefore a bloom can cause great danger for humankind and animals. This article gives a short overview over this interesting, yet dangerous species.

Key words: Toxic compounds, cyanobacteria, phytoplankton bloom, Microcystin-LR, Anatoxin-a, Saxitoxin, Nodularin.

Introduction

Over the last years, warning signs reporting danger of intoxication in shore areas have been emerging in many areas over the world (Figure 1). In many cases, people witnessed symptoms of nausea and dizziness while being exposed to a refreshing breeze from the ocean. More severe cases, from skin itching to acute intoxication, have been reported after immersion and ingestion of contaminated water.

These symptoms are often caused by certain types of microorganisms, such as cyanobacteria (formerly called blue-green algae.) These prokaryotic phototrophs are among the oldest species on our planet, as indicated by fossil records from western Australia, and they have been postulated to exist on earth since 3.5 billion years ago (Schopf and Packer, 1987). Cyanobacteria can be found in many habitats on earth, in particular freshwater, brackish as well as saline water environments (Carmichael and Li, 2006). In addition, they can adapt and survive environmental stress, such as dramatic temperature changes, dry periods and very high and low temperatures. For example, cyanobacteria are able to grow in polar regions, in antarctic terrestrial environment or in hot springs in Yellowstone National park at 74 °C (Briand et al., 2004; Jungblut et al., 2009; Reichwaldt and Ghadouani, 2012; Tang et al., 1997; Ward et al., 1998). Even extreme hostile environments such as volcanic ash and desert sand can be cultivated by cyanobacteria (Dor and Danin, 1996; Gerasimenko et al., 2013). Some species as *Synechococcus/Synechocystis* spp. can live in symbiosis with dinoflagellates, whereas others associate

with fungi and plants (Gordon et al., 1994; Hyvarinen et al., 2002; Meeks, 1998). This synergy is often due to the remarkable ability of cyanobacteria to fix nitrogen from the atmosphere, which then is further enzymatically converted into ammonium derivatives (Herrero et al., 2001; Latysheva et al., 2012). Responsible for this process are specialised cells, the so-called heterocysts, which have a large and round shape and possess thick cell walls with oxygen-binding glycolipids (Pitois et al., 2000). As energy metabolism for life, these photoautotrophic organisms use photosynthesis and likewise, this circumstance has been suggested to play a key role in evolution of life on Earth as cyanobacteria are assumed to have generated the oxygen in the atmosphere (Dismukes et al., 2001; Stanier and Bazine, 1977).

Cyanobacteria are unicellular to multicellular prokaryotes, which belong to the *Bacteria* domain, and these multicellular bacteria can be further categorized in filamentous, undifferentiated and differentiated species (Flores et al., 2006; Rokas, 2008). The multicellular bacteria are able to communicate intercellularly, to differentiate and form cell-cell adhesion (Flores and Herrero, 2010). An example for an unicellular bacteria would be *Microcystis aeruginosa* (Figure 2A). An example of a multicellular filamentous and undifferentiated bacteria is represented by *Planktothrix* and *Anabaena* sp. belongs in the group of multicellular differentiated cyanobacteria (Schirrmeister et al., 2011).

As any species, cyanobacteria produce a great variety of secondary metabolites such as carotenoids, fatty acids, lipopeptides, polysaccharides and as well as different bioactive molecules. A large number of cyanobacterial compounds are peptides or contain a peptic substructure and, so far, more than 600 members have been described (Dixit and Suseela, 2013; Welker and Döhren, 2006). A large structural diversity exists within these compounds, and the incorporation of unusual or modified amino acids is often found. As cyanobacteria have populated earth for a long time period, it is not surprising that many of the above mentioned molecules show significant bioactivity. It has been reported that some secondary metabolites of cyanobacteria show anti-




Fig. 1: Marion Reservoir, Kansas.
Photo Credit: Jennifer L. Graham USGS




Fig. 2:

A) *Microcystis aeruginosa*

B) *Planktothrix rubescens*;

Photo Credit: Thomas Posch

C) *Nostoc* sp.;

Photo Credit: Bettina Eugster & Esther Kohler.

cancer, cytotoxic, antibiotic, antifungal and anti-viral activity, as well have multi-drug resistance, antimycotic, immunosuppressive and anti-malarial features (Burja et al., 2001). As the compounds produced by cyanobacteria can be characterized by high chemical stability and water solubility, they can be considered also as interesting candidates for medical applications, such as the anticancer agent dolastatin, which failed




Fig. 3: Phytoplankton bloom in the Baltic Sea.
Photo Credit: European Space Agency, ESA

however in clinical trials due to toxicity effects (Luesch et al., 2001; Simmons et al., 2005).

In addition to these compounds that may have a positive impact by being potentially used as drugs, cyanobacteria produce several highly toxic molecules, the cyanotoxins, which pose a danger to humans, animals and the ecosystem alike. These toxins are also characterized by a high chemical stability and water solubility and thus have the potential for severe intoxication of humans and animals (Carmichael, 1992). In fact, due to eutrophication, cyanobacteria often prosper in aquatic environment and cause dangerous blooms all over the world. In addition, climate change and therefore rising temperatures on our planet, the growth of toxic cyanobacteria might be further facilitated and become an even greater problem (Paerl and Huisman, 2008; Pitois et al., 2000). Toxic blooms mainly affect warm regions as Africa or South America, where every year many people are effected and water supply is endangered (Oberholster et al., 2005; Lagos et al., 1999). In addition, Europe is also struggling with the problem of toxic cyanobacteria, for example in the Baltic Sea (Figure 3), where the occurrence of blooms increased over the past years (Kahru et al., 2007). These phenomena are visible from space, as an image from a satellite clearly demonstrates (Figure 3). Furthermore, European lakes are as well affected by increasing density of toxic cyanobacteria (Eiler et al., 2013; Kurmayer and Gumpenberger, 2006).

Various toxins of cyanobacteria

A major problem occurs after the collapse of those blooms. This way various toxins get released and thereby pollute fresh and marine water, which creates a serious threat to water supply, public health, livestock and wildlife (Paerl et al., 2001). Another danger for health is inflicted by direct contact with poisoned water through swimming in a lake with a cyanobacteria bloom or drinking contaminated water. Besides this, another serious problem poses the circumstance that these toxins can enter the food chain via accumulation in fish, mussels or shell fish and on this way find their way into humans (van Apeldoorn et al., 2007).




Fig. 4: **A)** Microcystin-LR; **B)** Anatoxin-a; **C)** Anatoxin-a(s); **D)** Saxitoxin; **E)** Nadularin; **F)** Cylindrospermopsin; **G)** β -N-methylamino-L-alanine

Over the past years, many cases of poisoning by cyanobacteria were reported, whereby the regular intoxication occurs via oral intake, although other uptake mechanisms are also possible, such as inhalation or skin contact (Stewart et al., 2006). The main cases of intoxication happen with animals, for example at a lake in Canada where approximately 1000 bats died after ingesting cyanobacterial polluted water (Pybus et al., 1986). In addition, also humans suffer from cyanobacterial poisoning. A tragic accident of acute intoxication with microcystin containing water occurred in Brazil in the year 1996, where dialysis patients were treated with contaminated water and this conjuncture lead to the death of 52 people (Azevedo et al., 2002). A more recent case of intoxication of a young person was reported from Argentina, who dived in a lake with a cyanobacteria bloom and thus showed the clear symptoms of intoxication, like e.g. nausea and respiratory distress (Giannuzzi et al., 2011). The main toxin, which has been suggested to be responsible for these poisonings is Microcystin-LR, which belongs to a wide group of cyclic peptides, the microcystins with a great variety of 80 derivatives known to date (Zeller et al., 2011). Besides the general core structure *cyclo(-D-Ala-Xaa-D-MeAsp-Yaa-Adda-D-Glu-Mdha-)*, two diverse amino acids (Xaa, Yaa) lead to this great structural diversity (Mikalsen et al., 2003). Within this compound class, MC-LR (Xaa = Leu, Yaa = Arg) has an exceptional position as most studied microcystin, due to its high LD₅₀ value of 50 µgkg⁻¹ (Rodríguez et al., 2008). In fact it is one of most potent and toxic compound produced by cyanobacteria and on this account the WHO set a guideline value of MC-LR in drinking water of 1 µgL⁻¹ (Gupta et al., 2003; WHO, 2003).

In addition to the hepatotoxic effect in humans through inhibition of protein phosphatases 1 and 2A, chronic consumption of MC polluted water can cause liver cancer (MacKintosh et al., 1990; Ueno et al., 1996; Fujiki and Suganuma, 2011). Moreover, MC-LR seems to cause DNA damage, apoptosis, cell-signaling disruption and endoplasmatic reticulum stress (Campos and Vasconcelos, 2010; Christen et al., 2013; Žegura et al., 2008). The known transport of MCs in the

cell is the organic anion polypeptide transporter (OATP), which is present e.g. in human hepatocytes (Fischer et al., 2010; Magalhães et al., 2003). In aquatic organisms, which are especially exposed to cyanobacterial toxins, microcystins shows also toxic effects. For example, in *Daphnia magna*, which is an important grazer using cyanobacteria as a food source, reproduction is disturbed, which is reflected in higher mortality and suppressed development of the offspring (Dao et al., 2010; Lürding, 2003; Ortiz-Rodríguez et al., 2012). In fish and other invertebrates, the toxin leads to hepatotoxic effects and accumulation in tissues (Cazenave et al., 2005). Nevertheless, although intensive research on MC-LR has been carried out since decades, the reason for the strong toxicity is not yet fully understood.

In addition to microcystins, cyanobacteria produce a wide range of other potent, various toxins and the structural variety of these toxic compounds reaches from peptides, linear or heterocyclic, to lipid compounds (Bláha et al., 2009). Based on their toxicity effect and their mode of action and uptake, they can be categorized into different groups; the neurotoxins, the cytotoxins, the dermatotoxins, the irritant toxins and the hepatoxins (Wiegand and Pflugmacher, 2005).

For example, the neurotoxic compounds anatoxin-a, anatoxin-a(s) and saxitoxin are among the most toxic compounds produced by cyanobacteria, but display, due to their rare occurrence and therefore reduced exposure, a minor danger to humans when compared to the hepatoxins such as the microcystins and nodularins (Mankiewicz et al., 2003). Anatoxin-a is a bicyclic, secondary amine, which is a strong agonist at the nicotinic acetylcholine receptors (Devlin et al., 1977). Toxicity symptoms reveal quickly after consumption of contaminated water and are manifested in paralysis, tremors, convulsions and death (Rogers et al., 2005). Anatoxin-a(s) constitutes an organophosphate, structurally not related to anatoxin-a, which irreversibly blocks acetylcholinesterase and enhances salivation (Dittmann et al., 2013; Matsunaga et al., 1989). Furthermore, the administration of pure toxin to mice leads to fast death and respiratory failure

within 240 min. (Onodera et al., 1997). Saxitoxin, produced by freshwater cyanobacteria as well as marine dinoflagellates, is a strong neurotoxin, which accumulates in the food chain and causes paralytic shellfish poisoning (Pearson et al., 2010). Symptoms occur rapidly after intake of poisoned food and are manifested as preliminary numbness around the lips, which later affects the whole body and can lead to respiratory paralysis (Cusick and Sayler, 2013). Its chemical core structure is a trialkyl tetrahydropurine, whereas 30 derivatives are known (Llewellyn, 2006). Another example for a cyanobacterial compound is cylindrospermopsin, a cytotoxic and genotoxic alkaloid, which covalently binds to DNA and shows a toxicity effect in liver, kidney and lung (Moreira et al., 2013). Additionally, it is demonstrated to be carcinogenic and has evidence to inhibit protein synthesis (Falconer and Humpage, 2006). The strong liver toxicity effect is explained by a oxidation of the toxin by CYP450, which leads to a more potent toxic product (Runnegar et al., 1995). Additional toxins are, for example, lyngbyatoxin-a, debromoaplysiatoxin and aplysiatoxin, representing common dermatotoxins (Mankiewicz et al., 2003). The typical common clinical symptom of these toxins are dermatitis, complemented by tumor promotor activity and protein kinase C activation in the case of debromoaplysiatoxin and aplysiatoxin, as well as gastrointestinal inflammation for lyngbyatoxin-a (Cardellina et al., 1979; Mynderse et al., 1977). Another class of toxins are the irritant toxins. The lipopolysaccharides fall in this category, as they represent a crucial component of the cell wall of all bacteria (Bláha et al., 2009). The toxicity effect occurs after contact with these toxins and appears as allergic response, gastroenteritis and inflammation (Mur et al., 1999). Due to the less toxic effect in comparison to other bacteria, the importance of these toxins is slightly smaller classified then the abovementioned (Keleti and Sykora, 1982). The last compound class are the hepatoxins, which represent the most studied toxins so far (see the discussion on MC above). As mentioned above, cyanobacteria are known to be hepatotoxic and exposure to blooms cause typical clinical symptoms for liver damage, such as

increased liver weight, haemorrhage and blood pooling and this deep red coloration, as well as the enhanced activity of glutamate pyruvate transaminase/alanine aminotransferase, lactate dehydrogenase and alkaline phosphatase (Kumar et al., 2006). The two common hepatoxins are represented by the microcystins (see above) and nodularin, whereas this toxin shows a similar toxicity effect as microcystin (Zurawell et al., 2005). The characteristic feature of both compounds is a strong inhibition effect of protein phosphatase 1 and 2A, as well as promotion of liver cancer (MacKintosh et al., 1990; Ullah, 2011). Nodularin has a similar structure as microcystin, containing the Adda function, D-Glu, D-MeAsp, L-Arg and MeDhb and thus leading to similar toxic effects. In contrast to the microcystins, however, a much smaller number has been described so far, as only seven isoforms of this toxin have been reported (Pearson et al., 2010).

In addition to the above-mentioned categorized toxins, cyanobacteria produce other classes of natural products, which share structural similarity and biological effects. These compounds can be further classified regarding their general core structure into aeruginosins, microginins, anabaenopeptins, cyanopeptolins, microcystins, microviridins and cyclamides (Welker and Döhren, 2006). A large group of these toxins are the aeruginosins, which are trypsin-type serine protease inhibitors produced by *microcystis* and *oscillatoria* (Murakami et al., 1995; Shin et al., 1997). More than 40 variants are known up to date, and the characteristic core structure of this peptide class is determined by the presence of the unusual amino acid 2-carboxy-6-hydroxyoctahydroindole (Choi) and the hydroxyphenyllactic acid (Hpla), further a variable amino acid (Leu, Ile, Phe, or Tyr) and an arginine derivative (Elkobi-Peer et al., 2013). Another group of serine inhibitors are the cyclic peptides belonging to the group of cyanopeptolins, which are characterized by the amino acid 3-amino-6-hydroxy-2-piperidone (Ahp) (Martin et al., 1993). These toxins show a great structural diversity based on the variation of several various amino acids as well as derivatization or chlorination (Matern et al., 2001; Weckesser et al., 1996). The synthesis of these large biomolecules is mostly performed

by large enzyme complexes, the nonribosomal peptide synthetases (NRPS) or a combination of polyketide synthases and NRPSs systems (Schwarzer et al., 2003). The gene clusters of those synthetases contain highly conserved NRPS operons, containing the *mcyABC* gene cluster, within the different strains and thus suggest a horizontal gene transfer as an explanation for the similarity of the produced compounds (Mikalsen et al., 2003; Rounge et al., 2008). Recently, B-N-methylamino-L-alanine, a neurotoxic amino acid produced by cyanobacteria, received a lot of attention as it is considered to be a trigger for human neurological diseases such as Amyotrophic Lateral Sclerosis, Parkinson and Alzheimer's disease (Banack et al., 2010; Pablo et al., 2009). The disease seems to be initiated by an overactivation of neuroexcitatory glutamate receptors, which leads to damage of the neurons (Vyas and Weiss, 2009).

Ecological and biological role of cyanobacterial toxins

Despite all the research done on cyanobacterial toxins, the ecological and biological role of cyanobacterial toxins is still under investigation, whereby the recent production purpose is mainly linked with the protection against grazers (Blom et al., 2001; Lüring, 2003; Wilson et al., 2006). In contrast to this theory, phylogenetic analyses suggests that the synthetase genes developed simultaneously with the housekeeping genes and thus are older than grazers (Rantala et al., 2004). Another purpose is assumed to act as allelochemicals to repel other photosynthetic competitors or

the possibility of a physiological function as for example cell signalling or iron scavenging is also thinkable, since the long existence of these compounds and the effort of production suggest a more important role in the organism (Berry et al., 2008; Holland and Kinnear, 2013). In any case the existence of mutants of cyanobacteria, which lost the ability to produce microcystin by inactivation of the microcystin synthetase through mutations in the gene cluster, could be an hint to exclude the toxic purpose of microcystin (Christiansen et al., 2008). But this issue is getting even more complex facing the fact that non-toxic cyanobacteria exist and also form blooms without the missing disadvantage in growth compared to their poisonous siblings (Merel et al., 2013; Ostermaier and Kurmayer, 2009). Therefore, the missing «final explanation» for the purpose of the toxin production renders toxic cyanobacteria still a mysterious species.

Conclusions

In conclusion, the occurrence of toxic cyanobacteria in marine and freshwater environments will likely increase in the near future, due to climate change and eutrophication. As shown in this highlight article, the presence of a manifold of toxic compounds in these phototrophic organisms continues to pose a challenge to a variety of recreational, nutritional and industrial activities. As the mechanism of action for toxicity for many compounds is yet poorly understood, continuous research activities are needed to deepen our understanding of the chemical potential of cyanobacteria.

References

- Azevedo, S.M.F.O., Carmichael, W.W., Jochimsen, E.M., Rinehart, K.L., Lau, S., Shaw, G.R., Eaglesham, G.K., 2002. Human intoxication by microcystins during renal dialysis treatment in Caruaru-Brazil. *Toxicology* 181–182, 441–446.
- Banack, S.A., Caller, T.A., Stommel, E.W., 2010. The cyanobacteria derived toxin Beta-N-methylamino-L-alanine and amyotrophic lateral sclerosis. *Toxins (Basel)* 2, 2837–2850.
- Berry, J.P., Gantar, M., Perez, M.H., Berry, G., Noriega, F.G., 2008. Cyanobacterial toxins as allelochemicals with potential applications as algacides, herbicides and insecticides. *Mar Drugs* 6, 117–146.
- Bláha, L., Babica, P., Maršálek, B., 2009. Toxins produced in cyanobacterial water blooms – toxicity and risks. *Interdiscip Toxicol* 2, 36–41.
- Blom, J.F., Robinson, J.A., Jüttner, F., 2001. High grazer toxicity of [D-Asp³,(E)-Dhb⁷]microcystin-RR of *Planktothrix rubescens* as compared to different microcystins. *Toxicon* 39, 1923–1932.
- Briand, J.F., Leboulanger, C., Humbert, J.F., 2004. *Cylindrospermopsis raciborskii* (cyanobacteria) invasion at mid-latitudes: Selection, wide physiological tolerance, or global warming? *J. Phycol.* 40, 231–238.
- Burja, A.M., Banaigs, B., Abou-Mansour, E., 2001. Marine cyanobacteria—a prolific source of natural products. *Tetrahedron* 57, 9447–9377.
- Campos, A., Vasconcelos, V., 2010. Molecular mechanisms of microcystin toxicity in animal cells. *Int J Mol Sci* 11, 268–287.
- Cardellina, J.H., Marnier, F.J., Moore, R.E., 1979. Seaweed dermatitis: structure of lyngbyatoxin A. *Science* 204, 193–195.
- Carmichael, W.W., 1992. Cyanobacteria secondary metabolites—the cyanotoxins. *J. Appl Microbiol.* 72, 445–459.
- Carmichael, W. W., Li, R., 2006. Cyanobacteria toxins in the salton sea. *Saline System* 19;2:5.
- Cazenave, J., Wunderlin, D.A., de Los Angeles Bistoni, M., Amé, M.V., Krause, E., Pflugmacher, S., Wiegand, C., 2005. Uptake, tissue distribution and accumulation of microcystin-RR in *Corydoras paleatus*, *Jenynsia multidentata* and *Odontesthes bonariensis*. A field and laboratory study. *Aquat. Toxicol.* 75, 178–190.
- Christen, V., Meili, N., Fent, K., 2013. Microcystin-LR Induces Endoplasmatic Reticulum Stress and Leads to Induction of NFKB, Interferon-Alpha, and Tumor Necrosis Factor-Alpha. *Environ. Sci. Technol.* 47, 3378–3385.
- Christiansen, G., Molitor, C., Philmus, B., Kurmayer, R., 2008. Nontoxic strains of cyanobacteria are the result of major gene deletion events induced by a transposable element. *Mol. Biol. Evol.* 25, 1695–1704.
- Cusick, K.D., Sayler, G.S., 2013. An overview on the marine neurotoxin, saxitoxin: genetics, molecular targets, methods of detection and ecological functions. *Mar Drugs* 11, 991–1018.
- Dao, T.S., Do-Hong, L.-C., Wiegand, C., 2010. Chronic effects of cyanobacterial toxins on *Daphnia magna* and their offspring. *Toxicon* 55, 1244–1254.
- Devlin, J.P., Edwards, O.E., Gorham, P.R., 1977. Anatoxin-a, a toxic alkaloid from *Anabaena flos-aquae* NRC-44h. *Can. J. Chem.* 55, 1367–1371.
- Dismukes, G.C., Klimov, V.V., Baranov, S.V., Kozlov, Y.N., DasGupta, J., Tyryshkin, A., 2001. The origin of atmospheric oxygen on Earth: the innovation of oxygenic photosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 98, 2170–2175.
- Dittmann, E., Fewer, D.P., Neilan, B.A., 2013. Cyanobacterial toxins: biosynthetic routes and evolutionary roots. *FEMS Microbiology Reviews* 37, 23–43.
- Dixit, R.B., Suseela, M.R., 2013. Cyanobacteria: potential candidates for drug discovery. *Antonie Van Leeuwenhoek* 103, 947–961.
- Dor, I., Danin A., 1996. Cyanobacterial desert crust in the dead sea valley, Israel. *Algological Studies* 83, 197–206.
- Eiler, A., Drakare, S., Bertilsson, S., Perntaler, J., Peura, S., Rofner, C., Simek, K., Yang, Y., Znachor, P., Lindström, E.S., 2013. Unveiling Distribution Patterns of Freshwater Phytoplankton by a Next Generation Sequencing Based Approach. 8, *Plos one* e53516.
- Elkobi-Peer, S., Singh, R.K., Mohapatra, T.M., Tiwari, S.P., Carmeli, S., 2013. Aeruginosins from a *Microcystis* sp. Bloom Material Collected in Varanasi, India. *J. Nat. Prod.* 76, 1187–1190.
- Falconer, I.R., Humpage, A.R., 2006. Cyanobacterial (blue-green algal) toxins in water supplies: *Cylindrospermopsins*. *Environ. Toxicol.* 21, 299–304.
- Fischer, A., Hoeger, S.J., Stemmer, K., Feurstein, D.J., Knobeloch, D., Nussler, A., Dietrich, D.R., 2010. The role of organic anion transporting polypeptides (OATPs/SLCOs) in the toxicity of different microcystin congeners in vitro: a comparison of primary human hepatocytes and OATP-transfected HEK293 cells. *Toxicol. Appl. Pharmacol.* 245, 9–20.
- Flores, E., Herrero, A., Wolk, C.P., Maldener, I., 2006. Is the periplasm continuous in filamentous multicellular cyanobacteria? *Trends Microbiol.* 14, 439–443.
- Flores, E., Herrero, A. 2010. Compartmentalized function through cell differentiation in filamentous cyanobacteria. *Nature Rev. Microbiology* 8, 39–50.
- Fujiki, H., Suganuma, M., 2001. Tumor Promoters – Microcystin-LR, Nodularin and TNF- α and human cancer development. *Anti-Cancer Agents in Medicinal Chemistry* 11, 4–18.
- Gerasimenko, L.M., Orleanskii, V.K., Karpov, G.A., Ushatinskaya, G.T., 2013. Interaction of cyanobac-

- teria with volcanic ashes. *Microbiology* 82, 111–118.
- Giannuzzi, L., Sedan, D., Echenique, R., Andrinolo, D., 2011. An acute case of intoxication with cyanobacteria and cyanotoxins in recreational water in Salto Grande Dam, Argentina. *Mar Drugs* 9, 2164–2175.
- Gordon, N., Angel, D. L., Neori, A., Kress, N., Kimor, B., 1994. Heterotrophic dinoflagellates with symbiotic cyanobacteria and nitrogen limitation in the Gulf of Aqaba. *Mar. Ecol. Prog. Ser.* 107, 83–88.
- Gupta, N., Pant, S.C., Vijayaraghavan, R., Rao, P.V.L., 2003. Comparative toxicity evaluation of cyanobacterial cyclic peptide toxin microcystin variants (LR, RR, YR) in mice. *Toxicology* 188, 285–296.
- Herrero, A., Muro-Pastor, A.M., Flores, E., 2001. Nitrogen Control in cyanobacteria. *Journal of Bacteriology* 183, 411–425.
- Aqaba Holland, A., Kinnear, S., 2013. Interpreting the Possible Ecological Role (s) of Cyanotoxins: Compounds for Competitive Advantage and/or Physiological Aide? *Mar Drugs* 11, 2239–2258.
- Hyvarinen, M., Hardling, R., Tuomi, J., 2002. Cyanobacterial lichen symbiosis: the fungal partner as an optimal harvester. *Oikos* 98, 498–504.
- Jungblut, A.D., Lovejoy, C., Vincent, W.F., 2009. Global distribution of cyanobacterial ecotypes in the cold biosphere. *ISME J* 4, 191–202.
- Kahru, M., Savchuk, O.P., Elmgren, R., 2007. Satellite measurements of cyanobacterial bloom frequency in the Baltic Sea: interannual and spatial variability. *Marine Ecology Progress Series* 343, 15–23.
- Keleti, G., Sykora, J.L., 1982. Production and properties of cyanobacterial endotoxins. *Appl. Environ. Microbiol.* 43, 104–109.
- Kumar, A., Kumar, A., Singh, D.P., Sinha, R.P., Tyagi, M.B., 2006. Eldorado: Hepatotoxicity of *Microcystis aeruginosa* Strains Growing as Blooms in Certain Eutrophic Ponds.
- Kurmayer, R., Gumpenberger, M., 2006. Diversity of microcystin genotypes among populations of the filamentous cyanobacteria *Planktothrix rubescens* and *Planktothrix agardhii*. *Mol. Ecol.* 15, 3849–3861.
- Lagos, N., Onodera, H., Zagatto, P. A., Andrinolo, D., Zevedo, S. M. F. Q., Oshima Y., 1999. The first evidence of paralytic shellfish toxins in the freshwater cyanobacterium *Cylindrospermopsis raciborskii*, isolated from Brazil. *Toxicon* 37, 1359–1373.
- Latysheva, N., Junker, V.L., Palmer, W.J., Codd, G.A., Barker, D., 2012. The evolution of nitrogen fixation in cyanobacteria. *Bioinformatics* 28, 603–606.
- Llewellyn, L.E., 2006. Saxitoxin, a toxic marine natural product that targets a multitude of receptors. *Nat. Prod. Rep.* 23, 200–222.
- Luesch, H., Moore, R.E., Paul, V.J., Mooberry, S.L., Corbett, T.H., 2001. Isolation of dolastatin 10 from the marine cyanobacterium *Symploca* species VP642 and total stereochemistry and biological evaluation of its analogue symplostatins 1. *J. Nat. Prod.* 64, 907–910.
- Lüring, M., 2003. Daphnia growth on microcystin-producing and microcystin-free *Microcystis aeruginosa* in different mixtures with the green alga *Scenedesmus obliquus*. *Limnol. Oceanogr.* 48, 2214–2220.
- MacKintosh, C., Beattie, K.A., Klumpp, S., Cohen, P., Codd, G.A., 1990. Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants. *FEBS Letters* 264, 187–192.
- Magalhães, V.F., Marinho, M.M., Domingos, P., Oliveira, A.C., Costa, S.M., Azevedo, L.O., Azevedo, S.M.F.O., 2003. Microcystins (cyanobacteria hepatotoxins) bioaccumulation in fish and crustaceans from Sepetiba Bay (Brasil, RJ). *Toxicon* 42, 289–295.
- Mankiewicz, J., Tarczynska, M., Walter, Z., 2003. Natural toxins from cyanobacteria. *Acta Biol. Cracov. Ser. Bot.* 45, 9–20.
- Martin, C., Oberer, L., Ino, T., König, W.A., Busch, M., Weckesser, J., 1993. Cyanopeptolins, new depsipeptides from the cyanobacterium *Microcystis* sp. PCC 7806. *J. Antibiot.* 46, 1550–1556.
- Matern, U., Oberer, L., Falchetto, R.A., Erhard, M., König, W.A., 2001. Scyptolin A and B, cyclic depsipeptides from axenic cultures of *Scytonema hofmanni* PCC 7110. *Phytochemistry* 58, 1087–1095.
- Matsunaga, S., Moore, R.E., Niemczura, W.P., 1989. Anatoxin-a (s), a potent anticholinesterase from *Anabaena flos-aquae*. *J. Am. Chem. Soc.* 11, 8021–8023.
- Meeks, J.C., 1998. Symbiosis between Nitrogen-Fixing Cyanobacteria and Plants. *BioScience* 48, 266–276.
- Merel, S., Walker, D., Chicana, R., Snyder, S., Baurès, E., Thomas, O., 2013. State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. *Environ Int.* 59, 303–327.
- Mikalsen, B., Boison, G., Skulberg, O.M., Fastner, J., Davies, W., Gabrielsen, T.M., Rudi, K., Jakobsen, K.S., 2003. Natural Variation in the Microcystin Synthetase Operon *mcyABC* and Impact on Microcystin Production in *Microcystis* Strains. *J. Bact.* 2774–2785.
- Moreira, C., Azevedo, J., Antunes, A., Vasconcelos, V., 2013. *Cylindrospermopsin*: occurrence, methods of detection and toxicology. *J. Appl. Microbiol.* 114, 605–620.
- Mur, R., Skulberg, O.M., Utkilen, H., 1999. Cyanobacteria in the environment. WHO.
- Murakami, M., Ishida, K., Okino, T., Okita, Y., Matsuda, H., 1995. Aeruginosins 98-A and B, trypsin inhibitors from the blue-green alga *Microcystis aeruginosa* (NIES-98). *Tetrahedron Letters* 36, 2785–2788.
- Mynderse, J.S., Moore, R.E., Kashiwagi, M., Norton, T.R., 1977. Antileukemia activity in the *Osillatoria*

- aceae: isolation of Debromoaplysiatoxin from *Lyngbya*. *Science* 196, 538–540.
- Oberholster, P. J., Botha, A.-M., Cloete, t. E., 2005. An overview of toxic freshwater cyanobacteria in south africa with special reference to risk, impact and detection by molecular marker tools. *Biokemistri* 17, 57–71.
- Onodera, H., Oshima, Y., Henriksen, P., Yasumoto, T., 1997. Confirmation of anatoxin-a(s), in the cyanobacterium *Anabaena lemmermannii*, as the cause of bird kills in Danish lakes. *Toxicon* 35, 1645–1648.
- Ortiz-Rodríguez, R., Dao, T.S., Wiegand, C., 2012. Transgenerational effects of microcystin-LR on *Daphnia magna*. *J. Exp. Biol.* 215, 2795–2805.
- Ostermaier, V., Kurmayer, R., 2009. Distribution and abundance of nontoxic mutants of cyanobacteria in lakes of the Alps. *Microb Ecol* 58, 323–333.
- Pablo, J., Banack, S.A., Cox, P.A., Johnson, T.E., Papapetropoulos, S., Bradley, W.G., Buck, A., Mash, D.C., 2009. Cyanobacterial neurotoxin BMAA in ALS and Alzheimer's disease. *Acta Neurol. Scand.* 120, 216–225.
- Paerl, H.W., Fulton, R.S., Moisaner, P.H., Dyble, J., 2001. Harmful Freshwater Algal Blooms, with an Emphasis on Cyanobacteria. *The Scientific World Journal* 1, 76–113.
- Paerl, H.W., Huisman, J., 2008. Blooms like it hot. *Science* 320, 57–58.
- Pearson, L., Mihali, T., Moffitt, M., Kellmann, R., Neilan, B., 2010. On the Chemistry, Toxicology and Genetics of the Cyanobacterial Toxins, Microcystin, Nodularin, Saxitoxin and Cylindrospermopsin. *Mar Drugs* 8, 1650–1680.
- Pitois, S., Jackson, M.H., Wood, B.J.B., 2000. Problems associated with the presence of cyanobacteria in recreational and drinking waters. *International Journal of Environmental Health Research* 10, 203–218.
- Pybus, M.J., Hobson, D.P., Onderka, D.K., 1986. Mass mortality of bats due to probable blue-green algal toxicity. *Journal of Wildlife Diseases* 22, 449–450.
- Rantala, A., Fewer, D.P., Hisbergues, M., Rouhiainen, L., Vaitomaa, J., Börner, T., Sivonen, K., 2004. Phylogenetic evidence for the early evolution of microcystin synthesis. *Proc. Natl. Acad. Sci. U.S.A.* 101, 568–573.
- Reichwaldt, E.S., Ghadouani, A., 2012. Effects of rainfall patterns on toxic cyanobacterial blooms in a changing climate: Between simplistic scenarios and complex dynamics. *Water Res.* 46, 1372–1393.
- Rodríguez, E.M., Acero, J.L., Spoo, L., Meriluoto, J., 2008. Oxidation of MC-LR and -RR with chlorine and potassium permanganate: toxicity of the reaction products. *Water Res.* 42, 1744–1752.
- Rogers, E.H., Hunter, E.S., Moser, V.C., Phillips, P.M., Herkovits, J., Muñoz, L., Hall, L.L., Chernoff, N., 2005. Potential developmental toxicity of anatoxin-a, a cyanobacterial toxin. *J. Appl. Toxicol.* 25, 527–534.
- Rokas, A., 2008. The origins of multicellularity and the early history of the genetic toolkit for animal development. *Annu. Rev. Genet.* 42, 235–251.
- Rounge, T.B., Rohrlack, T., Kristensen, T., Jakobsen, K.S., 2008. Recombination and selectional forces in cyanopeptolin NRPS operons from highly similar, but geographically remote *Planktothrix* strains. *BMC Microbiol.* 8, 141.
- Runnegar, M.T., Kong, S.M., Zhong, Y.Z., Lu, S.C., 1995. Inhibition of reduced glutathione synthesis by cyanobacterial alkaloid cylindrospermopsin in cultured rat hepatocytes. *Biochem. Pharmacol.* 49, 219–225.
- Schirrmeister, B.E., Antonelli, A., Bagheri, H.C., 2011. The origin of multicellularity in cyanobacteria. *BMC Evol. Biol.* 11, 45.
- Schopf, J. W., Packer, B. M., 1987. Early archean (3.3-billion to 3.5-billion-year old) microfossils from Warrawoona group, Australia. *Science* 237, 70–73.
- Schwarzer, D., Finking, R., Marahiel, M.A., 2003. Nonribosomal peptides: from genes to products. *Nat. Prod. Rep.* 20, 275.
- Shin, H.J., Matsuda, H., Murakami, M., Yamaguchi, K., 1997. Aeruginosins 205A and -B, Serine Protease Inhibitory Glycopeptides from the Cyanobacterium *Oscillatoria agardhii* (NIES-205). *J. Org. Chem.* 62, 1810–1813.
- Simmons, T.L., Andrianasolo, E., McPhail, K., Flatt, P., Gerwick, W.H., 2005. Marine natural products as anticancer drugs. *Mol. Cancer Ther.* 4, 333–342.
- Stanier, R.Y., Bazine, G.C., 1977. Phototrophic Prokaryotes: The Cyanobacteria. *Ann. Rev. Microbiol.* 31, 225–274.
- Stewart, I., Webb, P.M., Schluter, P.J., Shaw, G.R., 2006. Environmental Health | Full text | Recreational and occupational field exposure to freshwater cyanobacteria – a review of anecdotal and case reports, epidemiological studies and the challenges for epidemiologic assessment. *Environ Health* 5, 6.
- Tang, E., Tremblay, R., Vincent, W.F., 1997. Cyanobacterial dominance of polar freshwater ecosystems: are high-latitude mat-formers adapted to low temperature? *J. Phycol.* 33, 171–181.
- Ueno, Y., Nagata, S., Tsutsumi, T., Hasegawa, A., Watanabe, M.F., Park, H.-D., Chen, G.-C., Chen, G., Yu, S.-Z., 1996. Detection of microcystins, a blue-green algal hepatotoxin, in drinking water sampled in Haimen and Fusui, endemic areas of primary liver cancer in China, by highly sensitive immunoassay. *Carcinogenesis* 17, 1317–1321.
- van Apeldoorn, M.E., van Egmond, H.P., Speijers, G.J.A., Bakker, G.J.I., 2007. Toxins of cyanobacteria. *Mol. Nutr. Food Res.* 51, 7–60.
- Vyas, K.J., Weiss, J.H., 2009. BMAA – an unusual cyanobacterial neurotoxin. *Amyotroph Lateral Scler* 10 Suppl. 2, 50–55.

- Ward, D.M., Ferris, M.J., Nold, S.C., Bateson, M.M., 1998. A natural view of microbial biodiversity within hot spring cyanobacterial mat communities. *Microbiol. Mol. Biol. Rev.* 62, 1353–1370.
- Weckesser, J., Martin, C., Jakobi, C., 1996. Cyanopeptolins, depsipeptides from cyanobacteria. *Systematic and applied microbiology. System. Appl. Microbiol.* 19, 133–138.
- Welker, M., Döhren, von, H., 2006. Cyanobacterial peptides – Nature’s own combinatorial biosynthesis. *FEMS Microbiology Reviews* 30, 530–563.
- WHO, 2003. Cyanobacterial toxins: Microcystin-LR in drinking water. 1–18.
- Wiegand, C., Pflugmacher, S., 2005. Ecotoxicological effects of selected cyanobacterial secondary metabolites: a short review. *Toxicol. Appl. Pharmacol.* 203, 201–218.
- Wilson, A.E., Sarnelle, O., Tillmanns, A.R., 2006. Effects of cyanobacterial toxicity and morphology on the population growth of freshwater zooplankton: meta-analyses of laboratory experiments. *Limnol Oceanogr.* 51, 1915–1924.
- Zeller, P., Clément, M., Fessard, V., 2011. Similar uptake profiles of microcystin-LR and -RR in an in vitro human intestinal model. *Toxicology* 290, 7–13.
- Zurawell, R.W., Chen, H., Burke, J.M., Prepas, E.E., 2005. Hepatotoxic Cyanobacteria: A Review of the Biological Importance of Microcystins in Freshwater Environments. *Journal of Toxicology and Environmental Health, Part B* 8, 1–37.
- Žegura, B., Zajc, I., Lah, T.T., Filipič, M., 2008. Patterns of microcystin-LR induced alteration of the expression of genes involved in response to DNA damage and apoptosis. *Toxicol.* 51, 615–623.

Verena Grundler and Karl Gademann
Department of Chemistry
University of Basel
St. Johanns-Ring 19
4056 Basel

