

# **Terebratella indica sp. n. representing a new genus of Tetrastichinae (Hymenoptera : Eulophidae) from Muzaffarpur, India**

Autor(en): **Shafee, S. Adam / Rizvi, Seema**

Objektyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society**

Band (Jahr): **57 (1984)**

Heft 4: **Festschrift Prof. P. Bovey**

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-402135>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

*Terebratella indica* sp. n. representing a new genus of Tetrastichinae  
(Hymenoptera: Eulophidae) from Muzaffarpur, India

S. ADAM SHAFEE & SEEMA RIZVI

Department of Zoology, Aligarh Muslim University, Aligarh, India

*Terebratella* gen. n. is defined. It is a unique genus of Tetrastichinae in having a very long ovipositor. *T. indica* sp. n. is described and illustrated.

*Terebratella* gen. n.

Female

*Head* slightly wider than long in facial view; frontovertex twice as wide as long; malar space as long as eye width; mandibles with two teeth and a truncation; antennae with funicle 3-segmented, club 2-segmented, ring segments present; pedicel shorter than first funicle segment; mesoscutum with well developed parapsidal furrows, scutellum with two sublateral longitudinal grooves; fore wings with marginal vein very long, postmarginal vein rudimentary, stigmal vein well developed; legs long and slender, tarsi 4-segmented; abdomen sessile, distinctly longer than thorax; ovipositor very long, much longer than abdomen.


Type-species: *Terebratella indica* sp. n.

Comment: The new genus is assigned under the subfamily Tetrastichinae on the basis of having a mesoscutum with complete parapsidal furrows, a scutellum with two sublateral longitudinal grooves, and fore wings with a very long marginal vein, a rudimentary postmarginal vein, and a well developed stigmal vein. In the key to the genera of Tetrastichinae proposed by BOUCEK (1977), the new genus runs near *Tetrastichodes* ASHMEAD, but differs from that genus and the remaining genera of Tetrastichinae in having a very long ovipositor which is much longer than the abdomen.

*Terebratella indica* sp. n.

Female (Figs. A-C)

*Head* orange yellow, except a patch on dorsum, between eyes dark, slightly wider than long in facial view; frontovertex twice as wide as long; ocelli red, arranged in obtuse triangle, lateral ocellus separated by about twice its diameter from inner orbital margin and about thrice its diameter from occipital margin; eyes red; malar space as long as eye width, malar sutures present; antennae inserted above lower level of eyes, inter-antennal space about one-third the width of frons between eyes at median ocellus; mandibles with two teeth and a truncation. Antennae (fig. A) dark brown; scape slightly more than three times as long as wide; pedicel one and a half times as long as wide, less than one-half the length of



Figs. A-C: *Terebratella indica* sp. n., ♀: (A) Antenna; (B) Fore wing; (C) Abdomen with ovipositor.

first funicle segment; three ring segments present; funicle 3-segmented, first longest, slightly less than four times as long as wide, third slightly more than two times as long as wide; club 2-segmented, three and a half times as long as wide, slightly shorter than preceding two funicle segments together.

*Thorax* with metallic bluish green reflections; mesoscutum long with complete parapsidal furrows, scutellum with two sublateral longitudinal grooves; propodeum well developed. Fore wings (fig. B) hyaline, slightly less than three times as long as wide; costal cell narrow, slightly more than one-half the length of marginal vein; marginal vein very long with 13 long setae, postmarginal vein rudimentary, stigmal vein well developed; marginal fringe short, spaced by a distance equal to one-fifth their length. Legs orange yellow except pretarsus of all legs brown, hind coxae dark brown; long and slender; tarsi 4-segmented.

*Abdomen* dark except ventro-lateral margins yellow, much longer than thorax; ovipositor (fig. C) very long, longer than length of abdomen, arising from base of abdominal venter; subgenital plate reaching basal one-third of abdominal venter.

*Body* length excluding exerted part of ovipositor: 1.87 mm. Length of exerted part of ovipositor: 1.36 mm.

*Holotype* ♀. India: Bihar, Muzaffarpur, Jhapa, 5.XI.1969 (S. ADAM SHAFEE)  
Type deposited in Zoological Museum, Aligarh Muslim University, Aligarh, India.

#### ACKNOWLEDGEMENTS

The authors are deeply indebted to Prof. NAWAB H. KHAN, Chairman, Department of Zoology, Aligarh Muslim University, Aligarh, for providing research facilities. One of us (S. RIZVI) is thankful to U.G.C., New Delhi, for providing financial assistance.

#### LITERATURE

BOUCEK, Z. 1977. *Descriptions of Tachinobia gen. n. and three new species of Tetrastichinae (Hymenoptera. Eulophidae), with a tentative key to genera.* Bull. ent. Res. 67: 17-30.

(received June 14, 1984)