

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 40 (1960)

Heft: 2

Artikel: A note on the optical properties of strychnine

Autor: Rittmann, A. / El-Hinnawi, E.E.

DOI: <https://doi.org/10.5169/seals-31157>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A Note on the Optical Properties of Strychnine

By *A. Rittmann*¹⁾ and *E. E. El-Hinnawi*²⁾

With 1 figure in the text

Crystal-optical properties of strychnine $C_{21}H_{22}N_2O_2$ have been reported by MAYRHOFER (1928), KOFLER (1936) and WINCHELL (1954). These data include only the indices of refraction, for which, each author has given certain values. Data on the optical sign and the optical axial angle were not recorded, except for a merely qualitative expression given by WINCHELL (1954).

In what follows, the writers give the complete optical data they have measured on strychnine crystals grown on a glass carrier and also a discussion of the data given by previous authors.

Crystals of strychnine are colourless, elongated six sided, rarely about equant, with profile angles of 125° , 125° and 110° (Fig. 1). Using the immersion method, the indices of refraction N_z and N_x have been found to be 1.720 and 1.621 respectively. By calculation N_y has been found to be 1.672 and $N_z - N_x = 0.099$.

Between crossed nicols, the crystals are anisotropic with parallel extinction and positive elongation.

The conoscopic image is monosymmetric. The quotient of the extreme retardations $Q = 0.736$ while the quotient of the intermediate retardation and the maximum $Q' = 1.2$, from which the optic axial angle $2V$ has been found be -87° . The Crystals are orthorhombic, lying on a prismatic face.

From a comparison of the above data with those recorded in the literature (Table 1), the following facts emerge.

Acknowledgements. One of us (*E. E. El-Hinnawi*) wishes to express his deep appreciation to the National Research Centre and the Mission Bureau, Cairo for granting him a study leave at the Vulcanological Institute of Catania University, Italy.

¹⁾ Istituto di Vulcanologia dell'Università Catania, Sicilia.

²⁾ Present Adress: National Research Centre, Dokki, Cairo, Egypt.



Fig. 1. Photomicrograph of strychnine crystals lying on a glass carrier. Crossed Nicols, $42.5\times$.

Table 1.

	MAYRHOFER (1928) ³⁾	KOFLER (1936) ⁴⁾	WINCHELL (1954) ⁵⁾	Present data
Nz	1.730	\sim 1.740	1.74	1.720
Ny	1.631	1.653	1.68	1.672
Nx	1.620	1.612	1.61	1.621
Nz—Nx	0.110	\sim 0.128	0.13	0.099
sign	+ ?	+ ?	+ ?	- ve
2V	very high	very high	very high	87°

References

- MAYRHOFER, A. (1928): Mikrochemie der Arzneimittel und Gifte. Bd. 2. Berlin.
 KOFLER, A. (1936): Mikroskopische Methoden in der Mikrochemie. Emil Haim & Co. Leipzig.
 WINCHELL, A. N. (1954): Optical properties of organic compounds. 2nd Ed. Academic Press, New York.

Istituto di Vulcanologia dell'Università Catania, February 1960.

³⁾ The data of MAYRHOFER do not include the optic axial angle, furthermore the optic sign is doubtful. From the indices of refraction as measured by him, it appears that the optic sign is positive, while a calculation of the optic axial angle based on these indices would give an angle of about $+37^\circ$ which is surely not very high!

⁴⁾ The data of KOFLER lack an accurate determination of Nz. In addition the optic sign is doubtful and the optic axial angle undeterminable.

⁵⁾ The data recorded as WINCHELL also remain without a determination of the optic sign and angle, although a calculation of 2V based on the given indices of refraction points to a high negative optic angle. However the indices Nz and Ny are overestimated while Nx is lower than the value recorded in the present investigation.