Zeitschrift:	Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band:	54 (1974)
Heft:	1
Artikel:	On calcic amphiboles and amphibolites from the Lepontine Alps
Autor:	Wenk, E. / Schwander, H. / Stern, W.
DOI:	https://doi.org/10.5169/seals-42191

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 01.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On Calcic Amphiboles and Amphibolites from the Lepontine Alps

By E. Wenk, H. Schwander and W. Stern (Basel)*)

With 24 figures in the text and 14 tables in appendix I

Summary

Spectrographic, XF- and microprobe-analyses of 56 calcic amphiboles from amphibolites and related rocks of the Central Alps, 15 coexisting biotites and 35 host rocks are presented. In the area considered mainly rocks in amphibolite facies are exposed; transitions to greenschist facies occur in the marginal parts. The metamorphic grade of amphibolites is judged in accordance with the anorthite content of the coexisting plagioclase. Fabric criteria and parageneses of amphibolite facies rocks indicate close approach to equilibrium conditions during the main phase of Tertiary regional metamorphism.

In amphibolite facies rocks common hornblende, rarely associated with actinolite, shows limited chemical variation, in spite of the considerable and regular compositional change of the coexisting plagioclase from high to low grade metamorphic zones. Mg increases in amphiboles as well as in amphibolites with higher grade and there is clear interdependence between the Mg/Fe ratios of hornblende, biotite and host rock. The mean values for Fe₂O₃, Na₂O and H₂O of the amphibole appear to decrease with higher grade, but the standard deviations are considerable. The bulk of common hornblendes is concentrated between Al^{tot} 1.75 and 2.4. Our data for amphibolite facies hornblendes indicate a compositional gap in the range Al^{tot} 1 to Al 1.75. In greenschist facies rocks, however, a miscibility gap is not evident. Tschermakitic hornblendes with Al^{tot} > 2.5 are of common occurrence in the schist-belt south of Gotthard and Lukmanier passes.

In amphibolite facies rocks the pronounced increase in anorthite content, reflecting progressive metamorphism, is performed mainly by modal variation. The quantities of Na, Ca, Si and Al remain fairly constant in the system (though not in greenschists). Plagioclase has to consume the major part of Na, hornblende the minor part. With rising anorthite content of plagioclase the balance can only be preserved by the formation of more feldspar, mainly at the expense of epidote-clinozoisite, chlorite and biotite, and in later stages also at the expense of hornblende. Modal analyses and calculations from rock and mineral analyses both show that andesine-amphibolites contain twice as much plagioclase as albite-amphibolites.

^{*)} Authors address: Mineralogisch-Petrographisches Institut, Bernoullianum, CH-4056 Basel.

Index

1. Introduction		98
2. Methodological considerations	10	01
2.1. Chemical investigations		01
2.1.1. X-ray microprobe		01
2.1.2. Analyses of main constituents with X-ray fluores	cence 10	02
2.1.3. Spectrographical trace element analysis	10	02
2.2. Physical investigations	10	02
$2.2.1. Density \dots \dots$		02
2.2.2. Index of refraction		02
2.2.3. Modal analysis	10	02
3. Results	10	03
3.1. Calcic amphiboles s. l		03
3.2. Trioctahedral micas		09
3.3. Host rocks; new and literature data of some Lepontine a	mphibolites 1	10
4. Discussion	11	10
4.1. Correlation between chemical and physical data of calci	c amphiboles 1	10
4.2. Chemical relations between coexisting minerals and host	$t rocks \dots 1$	11
4.3. Relationship between grade of metamorphism, fabric, m	ode, An-content of	
plagioclase and composition of amphibolite and calcic a	mphibole 11	17
4.3.1. Fabric		17
4.3.2. Number of mineral phases		18
4.3.3. Mode and anorthite content of plagioclase		18
4.3.4. Rock composition	12	20
4.3.5. Composition of calcic amphibole		22
4.3.6. Distribution of Or Ab An between coexisting pla	gioclase and horn-	a -
Diende	12	25 90
	12	49
Appendix I	18	30
Bibliography	14	4 8

1. Introduction

During the past twelve years many authors have investigated the regional distribution of rock-forming minerals, their physical and chemical properties and parageneses in the Central Alps, especially in the Canton Ticino and adjacent parts of Italy and in the Cantons Valais and Grisons. In this wide area of the Lepontine Alps, covering about 7500 km², rocks mainly in the

amphibolite facies are exposed; transitions to greenschist facies occur in the marginal part of the area. The whole region is by now well subdivided by isograds of different types which form a concentric system, imprinted during various phases of Alpine regional progressive metamorphism, culminating in the Oligocene-Miocene.

Metamorphic carbonate-rocks, aluminous schists and ultramafic rocks have been studied extensively, and this highly rewarding research is still in progress. It appeared desirable to extend these investigations to the basic rocks after which the metamorphic facies are named: amphibolites and greenschists. These usually form extended zones of moderate thickness and, although minor in importance in comparison to gneisses and micaschists, they serve as geological key horizons. Amphibolites should be useful also as indicators of metamorphic grade. Often they are composed of only two main constituents and a few accessories, and they show a regular equidimensional fabric indicating close approach to equilibrium conditions.

Paragenetic studies require rich and representative collections. The first author started sampling in the Central Alps many years ago, and in 1969 E. WENK and F. KELLER reported on the results of modal anorthite determinations in 700 different amphibolites, listed in the appendix of the cited paper. These investigations outlined concentric zones of albite-, oligoclase-, andesine- and labradorite-amphibolites indicating progressive metamorphism. The highest grade rocks are confined to structural steep zones, especially to the so called root zone in the south and to the Maggia zone in the centre. The general pattern of these plagioclase isograds is in good agreement with the plagioclase isograds observed in silicate-marbles and calc-schists of the same area. The zones of highest and of lowest metamorphic grade coincide, only the absolute An-values differ. This indicates that there are no profound differences between the patterns of plagioclase isograds developed in CO_2 -free rocks and those found in series strongly influenced by CO_2 partial pressure.

The present study is mainly concerned with questions of hornblende composition, modal and chemical variation of the host rock. It relies upon the collections described by WENK and KELLER (1969), augmented by further sampling (see sample maps Fig. 1). The responsibility for the extraction of mineral concentrates, for the chemical analyses and for parts 2 to 4.2 of the text rests with the second and third authors and their collaborators. The cell parameters of most of the analysed amphiboles, and a discussion of their variation with chemical composition, have been published by H. R. WENK (1971).

A review of existing literature reveals little compositional variation of common hornblende in amphibolites of different grade, in spite of the given possibilities of complex crystallochemical substitution. The variation is controlled mainly by rock composition (especially Mg/Fe-ratio). Published evidence



Fig. 1a. Samples, Amphiboles s. l.



Fig. 1b. Samples, Amphibolite + Hornblende bearing rocks.



Fig. 1c. Samples, Host rock + Amphibole.

ascribing minor variations in hornblendes to metamorphic grade is in part consistent, in part conflicting. Also our contribution dealing with this complex subject leaves room for further discussion. We have the feeling that scientists often ask questions that nature does not pose, or to which our instrumental possibilities cannot reply. Obviously it seems more promising to concentrate upon those topics where clear statements can be made. Our discussion also deals with the arising difficult problem: How does plagioclase manage to change its anorthite content so well in response to metamorphic grade, even in rocks of similar bulk composition and without compensating change in hornblende composition?

Reference samples of the rocks and minerals described are in the collections of the Department of Mineralogy of the University of Basel (Switzerland). Research on these materials is still in progress by A. Steck (Lausanne, ore minerals), R. Steiger (Zürich, age determination of hornblendes) and H. Ulbrich (Basel, tschermakitic hornblendes of the Gotthard road-tunnel).

2. Methodological considerations

2.1. CHEMICAL INVESTIGATIONS

2.1.1. X-ray microprobe

Polished thin sections of amphibolites were investigated with a Jeol probe (for excitation conditions see table 1). Sample preparation was carried out with special care:

First grinding up to a gradation of emery-1000 and a slide thickness of 30 microns, subsequently polishing by hand on Metcloth 40-7158 support with one micron diamond paste, or on a Depiereux apparatus. Certain rocks had to be hardened prior to polishing with epoxy resins (araldite or Z 70 C).

After cleaning with brush and detergents, the thin section was coated first with a few angstroms of carbon, thereon with approximately 200 Å of gold (evaporation of a constant quantity of Au).

The use of different silicate minerals as reference samples causes generally different interelemental effects; working with low excitation conditions (as low as possible for an old type instrument) these elemental interactions may be reduced, and sample contamination as well. The lower limit of those excitation conditions is given by the counting statistics.

The overall reproducibility of our measurements has been checked by testing chemically homogeneous plagioclase and amphibole samples (e. g. Cal 21). The variation found corresponds reasonably well with the expected error of the probe. Thus, the quality of our thin sections seems to be sufficient.

2.1.2. Analysis of main constituents with X-ray fluorescence

Details of sample preparation and analyzing techniques have been described in a previous paper (STERN, 1972). Most of the analyses listed below have been executed according to these procedures, a few by means of light optical emission spectrometry (SCHWANDER et al., 1968).

Mineral separation was performed mainly according to STERN (1966). Due to frequent intergrowths of amphiboles, especially from low grade amphibolites, minerals became separated out of small siftings (grain diameter 0.1 to 0.2 mm).

In certain cases, no pure hornblende concentrates could be obtained; these samples with 2 to 10% of inclusions and intergrowths were labelled as "concentrates".

2.1.3. Spectrographical trace element analysis

Some trace elements were analyzed by means of light optical emission spectrography according to SCHWANDER et al. (1968).

2.2. PHYSICAL INVESTIGATIONS

2.2.1. Density

The density of the hornblende samples was determined using a 10 ml bottle, Gay-Lussac type, with a mixture of 10% alcohol in distilled water.

2.2.2. Index of refraction

The refractive indices of amphiboles have been determined on grains oriented with the aid of an Eulerian cradle. The microgoniometer/cuvette assemblage, developed and described by STECK and GLAUSER (1968), was used for the determinations listed in this paper.

2.2.3. Modal analyses

The mineral modes have been determined mainly with a point counter on the base of 2000 grains per thin section, step width 0.25 mm. Each slide was scanned in the two main directions; the accuracy of these determinations is approximatively $\pm 2\%$ rel.

The modes of some amphibolite samples were estimated; the accuracy may be assumed to reach $\pm 5\%$ rel.

3. Results

The bulk of analytical data is listed below, in tables 2–4, 6 and the evaluation follows in chapter 4.

Part of the analytical information had to be recalculated: structural formulae of minerals, QLM values of rocks, etc., have been computed by a Diehl Combitron calculator (program input with punch tape). Part of the programs had been obtained as software from Diehl Ltd. (regression calculations, student t test), another part has been set up by us.

3.1. CALCIC AMPHIBOLES (Tables 2, 6, Figures 2, 3)

With one exception (Bni 258) all analyses presented in tables 2 and 6 belong to the group of calcic amphiboles. Most of these monoclinic amphiboles from the Central Alps fit the conditions $(Na + K)_A < 0.5$ and Ti < 0.5, only in 13 cases is $(Na + K)_A > 0.5$. In fig. 21 the common hornblendes from amphibolites are concentrated in the field Al^{tot} 1.75–2.4, while actinolites and tremolites from silicate-marbles and mafic lenses found in the same area, occupy the corner Al^{tot} < 1. Accordingly our analyses have been grouped in the two tables 2a and 2b. Hornblendes with high alumina contents tending towards tschermakite, occur in a schist belt (garnet-hornblende-Garbenschiefer) which includes also some true amphibolites and which is localised in a zone south of the Gotthard massif (rock group III, Tremola, Frodalera). For these tschermakitic



Fig. 2a. Calcic amphiboles, plotted after LEAKE (1968): Ca+Na+K<2.50; Ti<0.50. New data: big circles; literature: small circles.

I Tremolite

- III Actinolite
- V Mg-hornblende
- VII Tschermakite
- IX Ferro-actinolitic hornblende
- XI Ferro-tschermakitic hornblende
- II Tremolitic hornblende
- IV Actinolitic hornblende
- VI Tschermakitic hornblende
- VIII Ferro-actinolite
- X Ferro-hornblende
- XII Ferro-tschermakite



Fig. 2b. Calcic amphiboles, plotted after LEAKE (1968): Ca+Na+K>2.50; Ti<0.50.

I Edenite

- **III** Pargasitic hornblende
- V Ferroan pargasitic hornblende
- VII Ferro-edenite
- IX Mg-hastingsitic hornblende
- XI Hastingsitic hornblende
- IV Pargasite VI Ferroan pargasite
- VIII Ferro-edenitic hornblende

II Edenitic hornblende

- X Magnesian hastingsite
- XII Hastingsite

amphiboles which are included in tables 2a and 6, ANGEL (1967) proposed the name "gotthardite".



Fig. 3a. QLM-plot of calcic amphiboles and amphibolites. New data: small dots = calcic amphiboles; black rectangles = actinolitic/tremolitic hornblendes; big dots = amphibolites/hornblende bearing rocks. Literature: open circles = amphibolites/hornblende bearing rocks.



Fig. 3b. Kp-Ne-Cal plot; for explanation of symbols see fig. 3a.



Fig. 3c. Fo-Fa-Cs plot; for explanation of symbols see fig. 3a.

While in the Ticino area proper, a clear miscibility gap exists between tremolite and common hornblende, the data presented by WETZEL et al. (1973) from lower grade metamorphic series adjacent to the W indicate miscibility (actinolitic hornblendes) and a trend toward edenitic and pargasitic compositions with $(Na + K)_A > 0.5$.

The analyzed calcic amphiboles are graphically presented on Figs. 2 and 3; some crystal chemical data are plotted on Figs. 4, 5 showing the distribution of octahedral sums (mean approximatively 5.00, probably more than one normal distribution) and of alkali sums (mean ± 2.25 , normal distribution).

The chemical homogeneity of the amphiboles was tested by microprobe. Spot analyses were taken along lines perpendicular and parallel to c of selected hornblende crystals; line scanning was not possible due to low sensitivity and



Fig. 4. Octahedral sum of calcic amphiboles; N = 60 new data, 90 from literature (triangles).



Fig. 5. Alkali sum of calcic amphiboles: normal distribution; N = 65.

the probable influence of topographical effects. Amphiboles from high grade amphibolites with well developed mosaic structure, showed a homogeneity corresponding with the relative error of the probe – point analyses along the mentioned lines showed the same variability as the respective averages of different mineral grains in the same slide:

1 to 2% rel. in case of SiO₂, 3 to 5% rel. in case of Al₂O₃, FeO_{tot}, MgO and CaO, 5 to 10% rel. in that of Na₂O, K₂O, see table 6. In these amphibolites it seems, that from the chemical standpoint, only one type of amphibole exists.

In the zone of low grade metamorphism however, especially in the marginal part of the Lepontine Alps, rather inhomogeneous hornblende patterns occur; spot analyses on one and the same crystal prove a better homogeneity – though being zoned – than the averages of different grains of the same thin section. Thus certain amphibolites of the lower metamorphic margin seem to contain at least two types of calcic amphiboles showing especially high variations of Si and Al and having high Si/Al-ratios. Whether the possible chemical variability of those amphiboles could even be greater or not, may not be decided on the base of the present data. If there exists furthermore an interdependence between chemical composition and casual zonality or domain texture should be tested with these chemically inhomogeneous amphiboles. This investigation should however be done by means of a new type microprobe with high geometrical resolution.

Microprobe analyses of calcic amphiboles and their structural formulae are listed on Table 6, the octahedral cations are plotted on Fig. 6.



Fig. 6. Microprobe analyses of calcic amphiboles; tie lines between corresponding spot analyses on one crystal. Mg-Fe-Al^{VI} plot of octahedral cations.

The number of analyses being rather high -106 amphibolites including literature values, and 86 calcic amphiboles of the Central Alps - a statistical evaluation of the data seemed to be of interest. The analyses were grouped according to petrographical criteria, i. e. to the optically determined An content of plagioclase in amphibolites (see Table 10), and the respective averages, standard deviations and variations of every main constituent were calculated (Table 11a, b).

From these data, student – t values were computed and the probabilities estimated for the chemical difference of the respective petrographical groups (see Fig. 7). It is obvious that the analyzed calcic amphiboles show clearer differences than the corresponding amphibolite groups: sodium, magnesium, aluminum, silicon (the latter in groups I, III only) in the case of calcic amphiboles, and calcium and H_2O^+ only in the case of amphibolites seem to have a certain diagnostic meaning in the Lepontine Alps, if any. Considering the often large variances in between these petrographical groups, one should not exaggerate the meaning of a single criterion.



Fig. 7a. Mean values of chemical components (amphiboles, amphibolites), for comparison see table 11.

	SiO ₂ G H	А і 2 ⁰ 3 G Н	Fe ₂ 0 ₃ G H	FeO G H	Mg0 GH	Са О G Н	Na ₂ 0 G Н	к ₂ 0 G Н	тіО ₂ G H	н ₂ 0 G Н
0/I II IV V VI	•	0 •	0	•		0	0 •		•	© ● ● ● ●
I / II IV V VI	0	•		0	• • •	• • •	0 0	•00	● 0	• •
II / IV V VI	0						•0			•
IV/V VI		0		٥		•	0 0	0	•	:
V/ VI	0	0				٥	0			
III / O I II IV V VI			I	 ● ○				• •		

Fig. 7b. Probabilities estimating the chemical differences between petrographical groups.

• 99.6–99.99% probability ⊙ 98.5–99.5

G rocks

○ 95 -98

H amphiboles s. l.

For roman ciphers see p. 144. Sources of chemical data: DE QUERVAIN et al. 1956, LEAKE 1968, STEIGER 1961, DIETRICH 1969, WEIBEL 1964; and analyses listed on tables.

3.2. TRIOCTAHEDRAL MICAS (Table 3, Fig. 8a-c)

From some amphibolites trioctahedral micas have been separated and analyzed as well. Moreover, published data (WENK et al., 1963) have been added on micas coexisting with hornblendes discussed in this paper.



Fig. 8a. QLM plot hornblende-mica.

Fig. 8b. Kp-Ne-Cal hornblende-mica.



Fig. 8c. Fo-Fa-Cs/Fo-Fa-C plot of coexisting trioctohedral micas and calcic amphiboles.

3.3. HOST ROCKS; NEW AND LITERATURE DATA OF SOME LEPONTINE AMPHIBOLITES (Table 4, Figures 9a-c)

Only for 21 of the analyzed amphiboles can we present analytical data of the respective host rocks; on the other hand, we list a number of amphibolite analyses without corresponding data on hornblendes.

Recent published data on amphiboles and amphibolites from the Lepontine Alps are rather scarce. It seems moreover necessary to omit most of the early analyses collected and published by DE QUERVAIN et al. (1956), because of their questionable sodium values.

4. Discussion

4.1. CORRELATION BETWEEN CHEMICAL AND PHYSICAL DATA

The ratio Mg/Mg + Fe + Mn represents not only the basis of classification of the structural formulae (Table 5), but shows positive correlation with some physical parameters, such as density and refractive index (Fig. 10–12). Insofar as both specific weight and index of refraction are related to the chemical character of a calcic amphibole, density itself seems to be connected linearly with the optic parameters (see Fig. 13).

The optic parameters show a similar trend for n_{α} and n_{γ} , both rising parallelly, whereas $2V_x$ decreases at the same time. This phenomenon may be explained by the fact, that $n_{\beta}-n_{\alpha}$ increases generally with higher values, whereas $n_{\gamma}-n_{\beta}$ diminishes (see Fig. 10).



Fig. 9a/b/c. QLM plots of calcic amphiboles and host rocks. Small black dots = calcic amphiboles; big black dots = host rocks.

4.2. CHEMICAL RELATIONS BETWEEN COEXISTING MINERALS AND HOST ROCKS

From the two coexisting minerals hornblende s. l. and trioctahedral mica, only one main chemical constituent – magnesium – shows a clear interdependence between not only the two mineral phases (Fig. 14), but between horn-



Fig. 10. Optical parameters of calcic amphiboles; grouping according to rising values of n_{α} .



Fig. 12. Density of calcic amphiboles. Mg-Fe-Al^{VI} plot of octohedral cations.

1	D < 3.00	2	3.01 - 3.05
3	3.06 - 3.10	4	3.11 - 3.15
5	3.16 - 3.20	6	3.21 - 3.25
7	3.26 - 3.30	8	$> \! 3.31$

.



Fig. 11. Density, and optical parameters vs mg (Mg/Mg+Fe_{tot}+Mn) of calcic amphiboles. Triangles: tremolitic/actinolitic hornblendes; open circles: common hornblendes; crosses: "gottharditic" hornblendes. Calculated regressions: full lines for common and trem./act. hornblendes, broken lines for "gottharditic" hornblendes.



Fig. 13. Density vs n of calcic amphiboles.



Fig. 14. mg (Mg/Mg + Fe'' + Fe'' + Mn) of coexisting calcic amphiboles and trioctohedral micas.



Fig. 15. MgO and FeO distribution (wt-%) between hornblende s. l. / host rock, and hornblende s. l./coexisting mica. r = coefficient of correlation.



Fig. 16. Al₂O₃ and MgO (wt-%) of calcic amphiboles s. l. (new data). Black dots: microprobe analyses; circles and triangles: XFA.

blende and host rock as well (Fig. 15). All other elements participate in other mineral phases occurring in variable modes. Consequently, the chemical correlation of these hornblendes with their host rocks is rather poor, except for Mg. It may be noted, however, that the total of analyzed amphibole/host rock pairs is still fairly limited.

From Fig. 7 chemical differences between rock groups of different metamorphic grade can be estimated: calcic amphiboles of group III differ from all other groups especially with respect to Al and Mg. Fig. 16 with Al_2O_3 vs MgO



Fig. 17. Wt-% FeO vs Fe₂O₃ of coexisting calcic amphiboles and trioctohedral micas.

(wt-%) shows consequently a well defined group of "gottharditic" hornblende corresponding with group III, whereas common amphiboles and actinolitic/ tremolitic hornblendes are not separated by a clear gap.

Fig. 17 informs about oxidation relations of calcic amphiboles and coexisting trioctahedral micas: the tie-lines between mineral pairs show two distinct trends: the ratio wt-% $Fe_2O_3/wt-\%$ FeO of hornblendes s. l. is in 70% of the cases higher than the one of coexisting trioctahedral micas. Note, however, that the vertical Fe_2O_3 scale is enlarged 2.5 times; thus, the analytical error of wet chemical FeO determinations appears enlarged as well.

The relations of coexisting micas and amphiboles are plotted on Fig. 8. Common and actinolitic/tremolitic hornblendes occupy distinct fields on the QLM plot, whereas most of the trioctahedral micas – biotites and phlogopites as well – are situated in a single, rather narrow field. Consequently, there are two different slopes of tie-lines between coexisting mineral pairs.

Obviously, trioctahedral micas can not be plotted on a Fo-Fa-Cs diagram (8c) because of their high aluminum and low calcium content. Their M relations were, therefore, calculated in a Fo-Fa-C triangle and connected – in a slightly unusual way – with their respective coexisting amphiboles.

4.3. RELATIONSHIP BETWEEN GRADE OF METAMORPHISM, FABRIC, MODE, An-CONTENT OF PLAGIOCLASE AND COMPOSITION OF AMPHIBOLITE AND CALCIC AMPHIBOLE

The following discussion concentrates upon common amphibolites with calcic hornblende and plagioclase as predominant constituents (75-95% vol), and with clinopyroxene, epidote-clinozoisite, biotite, chlorite or quartz as additional components. Sphene, rutile, iron ore, and rarely calcite are accessory minerals. Orthopyroxene is absent. Scapolite-, anthophyllite- and alkali-amphibole-bearing rocks are excluded. Also garnet-amphibolites and eclogites, both of which are rare in the Lepontine Alps, are not considered. This concentration upon the main rock types which form more than 95% of the total basic rocks occurring in the region investigated, infers that our conclusions pertain only to amphibolites, with a range of plagioclase from albite to labradorite.

Hornblende-rich rocks with plagioclases more calcic than An 60 are rare and our data insufficient. The transition zone from albite-amphibolites to hornblende-bearing greenschists (prasinites in the definition of KALKOWSKY 1886) is discussed by WETZEL (1973) and will be mentioned only occasionally.

At the western and northeastern margin of the area investigated (see Fig. 1) there is clear evidence that several zones of the Mesozoic ophiolitic suite in greenschist facies change into amphibolite facies rocks. However, in the higher grade central part of the Lepontine Alps few geological facts, e. g. the association of amphibolites with long stretched bands of silicate-marbles and calcschists (Mesozoic Bündnerschiefer) indicate an ophiolitic parentage. The origin of all those amphibolites which are intercalated in the vast gneiss complexes is not known (Mesozoic ophiolites, older basic rocks, or restites in Alpine migmatites). The only proven fact is that all these Lepontine rocks recrystallized and were deformed during the Alpine orogenesis. Whatever the early history of these rocks of basaltic composition may be, they now form a correlated metamorphic series.

4.3.1. Fabric

With rising grade of metamorphism the fabric of the basic rocks becomes more regular with regard to grain size, shape and grain boundaries (compare Fig. 4 in WENK and KELLER 1969). Sieve-structured porphyroblasts and diablastic intergrowths disappear, and a mosaic results, made up of polygonal or interlocking crystals of hornblende and plagioclase of approximately equal size. In these advanced stages of progressive regional metamorphism the recrystallization is often accompanied by mineral sorting, producing compositional layering parallel to the foliation; but the individual layers show regular equidimensional structure. All amphibolites show distinct foliation and lineation parallel to the Alpine fold axes.

4.3.2. Number of mineral phases

From the zone of albite – to the zone of andesine – amphibolites the number of mineral phases decreases. Thus, often in the latter, only two main constituents, viz. a homogeneous hornblende and zoned andesine occur. With the appearance of clinopyroxene at higher metamorphic grade a new phase is added.

4.3.3. Mode and anorthite content of plagioclase

The results of modal analyses and of anorthite determinations in 85 plagioclase-amphibolites from the Central Alps (65 sets of data refer to our samples, 20 are taken from the literature) were plotted in a large scale diagram, with modal composition as the vertical and An-range as the horizontal axis. In this detailed master diagram each mineral component of the 85 rocks was represented by a straight line at the modal amount given, the length of the line indicating the An-variation of the feldspar. From this original diagram average



Fig. 18. Average modes of amphibolites with plagioclase ranging from An 0 to An 60. The diagram shows that the amount of plagioclase increases with rising anorthite content.

modes were derived for the An-intervals listed in Table 12, and represented in Fig. 18. The evidence offered in this diagram is well founded in the albite range and from An 20 to 60; it is less reliable within the peristerite gap and at An > 60 (dotted lines). The modal averages considered refer to rock-types of similar composition, and the mean values give rough information on the variable volumes of the mineral phases in relation to changing An-content of plagioclase.

Fig. 18 shows that from the peristerite gap on, the amount of plagioclase increases, concomitant with the rise in anorthite content, indicating higher metamorphic grade. Andesine-amphibolites carry on average twice as much feldspar as albite-amphibolites, mainly at the expense of clinozoisite-epidote. The quantity of hornblende remains almost constant in this range: from An 40 on it decreases in favour of pyroxene and plagioclase. At An-values higher than 60 our evidence is insufficient. Such calcic rocks have compositions different from common amphibolites. Our Fig. 18 agrees well with Fig. 2 in ENGEL et al. (1964) and includes additional information on lower-grade rocks.

The main result of the modal analyses is the considerable increase of the volume of feldspar with rising anorthite content of the plagioclase. This fact demands a critical discussion of possible variations in mode and mineral composition of isochemical rocks as a function of prograde metamorphism. Rapid checks on this topic are facilitated by P. NIGGLI'S (1936a) efficient method of equivalent norms. The equivalent weights of the main constituents of amphibolites closely agree: albite 52.4, anorthite 55.6, zoisite 56.8, tremolite 54.1, tschermakite 54.5 and diopside 54.1. Therefore, the calculated cation percentages of plagioclase, hornblende, clinopyroxene and clinozoisite do not differ much from the actual mode. Only minor components, such as sphene 65.3, rutile 79.9 or magnetite 77.2, show considerably higher equivalent weights (see also the compilation of BURRI, 1958). Table 13 shows the change in quantitative and qualitative mineral composition of three amphibolites with rising anorthite content of their plagioclase. Cation percentages were calculated from rock and hornblende analyses published in this paper. The amount of feldspar increases with rising anorthite content, and the results in Table 13 can be interpreted as representing approximate quantitative mineral reactions in a rock undergoing prograde metamorphism.

Increase of the amount of feldspar with rising anorthite content sheds also some light on the important Na balance of the host rock. In amphibolites plagioclase is the main, hornblende the minor consumer of sodium, so that in first approximation the sodium balance of hornblende can be disregarded.

Assuming that the total amount of Na present in the plagioclase of the rock remains constant and that Ca, Al and Si are freely available in the rock system, a calculation in cation percentages gives the data for An-content of plagioclase and for the total cation percentages of feldspar specified in Table 14 and



Fig. 19. Relations between total cation % plagioclase and anorthite content, assuming that Na remains constant and is equivalent to 1, 2, 3, 4, 5 or 6 cation %.

Fig. 19. This simplified model shows that under those conditions the volume of plagioclase doubles from albite to An 50. This is in good agreement with the observed facts. This theoretical approach which does not consider the role of the hornblende, makes also clear why bytownites and anorthites can only be formed in rocks very low in Na, if other mineral components consume appreciable amounts of Na¹), or if sodium escapes.

4.3.4. Rock composition

Table 11 b gives information on the average chemical composition of amphibolites of different metamorphic grade. Data from recent literature on Alpine greenstones (diabases etc.) and hornblende-bearing greenschists (prasinites) are added for comparison.

From albite- to and esine-amphibolites the mean values for SiO_2 increase distinctly, those for Al_2O_3 very slightly, while Fe_2O_3 and H_2O decrease. The

¹) In this respect the joint occurrence of alkali-feldspar and anorthite in silicatemarbles of the Lepontine Alps may be mentioned.

Fig. 20. Na_2O -CaO diagram for greenstones (diabase etc.), hornblende-bearing greenschists (prasinites), amphibolites with different An content of plagioclase, and for hornblendes occurring in these rocks. Tielines connect hornblende and host rock of the same sample. The diagram shows the wide range of rock composition, contrasting with the limited chemical variation of hornblende.



other oxides show either no consistent trend or no significant variation. In particular, the average content in Na₂O remains almost constant in these amphibolite groups, a fact which is important for the discussion of the feldspar problem. The data for the small group of labradorite-bytownite-amphibolites have to be judged with reserve, as mentioned above; they show higher Al_2O_3 , MgO, CaO, lower Fe_2O_3 , Na₂O and H_2O than and esine-amphibolites. The basic rocks in low-grade greenschist facies differ considerably in composition from the amphibolites. Their mean values for SiO₂, FeO, MgO and CaO are lower, those for Al_2O_3 , Fe_2O_3 , Na₂O and H_2O distinctly higher.

Within the different groups of compared basic metamorphic rocks (see Table 11b), the individual members show wide compositional scatter. But, astonishingly, the mean values for albite-amphibolites, oligoclase-amphibolites and andesine-amphibolites agree fairly well. These amphibolites can be regarded as a rock-series of similar composition. Chemically they fit a gabbroic magma-type after P. NIGGLI (1936), and high-alumina basalt after Kuno (1968). The average values for ophiolitic rocks in greenschist facies, however, assemble in the field of alkali olivine basalt.

The differences between greenstones-greenschists (rocks with alkali-amphibole are excluded!) and amphibolites are well born out also by the Na₂O-CaO diagram of Fig. 20. Alpine ophiolitic rocks termed "diabase" in the literature show the highest Na₂O and lowest CaO contents. Sodium metasomatism may have played an important role in greenschist facies environment, not so in amphibolite facies series.

In the same diagram the projections for hornblendes are also entered and tie-lines connect corresponding rock and mineral analyses. Hornblendes contain generally more CaO and less Na_2O than their host rocks. The discussion will be continued in the next section (p. 125); but here we must emphasize the main impression given by the Na_2O -CaO diagram: The wide range of individual rock composition contrasts with the limited chemical variation of hornblende, concentrated near Na_2O 1.1, CaO 11.9 weight %. This compact field comprises the calcic amphiboles from low- and high-grade amphibolites. Only in some albite-epidote-amphibolite and in greenschist facies rocks do the amphiboles scatter in direction of higher Na_2O and lower CaO contents.

4.3.5. Composition of calcic amphibole

In metamorphic basic rocks in greenschist facies and in albite-epidoteamphibolites from the Alps a fairly broad range of amphibole composition is observed, leading from tremolite and actinolite to common hornblende and extending to pargasitic, edenitic and barroisitic types (WETZEL et al. 1973). In these low-grade rocks with strongly zoned amphiboles a miscibility gap between actinolite and magnesian hornblende is not evident, though it may be confined to a small part of the series; rather a concentration in the field of actinolitic hornblendes is found. In amphibolite facies rocks, with plagioclase > An 5, tremolites and actinolites still occur in rocks of appropriate composition, but there is a broad gap in the tremolite-tschermakite series between Al^{tot} 1.0 and 1.75 (Fig. 21). Not one analysis of a hornblende fits in this space. The bulk of common hornblendes is concentrated between Al^{tot} 1.75 and 2.4.



Fig. 21. Al^{VI}-Al^{IV} Si diagram for calcic hornblendes from the Central Alps, showing the fields of tschermakitic hornblendes from schists and amphibolites of the Gotthard-Tremola-Frodalera area, of Mg- and Fe-hornblendes from amphibolites of the Lepontine Alps, of actinolitic hornblendes and actinolites-tremolites. Circles and dots refer to amphibolite facies rocks, crosses to greenschist facies rocks.

In this compact field no distinction relative to metamorphic grade can be made, except for the fact, reported from rocks all over the world, that amphiboles from labradorite-pyroxene-amphibolites are arranged in the Al^{VI}-poor corner.

As mentioned above (p. 103), truly tschermakitic end members with $Al^{tot} > 2.5$ and $Al^{VI} > 0.9$ are confined to the meta-sedimentary belt south of the Gotthard massif (e. g. biotite-chlorite-garnet-oligoclase-hornblende-garben-schiefer of the Tremola series). The high Al^{VI} content is here hardly related to high pressure. Such tschermakitic hornblendes do not occur in the deeply eroded part of the Lepontine Alps and offer a special problem that needs further research.

The statistical data presented in Table 11a have been calculated from the new hornblende analyses contained in Table 2 of this paper and in the study of WETZEL et al. (1973); moreover the analyses of Alpine hornblendes included in the compilation of LEAKE (1968), and data published by FREY (1969) have been incorporated.

The studied section of the Central Alps is by now probably one of the regions best covered by hornblende analyses, though the number of analyses is still insufficient. The dominant first impression is the great homogeneity of the hornblendes from amphibolites (Gotthard-Tremola excluded). The mean values for Fe_2O_3 , Na_2O and H_2O appear to decrease with rising anorthite content of the coexisting plagioclase, while MgO increases, but the standard deviations are considerable.

Other oxides vary little or inconsistently, unfortunately also TiO_2 , which is only significantly lower in greenschist hornblendes. The hornblendes of greenschists differ irregularly from those of albite-amphibolites, but we are here in the poorly documented field of the strongly zoned actinolitic hornblendes, where the meaning of a chemical analysis of a mineral concentrate becomes uncertain.

Much more significant than the minor variability of hornblendes from amphibolites of different metamorphic grade is the clearly distinct chemical composition of the tschermakitic hornblendes of the Gotthard-Tremola-Frodalera group. Their mean value for Al_2O_3 is much higher, FeO higher and MgO much lower than the averages of all other hornblende groups. The rocks with tschermakitic hornblende occur in the zone of oligoclase-amphibolites and inside the kyanite and staurolite isograds. Most of the rocks from which these Al-rich amphiboles originated are aluminous schists or gneisses and a few are oligoclase-amphibolites proper. But as is now well displayed by the new outcrops in the Gotthard road tunnel, the hornblende-rich rocks are interlayered with a series of garnet-schists and gneisses. The significant differences between the hornblendes of the Tremola group and those of common oligoclase-amphibolites from other parts of the Lepontine Alps should make us cautious: rock composition and the environment within a layered series can influence the mineral composition more strongly than does changing metamorphic grade.

The Na₂O-CaO diagram (Fig. 20) supports graphically some conclusions, drawn from statistical data: the hornblendes of oligoclase-amphibolites (small dots), and esine-amphibolites (medium dots) and labradorite-bytownite-amphibolites (large dots) are crowded in a narrow field. Half of the hornblendes from albite-amphibolites occupy the same position; the other half scatter, as do the hornblendes of prasinites, as they approach higher Na₂O, lower CaO values (pargasitic and edenitic hornblendes). A significant change in hornblende composition is indicated only in these lower-grade rocks, not in the amphibolite facies series which forms the bulk of the material studied in this paper.

4.3.6. Distribution of Or Ab An between coexisting plagioclase and hornblende

In the examined amphibolite series of the Central Alps the proportion $\frac{An}{Or+Ab+An}$ of plagioclase increases with progressive regional metamorphism. This fact is well established (WENK and KELLER, 1969) and, in fact, is still the best indicator of metamorphic grade. The cations substituting in the plagioclase series are distributed between the following mineral components of the rocks:

Si	Al	Ca	Na
quartz			
plagioclase	plagioclase	plagioclase	plagioclase
hornblende	hornblende	hornblende	hornblende
pyroxene		pyroxene	
epidot-clinozoisite	epidote	epidote	—
chlorite	$\mathbf{chlorite}$		
biotite	biotite		(biotite)
muscovite	muscovite	—	(muscovite)
sphene	→	\mathbf{sphene}	
	,	apatite	
zircon		calcite	
<u> </u>		dolomite	

This list shows that Si, Al and Ca can be apportioned to several major and minor mineral components, depending upon the paragenesis, but that Na occurs essentially only in the two main constituents plagioclase and hornblende. Thus the sodium balance is critical, but the role of the other cations should be considered as well. The balance Or Ab An in the rock system and the partitioning of these feldspar "molecules" among the mineral components are discussed with the aid of the feldspar triangle. This procedure is justified also by the fact that 1. in the analysed amphibolites the sum of normative feldspar components amounts to 55–68 cation percent, 2. in the hornblendes 35 to 45% and 3. in plagioclase 100%. Normal plagioclase-amphibolites are made up of 2/3 hornblende and 1/3 feldspar; therefore, commensurable amounts of feldspar components are present in these two main constituents.

Following the methods introduced by P. NIGGLI (1936) the weight percentages of the rock and hornblende analyses were transformed into cation percentages and then Or Ab and An were calculated. In our rocks which carry neither alkali hornblende nor jadeite, Al is always in excess after the formation of Or and Ab. However, there never remains enough Al to form anorthite from the total of Ca present. In all our amphibolites and hornblendes (Na + K + 2Ca) > Al, and after the formation of normative feldspar Wo can



Fig. 22. Or Ab An diagrams for four different amphibolites:

a) Biotite-albite-hornblende-schist MF 147 (M. FREY, 1969), Val Casatscha.

b) Biotite-epidote-albite/oligoclase-amphibolite Cal. 25, Alpe Alögna, Calanca.

- c) Epidote-biotite-oligoclase-amphibolite Hu 1170a₃, Val di Campo.
- d) Andesine/labradorite-amphibolite 20.7.69.01, Monti Laura.

Open circles represent the rock composition.

126

be calculated from the surplus Ca. The bottle-neck in these calculations is Al, not Si^2).

Fig. 22 represents the projections of four different amphibolites with two or more of the components hornblende, plagioclase, biotite and epidote, ranging from albite- to and esine-amphibolite. It shows at first sight that - judged from the rock composition - none of these amphibolites should contain a plagioclase more acid than and esine. Furthermore, it demonstrates that with rising An-content the volumes of feldspar also increase (shorter distance between the projections of plagioclase and rock).

For technical reasons it proved impossible to assemble all the data for the studied rocks in one single diagram Or Ab An. Therefore, only the data for amphibolite and hornblende could be entered in Fig. 23. The rocks (open



Fig. 23. Or Ab An diagram showing hornblende composition (dots) and rock composition (open circles).

circles) occupy the sector An 40 to 70, less than half of the range of the plagioclases found in these amphibolites. Except for biotite-bearing varieties the Or content of the rocks is low. The amphiboles (dots) group in a still narrower field, concentrated between An 60 and 80. If some epidote- and garnet-bearing varieties are disregarded, the amphiboles are always richer in An, poorer in Or and Ab than their host rocks.

In Fig. 24a the normative An content of the hornblende is related to the modal An of the coexisting plagioclase. In hornblende An varies much less than in plagioclase and, except for two cases, the normative An of the hornblende is higher than the modal anorthite content, but a positive correlation is indicated. Thus, in spite of the numerous ionic substitutions possible in amphibole structures, the variability Or Ab An is limited in our calcic hornblendes. Fig. 24b represents the relation between modal and normative anor-

²) In garnet-bearing hornblende rocks too high normative An values result from this calculation scheme.



Fig. 24a. Relation between modal An content of plagioclase and normative An content of the coexisting hornblende.



Fig. 24b. Relation between modal An content of plagioclase and normative An content of the host rock.

thite content in the rock-series. This diagram is instructive in showing that the modal An content is not dictated by rock composition. Albite and oligoclase are not confined to sodium-rich rocks. In Fig. 24 b the rocks lowest and those highest in normative An both contain albite! Once the peristerite gap is passed, a vague positive correlation between An mode and norm may be read from the diagram. Bytownite, though, is restricted to calcic rocks.

4.3.7. Conclusions

In the studied plagioclase-amphibolite series the pronounced increase in modal anorthite content, reflecting progressive metamorphism, takes place without appreciable change in rock composition and without compensating substitution in the coexisting hornblende. It is performed mainly by modal variation. The quantities of Na, Ca, Si and Al remain fairly constant in the system (though not in greenschists and greenstones). Plagioclase has to consume the major part of Na, hornblende the minor part. With rising anorthite content of plagioclase the balance can only be preserved by the formation of more feldspar, mainly at the expense of epidote-clinozoisite, chlorite and biotite, and in later stages also at the expense of hornblende. Modal analyses and calculations from rock and mineral analyses both show that andesineamphibolites contain twice as much plagioclase as albite-amphibolites.

This implies that for each rock-type of given normative Or Ab An ratio and of known paragenesis the maximum mean anorthite content of plagioclase and the maximum normative feldspar content, reached at high-grade metamorphism, can be deduced. In the high-grade metamorphic zone of the Lepontine Alps the highest modal An values observed in plagioclase amphibolites reach An 50 to 60, as can be calculated also from chemical analyses. The An content rarely exceeds An 60; bytownites occur only in hornblendic rock of more calcic composition.

In the same metamorphic zone bytownites/anorthites are common in silicate-marbles, while andesines occur in aluminous schists and oligoclase in the gneiss complexes. In alternating series of these rock-types irregularities in An-content, due to metasomatic processes may be observed, but they are usually confined to the cm-range.

The poor correlation between hornblende composition and metamorphic grade within the different groups of plagioclase-amphibolites confirms observations made in other parts of the world. Therefore, it is really surprising that plagioclase shows such strong compositional variation just in this range of metamorphism.

Acknowledgments

This study was made possible by grants 4282 and 5281 of the Swiss National Foundation for Scientific Research. The authors benefited from discussions with Prof. G. Ernst, Los Angeles, during his stay in Basle. We are especially obliged to our collaborators who contributed in many ways:

Mrs. E. Schedler, Mrs. E. Filisetti, Mrs. R. Mächler and Mr. M. Kristmann for analytical help; T. Fischer and H. Hänni for thin sections and microprobe preparations, Drs. A. Steck and R. Wetzel as well as H. Hänni for the determination of refractive indices; Miss Barbara Hiss for modal analyses; Mrs. A. Lüdin and Mr. H. H. Klein for draughting work; Mrs. S. Lüthi for her skill and patience in typing tables and manuscript. One of the authors is obliged to Julia Wenk (Berkeley) for the careful correction of the text of part 4.3.

Appendix I

Table 1. Analyzing conditions

Apparatus	Jeol JXA-3A	Analyzing crystals	Line
Excitation	Si, Al 10 kV, 0.14 μA Fe, Ca, K 15 kV, 0.09 μA Mg, Na 10 kV, 0.09 μA	RAP, KAP Quartz KAP	ΚαΙ ΚαΙ ΚαΙ
Beam Integration time Counting technique Calibration	1 to 5 microns diameter 30 seconds. point analysis; at least 3 sp standardization with different composition e. g. hornblend	ots per crystal. nt silicate minerals of es, pyroxenes, garnets	known s.

			C	lal	cio	e A	4n	np	hil	00	les	s a	nċ	l A	٩n	nb]	hil	00	lit	\mathbf{es}	fr	on	n 1	the	э]	Lej	po	\mathbf{nt}	in	ел	A 1 _]	\mathbf{ps}					1	31	L
D	3.16	3.25	3.19		3.21	3.24	3.21	3.35	3.21	3.13	3.13	3.28	3.26	3.15	3.24	3.24	3.18	3.26			3.23	3.24	3.20	3.24	3.16	3.29	3.24	3.31	8	3.14	3.16	3.14	3.19	3.18		3.19	3.18	3.22	
	100.1	100.1	100.1	99.4	100.0	100.1	100.1	100.0	99.2	100.0	100.2	100.1	100.0	100.1	100.1	99.9	100.0	100.0	99.7	99.5	100.0	100.0	100.0	100.0	100.1	100.1	100.1	100.1	99.9	100.0	100.0	100.1	100.0	99.8	100.2	99.8	100.9	99.9	99.9
H_2O	2.1	1.8	1.6	3.5	2.9	1.6	2.0	2.6	2.9	1.8	2.2	1.6	2.5	2.0	1.7	2.3	1.6	1.8	2.4	1.5	2.0	1.9	1.9	1.6	2.0	1.6	2.4	2.1	2.1	1.9	1.9	1.7	1.6	2.1	2.3	2.0	2.3	2.4	1.8
P_2O_5				0.1					0.0										0.1	0.0									0.0					0.2	0.2	0.2	0.0	0.1	0.1
TiO_2	0.7	1.0	0.7	2.3	1.7	1.1	1.5	0.2	0.9	1.1	0.4	0.8	1.9	0.5	1.5	1.3	1.4	0.9	1.1	0.8	1.3	0.9	0.7	1.1	0.8	1.0	0.9	0.7	0.6	0.5	0.4	0.8	0.4	0.9	1.2	0.9	0.4	1.2	0.7
K_2O	0.3	0.3	0.6	0.1	0.8	0.6	0.4	0.6	0.0	0.3	0.7	1.1	0.9	0.6	0.5	0.9	0.2	0.9	0.3	0.3	0.6	0.4	0.5	0.3	0.1	0.5	1.2	0.5	0.9	0.5	0.4	0.6	0.8	0.3	0.4	0.5	0.3	0.2	0.3
Na_2O	1.4	1.6	1.6	2.6	2.6	1.1	1.4	0.8	1.4	1.0	1.1	1.1	1.2	0.8	1.3	1.0	2.0	0.9	2.1	2.4	0.1	1.6	1.0	1.5	0.8	1.8	0.9	1.1	1.5	0.8	1.0	1.1	1.2	1.0	1.4	1.0	0.6	1.0	0.8
CaO	10.5	11.7	12.1	8.3	11.5	11.7	11.6	11.6	9.9	12.2	12.0	12.2	11.4	12.4	11.7	11.4	12.1	12.1	9.2	9.6	11.6	11.8	12.3	11.3	12.4	12.2	11.8	11.0	11.0	11.2	12.0	11.9	11.7	12.4	12.8	12.4	13.1	11.9	11.8
MgO	11.9	10.4	12.2	13.2	10.5	11.3	11.9	2.9	7.3	11.9	13.2	9.6	6.8	13.2	11.3	10.3	13.8	10.5	14.0	12.8	10.4	11.5	12.5	11.0	13.8	12.8	10.0	11.8	10.0	14.1	13.7	12.7	11.3	11.5	11.7	11.3	11.4	13.8	13.2
MnO	0.3	0.4	0.5	0.3	0.3	0.3	0.4	0.2	0.4	0.3	0.3	0.5	0.4	0.3	0.3	0.5	0.3	0.5	0.1	0.2	0.3	0.4	0.4	0.2	0.2	0.3	0.3	0.1	0.3	0.1	0.2	0.2	0.7	0.3	0.2	0.3	0.3	0.2	0.2
FeO	10.0	13.3	9.3	10.1	9.7	12.1	12.6	20.4	12.2	8.1	5.7	9.4	14.4	8.6	11.7	12.1	8.8	12.5	6.9	6.5	13.3	12.4	10.1	12.2	9.9	10.1	16.9	10.4	12.2	8.6	9.8	8.1	10.5	10.5	10.1	11.3	10.3	12.5	11.3
${\rm Fe_2O_3}$	5.3	4.7	5.3	2.6	5.2	5.4	2.6	5.5	6.2	4.9	5.6	7.7	6.5	3.4	5.5	5.4	4.6	6.9	1.1	2.8	4.2	4.8	4.8	4.0	3.7	4.8	0.1	3.8	2.6	2.2	2.4	4.6	4.9	3.6	4.0	3.2	6.1	1.3	0.9
Al_2O_3	12.7	12.2	13.4	12.4	12.7	13.1	11.1	13.2	16.8	11.4	11.3	13.7	13.0	10.3	10.9	11.8	11.1	12.5	12.9	14.1	12.0	11.5	12.2	14.6	11.1	11.8	13.2	14.1	9.2	12.1	11.7	12.5	11.9	11.0	12.6	12.1	10.4	10.7	14.1
SiO_2	44.9	42.7	42.8	43.9	42.1	41.8	44.6	42.0	41.2	47.0	47.7	42.4	41.0	48.0	43.7	42.9	44.1	40.5	49.5	48.5	44.2	42.8	43.6	42.2	45.3	43.2	42.4	44.5	49.5	48.0	46.5	45.9	45.0	46.0	43.3	44.6	45.7	44.6	44.7
	AS 912	AS 941	Blen 48 b	Brg 75 conc.	Cal 25	Cal 29	Cal 30	FK 644	GdT S 580	Hu 1164d (Hu)	Hu 1170 (Hu)	Hu 1170a ₃ (Hu)	Hu 1202a (Hy)	Luk 24	Mas 9z	Mas 19	Mera 33a	Mera 34	Mis 52b	Mis 52c conc.	Mto 798a	Mto 798b	Mto 823	Pior 9	PK 834	Riv 17	ShLz A 22	Spl 125	Splug IId cone.	$\dot{Vz} 631$	Vz 673	Wurz 138	Wurz 150	Wurz 175	Wurz 176	Wurz 177b	Wurz 179	20.7.69.01	22.7.69.02

Table 2a. XF Analyses: calcic amphiboles

D	2.91	3.04	3.09	3.06	3.07	3.13	3.17	2.96	10.6	9.04 9.00	2.93 2.05	9.00 2.11	3.06		D	9 10	3.05 2.05	000				100.1	100.1	6.66	100.0	99.7	99.9	100.0	100.0	100.0	99.77	100.06	99.3	100.4	99.67
	100.1 99.8 100.6	100.0	(100.0)	99.8	100.1	100.0	100.1	100.0	99.99	100.0	0.001	100.0	100.1			6 001	2.001 2.001	00 00	99.9		H_2O	3.2	4.3	5.0	4.9	3.8	3.6	3.4	4.1	3.1	4.0	2.6	3.7	5.8	3.7
H_2O	$\begin{array}{c} 13.2 \\ \mathrm{n.b.} \\ 7.4 \end{array}$	1.4 1.0	1.9 0.8	2.4	2.6	2.8 -	1.4	4.2	9.8 -	0.1 V	9.1 0	1 1	1.3		H_2O	0 7		0.7	2.4		P_2O_5														
P_2O_5	$0.2 \\ 0.1 \\ 0.1$	1.0													P_2O_5		0.0	0.0	0.0		TiO_2	2.1	3.1	1.4	1.3	1.9	1.6	2.4	2.4	2.3	0.9	1.1	2.5	1.3	1.5 9 0
TiO_2	0.0	0.1	0.2	0.1	0.1	0.1	0.3	0.0	0.0	1.0	1.0		0.2		TiO_2	и С	0.3	0.0 X	0.2		${\rm K_2O}$	8.2	7.6	8.4	8.4	8.4	9.0	8.0	11.7	10.8	9.6	9.5	9.1	9.1	9.5 7.8
${\rm K_2O}$	0.0	0.2	0.2	0.1	0.1	0.1	0.4	0.3	1.0	1.0	0.0	7.0 0	0.3	les s. l.	${ m K_20}$	80	0.0	10	0.1	nicas	Na_2O	0.4	0.4	0.5	0.3	0.4	0.4	0.4	0.4	0.5	0.3	0.3	0.4	0.3	0.4
Na_2O	0.3 0.2	2.1 6 1)	(1.2)	0.4	0.5	0.5	L.3	0.9	1.4 -	1.0	0.0 0	400	0.3	amphibo	Na_2O	0 6	9.6	200	1.7	ahedral 1	CaO	0.2	0.4	0.0	0.9	0.6	0.2	1.4	0.3	0.4	0.1	0.1	0.1	0.4	0.1
CaO	24.3 15.4 14 1	14.1	12.6	11.7	12.0	12.1	12,0	16.1	20.9	2.11	0 11	0.11	12.3	various	CaO	61	7.T 8 []	8.0	10.1	es: trioct	000	4.0	4.3	3.6	6.9	3.3	6.0	2.2	2.3	3.5	1.3	9.6	6.4	7.1	9.4
MgO	19.2 24.7 93.5	23.5 23.5	(1.62)	21.0	21.0	19.7	14.9	23.4	22.0	4.17 90.4	1.07 03 0	1 0 1	19.2	nalyses:	MgO	10 5	16.3	6.01 6.1	13.4	$^{T}Analys$	M (ľ	Ĺ	7	7	1	7	T	Ĩ	Ξ	1 2	3	H		5T 2
MnO	0.0	0.0	0.0	0.2	0.2	0.2	0.2	0.0	0.0	7.0	1.0	1.0	0.2	c. XFA	MnO	11	1.1	0.1	0.3	ole 3. XI	Mn(0.1	0.2	0.2	0.1	0.3	0.2	0.6	0.4	0.2	0.0	0.0	0.1	0.1	0.0
FeO	$\begin{array}{c} 0.5 \\ \mathrm{n.b.} \\ 0.4 \end{array}$	0.0	0.0 4.4	4.8	4.8	4.6	6.9	0.4	ר ט.א די	0.0 9 1	1.0	1.5	5.6	Table 2	FeO	8 5	5.7	. x c	12.0	Tat	FeO	12.9	14.8	13.6	8.0	13.6	12.7	15.2	14.4	16.0	8.1	6.1	10.3	10.0	8.9 101
${\rm Fe_2O_3}$	0.0 0.9	0.4	1.8 1.8	2.1	1.9	0.9	2.1	0.0	0.0		±.1	1.7	1.3		e ₂ O ₃	0 0	1.3	1.7	2.9		${\rm Fe_2O_3}$	3.2	3.5	2.9	0.1	3.6	2.0	4.1	4.9	3.5	1.5	1.9	1.6	1.9	9.5 9.5
Al_2O_3	0.5 7.5 7	9.7 9.1	5.8	4.2	4.5	3.4	7.9	0.9	0.8 0	0.0 0	0.0 1 6	6.9	5.1	,	M_2O_3 F	34 1	12.0	14.6	6.1		Al_2O_3	18.0	16.0	17.8	17.8	17.6	16.6	16.4	15.9	15.3	14.3	19.2	17.6	16.8	17.0
SiO_2	41.9 53.8 48.5	48.5 47.7	41.1 54.0	52.8	52.4	55.6 20 -	52.7	53.8	44.0 z 4 o	04.0 51 6	0.10	52.2	54.3		iO ₂ A	3 4	3.1	22	0.7		SiO_2	37.8	35.5	36.5	41.3	36.2	37.6	35.9	33.2	34.4	39.6	39.6	37.5	37.6	38.9 38 1
	Brg 2 Cadonigo grau Cadonigo grau	CT 19 «L»	Hu 1015 (Hu)	Hu 1026 I (Hu)	Hu 1026e I (Hu)	Hu 1026 II (Hu)	Hu 1026e 11	Mera 40		Varzo 0 Varzo 7	V_{π} 104	V_{Z} 394	Vz 428		Ω.	Bni 958		KAW 657 5	KAW 658 50			Cal 25	Cal 30	GdT S 580	Hu 1015 (Hu)	Hu 1170a 3 (Hu)	Luk 24	Mas 19	Mera 34	Mto 798b	Varzo 6	Varzo 7	Vz 194	Vz 394	Vz 428 Wiirz 138

Table 2 b. XFAnalyses: actinolitic/tremolitic hornblendes

132

E. Wenk, H. Schwander and W. Stern

	(Ca	lci	ic	A	mp	b	ib	ole	es i	an	d	Aı	ՠֈ	b	ib	oli	tes	s f	ro	m	tł	ıe	Le	əp	on	tir	ne	A	lp	3				
	100.0	100.0	99.8	100.5	99.8	99.7	100.9	100.2	100.43	100.2	99.7	99.9	99.8	99.75	100.5	100.6	98.8	99.3	100.1	99.7	100.4	101.3	100.0	99.9	100.0	99.5	100.1	100.5	100.8	99.2	100.2	99.1	100.0	99.8	99.8
H_2O	2.9	1.9	1.2	2.0	3.9	2.9	1.9	1.3	2.1	1.3	1.8	3.5	2.9	2.7	1.7	1.3	2.2	1.6	1.6	1.4	2.2	1.6	1.4	1.9	1.5	1.7	1.3	1.6	1.6	1.8	1.9	2.0	1.8	2.1	7.0
P_2O_5			0.1	0.2	0.2	0.3				0.2			0.0	0.1	0.2					0.1			0.1		0.2	0.4	n.b.	0.1			0.3		0.0		
TiO_2	1.4	1.9	1.3	1.3	1.5	2.4	1.3	1.1	2.3	1.3	1.0	1.5	1.2	1.1	1.6	0.3	1.3	0.9	2.0	0.6	0.5	1.2	1.2	1.7	1.6	1.7	1.7	1.4	1.3	1.4	1.3	0.9	0.3	0.6	1.1
${\rm K_2O}$	0.3	0.6	0.8	0.1	0.8	1.2	0.5	0.9	0.7	0.3	2.3	1.1	0.5	0.4	0.6	0.3	0.5	2.0	1.8	2.1	3.0	0.4	0.3	1.5	0.3	0.3	0.4	0.2	0.2	0.2	0.5	1.6	1.1	0.6	0.2
Na_2O	3.1	2.7	3.0	4.2	3.7	1.3	2.9	2.8	3.4	3.2	2.2	2.7	2.0	1.9	2.2	2.6	2.8	3.6	0.8	3.4	0.3	1.8	3.2	2.6	3.4	2.4	1.7	1.6	4.1	2.7	3.2	2.9	1.8	1.8	3.5
CaO	10.2	11.4	10.6	7.2	12.0	7.6	11.3	9.8	8.9	11.8	8.1	8.5	8.6	8.8	12.6	13.8	12.9	6.5	9.7	6.0	5.7	11.8	10.3	11.2	10.8	12.0	12.3	13.9	10.7	11.5	10.1	9.0	9.3	9.5	9.6
MgO	8.4	6.7	5.5	8.0	6.8	5.8	8.5	3.2	3.0	7.6	5.5	6.1	11.7	11.5	8.4	14.3	11.8	4.5	6.6	3.4	18.1	9.1	7.1	6.4	7.4	8.0	7.7	6.9	9.2	8.8	7.6	11.4	11.5	8.9	8.2
MnO	0.2	0.2	0.2	0.2	0.4	0.3	0.2	0.2	0.21	0.0	0.0	0.1	0.0	0.05	0.2	0.1	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.1	0.2		0.1
FeO	6.6	9.7	5.4	7.6	6.5	10.3	8.6	6.2	8.32	6.0	5.1	6.5	7.9	8.3	7.1	3.9	7.9	4.6	8.1	5.4	7.0	6.1	7.9	6.7	6.6	8.0	7.6	6.1	6.2	7.2	7.8	6.4	7.4	6.5	5.2
${\rm Fe_2O_3}$	3.7	3.1	3.6	2.0	3.2	3.5	0.2	3.8	2.4	1.4	2.7	5.3	1.9	2.2	2.1	0.8	4.0	3.1	2.6	0.2	1.6	2.1	1.5	4.8	1.9	1.5	2.3	4.2	2.1	1.8	2.3	1.7	1.4	2.0	5.4
Al_2O_3	15.3	13.6	18.8	16.9	17.5	14.5	14.9	16.9	16.1	18.8	17.8	13.3	15.7	15.5	17.4	14.8	13.8	16.2	14.3	15.3	14.6	16.9	16.3	16.5	16.6	15.2	17.0	17.0	16.0	17.0	15.6	13.9	13.2	17.4	16.4
SiO_2	47.9	48.2	49.3	50.8	43.3	49.6	50.6	54.0	53.0	48.3	53.2	51.3	47.4	47.2	46.4	48.4	41.4	56.1	52.4	61.7	47.3	50.1	50.5	46.4	49.5	48.1	48.0	47.4	49.2	46.7	49.4	49.2	52.0	50.4	43.1
	As 912	As 941	Brg 73	Brg 75	Cal 25	GdT S 580	Hu 1164d	Hu 1170a 3 (Hu)	Hu 1202a	Mera 33a	Mera 34	Mis 52	Mis 52b	Mis 52c	Riv 17	Sci 171	Sci 186	Sci 244	Sh Lz A 22a	Spluga 11 d	Vz 428	Vz 631	Vz 703	Wurz 166	Wurz 175	Wurz 176	Wurz 177b	Wurz 179	19.7.69.01	20.7.69.01	20.7.69.02	20.7.69.03	22.7.69.01	22.7.69.02	29.9.69.01

uM 0	Mg·10(⁺ fot9 ⁺ F ⁺ gM	99.2 99.2 99.0 99.0	$\begin{array}{c} 90.5\\ 90.2\\ 89.1\\ 86.1\\ 85.0\end{array}$	84.8 84.5 84.2 83.1 83.1	75.6 75.4 74.8 71.3 70.5	70.0 68.1 66.9 66.3 66.2	$\begin{array}{c} 65.7 \\ 65.0 \\ 64.8 \\ 64.6 \\ 64.5 \end{array}$
	$M_{g}^{\prime\prime}$	$\begin{array}{c} 91.0\\ 99.0\\ 99.0\\ 91.9\\ 89.4\end{array}$	98.7 98.3 88.8 82.6 75.8	60.2 83.3 80.7 75.4 67.1	58.1 69.3 62.5 52.6 84.2	57.6 57.8 57.0 30.9 57.0	53.7 58.1 56.7 57.6 53.7
	$ m R^{2+}_{2+}$	$\frac{1.0}{9.2}$	$1.3 \\ 0.9 \\ 7.8 \\ 11.3 \\ 9.9 \\ 9.9 \\ 0.9$	$18.5 \\11.1 \\12.1 \\12.8 \\13.4 \\13.4$	$16.3 \\ 19.6 \\ 16.7 \\ 15.5 \\ 11.3 \\ $	$19.9 \\ 14.8 \\ 23.4 \\ 11.1 \\ 21.6 \\ $	$\begin{array}{c} 26.2 \\ 21.5 \\ 25.1 \\ 23.7 \\ 19.7 \end{array}$
	$ m R_{3+}^{\%}$	9.0 8.1 1.4	$\begin{array}{c} 0.9\\ 3.6\\ 6.1\\ 14.3\end{array}$	21.3 5.5 7.2 11.9 19.5	25.5 11.1 20.8 32.0 4.5	$\begin{array}{c} 22.5\\ 27.4\\ 19.6\\ 58.0\\ 21.5\end{array}$	$\begin{array}{c} 20.1 \\ 20.3 \\ 18.2 \\ 18.7 \\ 26.6 \end{array}$
	Alk.	$\begin{array}{c} 2.66\\ 3.17\\ 2.63\\ 2.72\\ 1.83\end{array}$	$\begin{array}{c} 3.46 \\ 2.09 \\ 2.32 \\ 1.93 \\ 2.04 \end{array}$	$\begin{array}{c} 2.19\\ 1.87\\ 1.72\\ 1.99\\ 2.53\end{array}$	$\begin{array}{c} 2.00\\ 1.50\\ 2.25\\ 2.18\\ 1.94\end{array}$	$\begin{array}{c} 2.02\\ 2.26\\ 2.21\\ 2.73\\ 2.24\\ 2.24\end{array}$	$\begin{array}{c} 2.11 \\ 2.50 \\ 2.17 \\ 2.26 \end{array}$
	Okt.	5.23 4.87 4.78 5.32 5.30	3.77 4.66 4.68 4.88 5.45	5.25 5.24 5.52 5.33 5.15	5.01 5.83 5.03 5.15 5.17	5.20 4.84 5.17 4.02 4.96	5.31 5.18 4.95 5.17 5.09
(HO)	но	$\begin{array}{c} 1.75\\ 2.33\\ 3.80\\ 1.30\\ 1.93\end{array}$	$\begin{array}{c} 11.45 \\ 6.46 \\ 4.11 \\ 2.57 \\ 0.75 \end{array}$	$\begin{array}{c} 1.43 \\ 2.23 \\ 1.49 \\ 1.22 \\ 2.18 \end{array}$	$\begin{array}{c} 2.23\\ 1.42\\ 1.32\\ 1.42\\ 2.41\end{array}$	$\begin{array}{c} 1.81\\ 2.09\\ 1.83\\ 1.73\\ 1.92\end{array}$	$\begin{array}{c} 1.74 \\ 1.56 \\ 3.33 \\ 1.93 \\ 1.64 \end{array}$
<i>84 (0</i> ,	а		$\begin{array}{c} 0.02 \\ 0.01 \\ & \end{array}$		0.01	0.01	$0.01 \\ -0.01$
boles, 2	Ti	$0.01 \\ - 0.01 \\ 0.01 \\ 0.01$	$\begin{array}{c} - \\ 0.01 \\ 0.01 \\ 0.01 \\ 0.02 \end{array}$	$\begin{array}{c} 0.07 \\ 0.01 \\ 0.01 \\ 0.02 \\ 0.04 \end{array}$	$\begin{array}{c} 0.13 \\ 0.02 \\ 0.04 \\ 0.10 \\ 0.01 \end{array}$	$\begin{array}{c} 0.06 \\ 0.05 \\ 0.05 \\ 0.05 \\ 0.21 \\ 0.06 \end{array}$	$\begin{array}{c} 0.08\\ 0.17\\ 0.17\\ 0.28\\ 0.10\\ 0.10\end{array}$
umphil	К	$\begin{array}{c} 0.04 \\ 0.02 \\ 0.05 \\ 0.04 \\ 0.02 \end{array}$	$0.02 \\ 0.05 \\ 0.02 \\ 0.04 $	$\begin{array}{c} 0.04 \\ 0.02 \\ 0.02 \\ 0.05 \\ 0.07 \end{array}$	$\begin{array}{c} 0.05 \\ 0.04 \\ 0.07 \\ 0.05 \\ 0.02 \end{array}$	$\begin{array}{c} 0.09\\ 0.13\\ 0.07\\ 0.02\\ 0.11\\ 0.11\end{array}$	$\begin{array}{c} 0.06 \\ 0.04 \\ 0.02 \\ 0.02 \\ 0.11 \end{array}$
lyzed c	Na	$\begin{array}{c} 0.56\\ 0.46\\ 0.24\\ 0.57\\ 0.65\end{array}$	$\begin{array}{c} 0.08\\ 0.10\\ 0.08\\ 0.08\\ 0.13\\ 0.11\\ 0.11\end{array}$	$\begin{array}{c} 0.39\\ 0.11\\ 0.03\\ 0.03\\ 0.08\\ 0.69\end{array}$	$\begin{array}{c} 0.57\\ 0.11\\ 0.11\\ 0.36\\ 0.66\\ 0.13\end{array}$	$\begin{array}{c} 0.22 \\ 0.30 \\ 0.28 \\ 1.41 \\ 0.22 \end{array}$	$\begin{array}{c} 0.22\\ 0.57\\ 0.72\\ 0.72\\ 0.22\\ 0.31\end{array}$
of ana	Ca	$\begin{array}{c} 2.06\\ 2.69\\ 2.34\\ 2.11\\ 1.76\end{array}$	3.39 1.98 2.18 1.78 1.78 1.90	$1.77 \\ 1.75 \\ 1.68 \\ 1.85 \\ 1.77 \\ $	$\begin{array}{c} 1.38 \\ 1.35 \\ 1.82 \\ 1.46 \\ 1.79 \end{array}$	$1.71 \\ 1.83 \\ 1.86 \\ 1.30 \\ 1.91 \\ 1.91 \\$	$\begin{array}{c} 1.83 \\ 1.90 \\ 1.27 \\ 1.93 \\ 1.84 \end{array}$
vulae e	Mg	4.76 4.82 4.73 4.73 4.73 4.73	3.72 4.58 4.16 4.03 4.13	3.16 4.37 4.46 4.02 3.46	$\begin{array}{c} 2.91 \\ 4.04 \\ 3.15 \\ 2.71 \\ 4.35 \end{array}$	$\begin{array}{c} 2.99\\ 2.80\\ 2.95\\ 1.24\\ 2.82\\ 2.82\end{array}$	$\begin{array}{c} 2.85\\ 3.01\\ 2.80\\ 2.98\\ 2.73\end{array}$
ul forn	Mn	0.0	$\begin{array}{c} - \\ 0.01 \\ 0.02 \\ 0.02 \end{array}$	$\begin{array}{c} 0.04 \\ 0.02 \\ 0.02 \\ 0.02 \\ 0.01 \end{array}$	$\begin{array}{c} 0.01 \\ 0.02 \\ 0.02 \\ 0.02 \\ 0.02 \end{array}$	$\begin{array}{c} 0.01\\ 0.04\\ 0.02\\ 0.01\\ 0.04\\ 0.04\end{array}$	$\begin{array}{c} 0.02 \\ 0.04 \\ 0.04 \\ 0.02 \\ 0.02 \end{array}$
ructure	Fe ²⁺	$\begin{array}{c} 0.05 \\ 0.05 \\ - \end{array}$	$\begin{array}{c} 0.05\\ 0.04\\ 0.35\\ 0.53\\ 0.53\\ 0.52\end{array}$	$\begin{array}{c} 0.93\\ 0.56\\ 0.64\\ 0.66\\ 0.68\\ 0.68\end{array}$	$\begin{array}{c} 0.81\\ 1.12\\ 0.82\\ 0.77\\ 0.56\end{array}$	$\begin{array}{c} 1.02 \\ 0.68 \\ 1.18 \\ 0.43 \\ 1.03 \end{array}$	$\begin{array}{c} 1.37\\ 1.08\\ 1.08\\ 1.20\\ 1.20\\ 0.98\end{array}$
5. Sta	Fe ³⁺	$\begin{array}{c} 0.04 \\ \\ 0.04 \\ 0.01 \end{array}$	$\begin{array}{c} - & - & - & - & - & - & 0 \\ 0.03 & 0.04 & 0.09 & 0.19 & 0.19 \end{array}$	$\begin{array}{c} 0.34 \\ 0.22 \\ 0.18 \\ 0.14 \\ 0.14 \end{array}$	$\begin{array}{c} 0.12 \\ 0.18 \\ 0.22 \\ 0.30 \\ 0.20 \end{array}$	$\begin{array}{c} 0.24 \\ 0.60 \\ 0.26 \\ 0.17 \\ 0.37 \end{array}$	$\begin{array}{c} 0.10\\ 0.51\\ 0.28\\ 0.40\\ 0.50\end{array}$
Table	Alvī	$\begin{array}{c} 0.42 \\ - \\ 0.38 \\ 0.05 \end{array}$	0.19 0.56	$\begin{array}{c} 0.70 \\ 0.06 \\ 0.20 \\ 0.47 \\ 0.83 \end{array}$	$\begin{array}{c} 1.03 \\ 0.44 \\ 0.79 \\ 1.25 \\ 0.02 \end{array}$	$\begin{array}{c} 0.87\\ 0.68\\ 0.68\\ 0.70\\ 1.95\\ 0.64\end{array}$	$\begin{array}{c} 0.89\\ 0.37\\ 0.35\\ 0.35\\ 0.46\\ 0.76\end{array}$
	AlIV	$\begin{array}{c} 1.41 \\ 0.16 \\ 0.14 \\ 1.22 \\ 0.21 \\ 0.21 \end{array}$	$\begin{array}{c} 0.08\\ 0.88\\ 0.53\\ 0.36\\ 0.40\\ \end{array}$	$\begin{array}{c} 1.07 \\ 0.63 \\ 0.34 \\ 0.37 \\ 1.83 \end{array}$	$\begin{array}{c} 1.09 \\ 0.59 \\ 0.53 \\ 1.11 \\ 1.11 \\ 0.71 \end{array}$	$1.16\\1.21\\1.21\\1.29\\0.41\\1.11$	1.52 1.54 1.74 1.43 1.37
	Al_{tot}	$\begin{array}{c} 1.82\\ 0.16\\ 0.14\\ 1.60\\ 0.26\\ \end{array}$	$\begin{array}{c} 0.08\\ 0.88\\ 0.53\\ 0.55\\ 0.96\\ 0.96 \end{array}$	1.77 0.69 0.54 0.84 2.67	$\begin{array}{c} 2.12\\ 1.04\\ 1.32\\ 2.36\\ 0.74\end{array}$	$\begin{array}{c} 2.03\\ 1.89\\ 2.00\\ 2.35\\ 1.74\end{array}$	2.41 1.92 2.08 1.90 2.13
	Si	$\begin{array}{c} 6.59 \\ 7.40 \\ 7.31 \\ 6.78 \\ 7.79 \end{array}$	5.45 6.35 7.07 7.64 7.60	6.93 7.37 7.66 7.63 6.17	$\begin{array}{c} 6.91 \\ 7.41 \\ 7.47 \\ 6.89 \\ 7.29 \end{array}$	$\begin{array}{c} 6.84 \\ 6.79 \\ 6.71 \\ 7.59 \\ 6.89 \end{array}$	$\begin{array}{c} 6.48 \\ 6.46 \\ 6.26 \\ 6.57 \\ 6.63 \end{array}$
		CT 19 «S» TV 116e, corr. Mera 40 CT 19 «L» Vz 194	Brg 2 Trem. conc. Cadonigo «grun» Varzo 7 Strahlst. Hu 1026 II Hu 1015	Hu 1026e I Hu Hu 1026 I Varzo 6 Vz 428 Ho Vo	Mis 52b Vz 394 Hu 1026e II Mis 52c Hu 1026e I Hy	Vz 631 Hu 1170 Vz 673 KAW 657 Luk 24	22.7.69.02 Mera 33a Brg 75 conc. PK 834 Wurz 138

Table 5. Structural formulae of analyzed amphiboles, 24 (0, 0H)

134

$\begin{array}{c} 63.9\\ 62.4\\ 61.7\\ 60.7\\ 60.1\\ 60.1\end{array}$	$\begin{array}{c} 60.0\\ 60.0\\ 60.0\\ 59.3\\ 59.3\\ 58.4\end{array}$	58.0 58.0 56.4 55.9 55.9	55.0 54.6 54.5 54.2 54.2 53.9	51.6 51.2 50.9 50.8 50.4	50.0 49.4 41.5 37.1 16.9
58.0 51.7 57.1 54.3 49.4	52.4 51.6 53.1 51.0 49.8	$\begin{array}{c} 49.6 \\ 50.7 \\ 48.2 \\ 49.5 \\ 50.7 \end{array}$	$\begin{array}{c} 45.6\\ 49.3\\ 45.1\\ 48.4\\ 47.2\\ \end{array}$	44.1 45.4 44.7 43.4 43.4	47.0 45.2 32.7 31.8 14.0
$\begin{array}{c} 30.0\\ 20.5\\ 29.4\\ 24.8\\ 24.6\\ 24.6\end{array}$	$\begin{array}{c} 25.9\\ 25.0\\ 25.0\\ 26.9\\ 24.2\\ 24.2\end{array}$	28.6 31.1 26.8 26.4 26.5	$\begin{array}{c} 28.8\\ 30.8\\ 31.6\\ 28.8\\ 29.1\\ \end{array}$	32.4 31.2 33.0 24.6	$\begin{array}{c} 22.4\\ 31.4\\ 31.7\\ 38.8\\ 55.8\end{array}$
$12.0 \\ 27.8 \\ 13.5 \\ 21.0 \\ 26.0 \\ $	21.7 25.1 21.8 22.1 26.0	21.8 18.2 25.0 24.1 22.9	$\begin{array}{c} 25.6 \\ 19.9 \\ 23.3 \\ 22.8 \\ 23.8 \\ 23.8 \end{array}$	$\begin{array}{c} 23.5\\ 23.3\\ 22.3\\ 14.6\\ 32.9\\ 32.9\end{array}$	$\begin{array}{c} 30.6\\ 23.3\\ 35.7\\ 29.4\\ 30.2\end{array}$
$\begin{array}{c} 2.17\\ 2.22\\ 2.25\\ 2.54\\ 2.10\\ 2.10 \end{array}$	2.46 2.48 2.31 2.27 2.08	$\begin{array}{c} 2.32\\ 2.29\\ 2.34\\ 2.68\\ 2.26\\ 2.26\end{array}$	$\begin{array}{c} 2.27 \\ 2.41 \\ 2.30 \\ 2.32 \\ 2.29 \end{array}$	$\begin{array}{c} 1.97 \\ 2.26 \\ 2.38 \\ 2.37 \\ 2.46 \end{array}$	$\begin{array}{c} 2.52 \\ 2.39 \\ 1.95 \\ 2.35 \\ 2.24 \end{array}$
5.16 4.94 5.04 5.19 5.15	$\begin{array}{c} 4.84\\ 5.18\\ 5.14\\ 4.88\\ 5.16\\ 5.16\end{array}$	4.97 5.13 5.14 4.62 4.85	5.31 5.16 4.80 5.15 5.31	5.17 5.00 5.16 5.10 4.99	4.77 5.21 4.88 4.77 4.77
$\begin{array}{c} 2.33\\ 1.73\\ 2.31\\ 1.57\\ 2.03\\ 2.03\end{array}$	$2.23 \\ 1.57 \\ 1.86 \\ 2.04 \\ 2.03 $	$\begin{array}{c} 1.95 \\ 1.96 \\ 1.57 \\ 2.82 \\ 2.22 \end{array}$	$\begin{array}{c} 1.58\\ 1.88\\ 2.03\\ 1.68\\ 1.59\\ 1.59\end{array}$	$\begin{array}{c} 1.96\\ 2.27\\ 1.79\\ 2.38\\ 2.38\\ 1.58\end{array}$	2.58 1.81 2.84 2.50 2.64
0.01	$\begin{array}{c c} 0.02 \\ 0.02 \\ 0.02 \end{array}$	0.02			
$\begin{array}{c} 0.15 \\ 0.13 \\ 0.02 \\ 0.13 \\ 0.13 \\ 0.09 \end{array}$	$\begin{array}{c} 0.15\\ 0.09\\ 0.09\\ 0.11\\ 0.11\\ 0.09\end{array}$	$\begin{array}{c} 0.11\\ 0.19\\ 0.05\\ 0.21\\ 0.05\\ 0.05\end{array}$	$\begin{array}{c} 1.14 \\ 0.11 \\ 0.07 \\ 0.19 \\ 0.14 \end{array}$	$\begin{array}{c} 0.16\\ 0.16\\ 0.13\\ 0.13\\ 0.11\\ 0.10\\ 0.10 \end{array}$	$\begin{array}{c} 0.06\\ 0.11\\ 0.11\\ 0.24\\ 0.28\\ 0.03\end{array}$
$\begin{array}{c} 0.04 \\ 0.05 \\ 0.02 \\ 0.09 \\ 0.09 \end{array}$	$\begin{array}{c} 0.07\\ 0.11\\ 0.09\\ 0.06\\ 0.06\\ 0.06\end{array}$	$\begin{array}{c} 0.09 \\ 0.07 \\ 0.15 \\ 0.15 \\ 0.15 \\ 0.06 \end{array}$	$\begin{array}{c} 0.06\\ 0.08\\ 0.17\\ 0.09\\ 0.09\\ 0.11\end{array}$	$\begin{array}{c} 0.11\\ 0.17\\ 0.16\\ 0.06\\ 0.23\\ 0.23\\ 0.21\end{array}$	$\begin{array}{c} 0.14 \\ 0.17 \\ \\ 0.17 \\ 0.12 \end{array}$
$\begin{array}{c} 0.28\\ 0.28\\ 0.47\\ 0.51\\ 0.31\end{array}$	$\begin{array}{c} 0.39\\ 0.46\\ 0.28\\ 0.28\\ 0.39\\ 0.39\end{array}$	$\begin{array}{c} 0.28\\ 0.40\\ 0.34\\ 0.74\\ 0.17\end{array}$	$\begin{array}{c} 0.43\\ 0.46\\ 0.42\\ 0.37\\ 0.32\\ 0.32\end{array}$	$\begin{array}{c} 0.03\\ 0.29\\ 0.46\\ 0.26\\ 0.32\\ 0.32\end{array}$	$\begin{array}{c} 2.19\\ 0.26\\ 0.40\\ 0.35\\ 0.24\\ 0.35\\ 0.24\\ \end{array}$
$\begin{array}{c} 1.86\\ 1.88\\ 1.56\\ 1.56\\ 1.93\\ 1.70\end{array}$	$\begin{array}{c} 2.00\\ 1.91\\ 1.93\\ 1.93\\ 1.93\\ 1.63\end{array}$	$\begin{array}{c} 1.94 \\ 1.82 \\ 1.85 \\ 1.86 \\ 1.80 \\ 2.03 \end{array}$	$\begin{array}{c} 1.79 \\ 1.88 \\ 1.71 \\ 1.71 \\ 1.86 \\ 1.87 \\ 1.87 \end{array}$	$\begin{array}{c} 1.83\\ 1.81\\ 1.81\\ 1.87\\ 1.88\\ 1.94\\ 1.94\end{array}$	$\begin{array}{c} 0.18\\ 1.95\\ 1.55\\ 1.55\\ 1.83\\ 1.89\\ 1.89\end{array}$
$\begin{array}{c} 3.00\\ 2.56\\ 2.88\\ 2.81\\ 2.54\end{array}$	2.54 2.68 2.49 2.57	2.46 2.60 2.48 2.28 2.48 2.46	2.42 2.55 2.16 2.49 2.51	2.28 2.21 2.31 2.22 2.22 2.12	$\begin{array}{c} 2.24\\ 2.36\\ 1.59\\ 1.52\\ 0.66\end{array}$
$\begin{array}{c} 0.02\\ 0.04\\ 0.04\\ 0.04\\ 0.04\\ 0.01\end{array}$	$\begin{array}{c} 0.02\\ 0.06\\ 0.05\\ 0.04\\ 0.04\end{array}$	$\begin{array}{c} 0.04\\ 0.05\\ 0.09\\ 0.04\\ 0.04\\ 0.04\end{array}$	$\begin{array}{c} 0.03\\ 0.05\\ 0.04\\ 0.04\\ 0.04\end{array}$	$\begin{array}{c} 0.04 \\ 0.06 \\ 0.05 \\ 0.04 \\ 0.06 \end{array}$	$\begin{array}{c} 0.13 \\ 0.06 \\ 0.05 \\ 0.05 \\ 0.03 \end{array}$
$\begin{array}{c} 1.52\\ 0.98\\ 1.45\\ 1.25\\ 1.26\end{array}$	$1.23 \\ 1.14 \\ 1.24 \\ 1.28 \\ 1.28 \\ 1.21 \\ $	$\begin{array}{c} 1.38 \\ 1.55 \\ 1.29 \\ 1.18 \\ 1.25 \end{array}$	$1.51 \\ 1.54 \\ 1.48 \\ 1.45 \\ 1.51 \\ 1.51$	$\begin{array}{c} 1.64 \\ 1.50 \\ 1.66 \\ 2.10 \\ 2.10 \\ 1.17 \end{array}$	$\begin{array}{c} 0.93 \\ 1.57 \\ 1.50 \\ 1.80 \\ 1.80 \\ 2.60 \end{array}$
$\begin{array}{c} 0.14 \\ 0.53 \\ 0.31 \\ 0.53 \\ 0.53 \\ 0.41 \end{array}$	$\begin{array}{c} 0.44 \\ 0.59 \\ 0.53 \\ 0.39 \\ 0.58 \end{array}$	$\begin{array}{c} 0.35\\ 0.29\\ 0.54\\ 0.57\\ 0.66\end{array}$	$\begin{array}{c} 0.44\\ 0.54\\ 0.54\\ 0.28\\ 0.61\\ 0.60\end{array}$	$\begin{array}{c} 0.47 \\ 0.60 \\ 0.53 \\ 0.01 \\ 0.86 \end{array}$	$\begin{array}{c} 1.17\\ 0.78\\ 0.68\\ 0.73\\ 0.73\\ 0.63\end{array}$
$\begin{array}{c} 0.33\\ 0.71\\ 0.34\\ 0.43\\ 0.43\\ 0.84\end{array}$	$\begin{array}{c} 0.46\\ 0.63\\ 0.51\\ 0.57\\ 0.68\\ 0.68\end{array}$	$\begin{array}{c} 0.62 \\ 0.46 \\ 0.69 \\ 0.33 \\ 0.40 \end{array}$	$\begin{array}{c} 0.78 \\ 0.38 \\ 0.36 \\ 0.37 \\ 0.37 \\ 0.52 \end{array}$	$\begin{array}{c} 0.59 \\ 0.40 \\ 0.50 \\ 0.62 \\ 0.68 \\ 0.68 \end{array}$	$\begin{array}{c} 0.23\\ 0.32\\ 0.95\\ 0.43\\ 0.43\\ 0.76\end{array}$
$1.51 \\ 1.22 \\ 0.69 \\ 1.62 \\ 1.56 \\ $	1.70 1.70 1.60 1.31 1.31 1.49	$1.47 \\ 1.46 \\ 1.37 \\ 1.37 \\ 1.38 \\ $	$\begin{array}{c} 1.76 \\ 1.64 \\ 0.82 \\ 0.82 \\ 1.53 \\ 1.78 \end{array}$	1.49 1.65 1.64 1.64 1.71	$\begin{array}{c} 0.35\\ 1.90\\ 1.96\\ 1.86\\ 1.86\\ 1.61\end{array}$
$1.84 \\ 1.94 \\ 1.04 \\ 2.05 \\ 2.40 \\$	$\begin{array}{c} 2.16\\ 2.32\\ 2.11\\ 1.88\\ 2.17\\ 2.17\end{array}$	$\begin{array}{c} 2.09 \\ 1.92 \\ 2.06 \\ 2.18 \\ 2.18 \\ 1.78 \end{array}$	2.54 2.01 1.57 1.90 2.30	$\begin{array}{c} 2.08\\ 2.06\\ 2.14\\ 2.31\\ 2.39\\ 2.39\end{array}$	$\begin{array}{c} 0.57\\ 2.22\\ 2.90\\ 2.29\\ 2.37\\ 2.37\end{array}$
$\begin{array}{c} 6.49 \\ 6.78 \\ 7.31 \\ 6.38 \\ 6.44 \end{array}$	$\begin{array}{c} 6.30 \\ 6.30 \\ 6.40 \\ 6.69 \\ 6.51 \end{array}$	$\begin{array}{c} 6.53 \\ 6.54 \\ 6.63 \\ 6.15 \\ 6.62 \\ 6.62 \end{array}$	$\begin{array}{c} 6.24 \\ 6.36 \\ 7.18 \\ 6.47 \\ 6.22 \end{array}$	$\begin{array}{c} 6.51 \\ 6.35 \\ 6.36 \\ 6.31 \\ 6.31 \\ 6.29 \\ 6.29 \end{array}$	$\begin{array}{c} 7.65 \\ 6.10 \\ 6.04 \\ 6.14 \\ 6.14 \\ 6.39 \end{array}$
20.7.69.01 Hu 1164d KAW 658 Riv 17 Spl 125	Wurz 176 Blen 48b Mto 823 Wurz 175 AS 912	Wurz 177 b Cal 30 Wurz 150 Cal 25 Wurz 179	Pior 9 Mto 798b Splug 11d cone. Mas 9z Cal 29	Mto 798a Mas 19 AS 941 Sh Lz A 22 Hu 1170a ₃	Bni 258 Mera 34 GdT S 580 Hu 1202a FK 644

Table 6. Microprobe analyses of calcic amphiboles, structural formulae (basis 23 O)

		Ad 36e			AS 907			AS 907I	I
SiO,	47.5	49.7	51.0	47.5	51.5	49.8	51.5	45.7	46.8
$Al_2 \tilde{O}_3$	11.9	11.5	11.3	12.0	9.1	8.8	6.3	13.1	13.8
FeO	8.9	8.9	7.4	13.8	12.5	12.5	13.9	15.0	14.2
MgO	14.6	15.2	16.7	12.0	11.1	13.9	10.2	9.8	9.8
CaO	11.0	8.6	9.4	12.9	12.7	12.6	12.5	12.8	12.5
Na ₂ O	1.6	2.5	1.8	0.9	1.5	1.5	1.2	0.8	1.3
K ₂ Ō	0.4	0.4	0.4	0.4	0.4	0.4	0.9	0.7	0.8
$\overline{\text{TiO}}_2$	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0.6
	95.9	96.8	98.0	99.7	99.0	100.1	97.0	98.4	99.8
Si	6.88	7.07	7.10	6.79	7.33	7.05	7.55	6.68	6.71
Al^{IV}	1.12	0.93	0.90	1.21	0.67	0.95	0.45	1.32	1.29
Alvi	0.91	1.00	0.96	0.81	0.85	0.52	0.64	0.93	1.04
$\mathrm{Fe^{+2}}$	1.08	1.06	0.86	1.65	1.49	1.48	1.70	1.83	1.70
Mg	3.15	3.22	3.47	2.56	2.35	2.93	2.23	2.13	2.09
Ca	1.70	1.31	1.40	1.98	1.94	1.91	1.96	2.00	1.92
Na	0.45	0.69	0.49	0.25	0.41	0.41	0.34	0.23	0.36
Κ	0.07	0.07	0.07	0.07	0.07	0.07	0.17	0.13	0.15
Ti	0.0	0.0	0.0	0.02	0.02	0.06	0.06	0.05	0.06
Okt.	5.14	5.28	5.29	5.04	4.71	5.00	4.63	4.96	4.89
Alk.	2.23	2.07	1.96	2.30	2.42	2.39	2.47	2.36	2.43
R^{+3}	17.7	18.9	18.1	16.6	18.5	11.7	15.1	20.0	22.5
$\mathrm{Fe^{+2}}$	21.0	20.1	16.3	32.7	31.6	29.6	36.8	37.0	34.8
Mg	61.3	61.0	65.6	50.7	49.9	58.7	48.1	43.0	42.7
		Bi 22			Cal 21			Frod. 1	
SiO,	48.5	47.9	54.0	45.5	45.1	44.0	43.3	43.5	43.3
Al,Õ,	13.0	13.0	5.5	13.6	13.2	14.4	18.1	18.4	18.0
FeO	12.4	12.4	9.7	12.9	13.9	14.7	15.5	15.8	14.8
MgO	12.5	11.8	16.8	11.2	11.2	10.6	8.8	8.5	9.3
CaO	11.6	11.9	13.1	12.7	12.7	12.7	10.4	10.5	11.2
Na ₂ O	1.2	1.0	0.9	1.1	1.2	1.1	1.4	1.4	1.4
K _a Õ	.0.3	0.3	0.3	0.4	0.4	0.4	0.0	0.0	0.0
TiO_2	0.2	0.2	0.2	0.6	0.3	0.3	0.4	0.4	0.4
	99.7	98.5	100.2	98.0	98.0	98.2	98.0	98.6	98.5
Si	6.84	6.85	7.48	6.51	6.59	6.45	6.31	6.31	6.28
Al^{IV}	1.16	1.15	0.52	1.49	1.41	1.55	1.69	1.69	1.72
Al^{VI}	1.00	1.04	0.38	0.86	0.87	0.94	1.42	1.45	1.35
Fe^{2+}	1.46	1.48	1.12	1.70	1.70	1.80	1.89	1.91	1.79
Mg	2.63	2.51	3.47	2.44	2.44	2.31	1.91	1.83	2.01
Ca	1.75	1.82	1.94	1.99	1.99	1.99	1.69	1.63	1.74
Na	0.33	0.28	0.24	0.31	0.34	0.31	0.40	0.39	0.39
K	0.05	0.05	0.0	0.07	0.07	0.07	0.0	0.0	0.0
Ti	0.02	0.02	0.02	0.07	0.03	0.03	0.04	0.04	0.04
Okt.	5.11	5.05	4.99	5.07	5.04	5.09	5.27	5.25	5.21
Alk.	2.13	2.15	2.18	2.38	2.40	2.38	2.02	2.02	2.13
\mathbf{R}^{+3}	20.0	20.9	8.0	18.2	17.8	19.1	27.7	28.4	26.8
F ,e+5	28.6	29.3	22.5	33.6	33.7	35.4	36.1	36.7	34.7
Mg	51.4	49.7	69.5	48.2	48.4	45.5	36.2	34.9	38.6

.

	\mathbf{Fro}	d. 2		Jo 545			Sci 123		ŝ	Spl 104 k)
SiO.	43.2	41.7	50.0	51.8	48.8	47.1	46.7	46.7	55.5	54.7	51.9
Al.Ô.	18.8	18.1	9.8	7.6	11.0	8.9	9.4	8.9	4.0	5.2	6.2
FeO	15.1	18.4	6.5	6.5	7.5	14.3	15.4	15.4	10.1	10.1	10.7
MgO	7.9	6.3	18.7	17.0	17.0	11.9	11.9	11.2	14.9	14.9	14.8
CaO	10.9	10.1	12.6	12.2	12.7	11.8	12.3	11.7	11.3	10.3	10.5
Na ₀ O	1.4	1.2	0.8	1.2	1.0	1.0	0.8	1.3	1.4	1.2	1.4
Ko	0.0	0.0	0.4	0.4	0.4	0.1	0.1	0.1	0.3	0.3	0.3
TiÔ.	0.0	0.0	0.3	0.3	0.3	0.3	0.3	0.6	0.0	0.0	0.0
2	96.3	95.7	99.1	97.0	98.7	95.4	96.9	95.9	97.5	96.7	95.8
Si	6.27	6.30	6.95	7.32	6.85	7.05	6.93	7.00	7.86	7.78	7.53
Aliv	1.73	1.70	1.05	0.68	1.15	0.95	1.07	1.00	0.14	0.22	0.47
AlVI	1.54	1.53	0.55	0.59	0.67	0.62	0.57	0.57	0.53	0.66	0.59
Fe ⁺²	1.87	2.33	0.75	0.77	0.88	1.79	1.91	1.93	1.20	1.20	1.30
Mø	1.74	1.42	3.87	3.58	3.55	2.65	2.63	2.50	3.14	3.16	3.20
Ca	1.73	1.64	1.87	1.85	1.91	1.89	1.95	1.88	1.71	1.57	1.63
Na	0.4	0.35	0.22	0.33	0.27	0.29	0.23	0.38	0.38	0.33	0.39
K	0.0	0.0	0.07	0.07	0.07	0.02	0.02	0.02	0.05	0.05	0.06
Ti	0.0	0.0	0.03	0.03	0.03	0.03	0.03	0.07	0.0	0.0	0.0
Okt_	5.17	5.27	5.20	4.97	5.13	5.09	5.14	5.07	4.87	5.02	5.09
Alk	2 13	1.99	2.16	2.25	2.25	2.20	2.20	2.27	2.15	1.96	2.08
R^{+3}	29.8	29.0	11.2	12.5	13.6	12.8	11.7	12.6	10.9	13.1	11.6
Fe^{+2}	36.4	44.1	14.5	15.5	17.2	35.1	37.1	38.1	24.6	23.9	25.5
Mo	33.8	26.9	74.3	72.0	69.2	52.1	51.1	49.3	64.5	62.9	62.9
	0010	-010									
		Spl 119			Toce 8	82a			Tren	n. 4	
SiO.	43.3	45.0	42.5	42.	7 42.7	44.	8	42.3	43.8	43.9	44.6
Al	12.3	12.5	12.7	11.	6 12.8	s 11.	5	17.2	17.5	17.7	17.2
FeO	17.4	17.4	17.6	18.	0 18.0	18.	2	18.2	18.0	18.7	18.6
MgO	9.5	9.8	9.6	9.	2 9.2	9.	3	7.1	6.6	6.8	7.0
CaO	12.5	12.5	12.0	12.	8 12.4	12.	8	10.6	10.6	10.4	10.7
Na _o O	1.2	1.1	1.4	1.	2 1.0) 1.	0	1.6	1.9	1.8	1.4
K ₀	0.8	0.8	0.9	1.	1 1.1	. 1.	0	n.d.	n.d.	n.d.	n.d.
TiO.	0.5	0.5	0.5	0.	6 0.5	6 0.	5	0.4	0.4	0.4	0.4
2	97.5	99.6	97.2	97.	2 97.7	99.	.1	97.4	98.8	99.7	99.9
Si	6.52	6.59	6.43	6.	50 6.4	14 6.	.65	6.30	6.40	6.37	6.45
AIIV	1.48	1.41	1.57	1.	50 1.5	56 I.	.35	1.70	1.60	1.63	1.55
AIVI	0.70	0.75	0.70	0.	58 0.7	12 0.	.66	1.32	1.41	1.40	1.38
Fe^{+2}	2.19	2.13	2.23	2 .	29 2.2	27 2.	.26	2.27	2.20	2.27	2.25
Mo	2.13	2.14	2.16	2.	09 2.0)7 2.	.05	1.58	1.44	1.47	1.51
Ca	2.01	1.96	1.94	2.	.09 2.0	00 2.	.03	1.69	1.66	1.62	1.66
Na	0.35	0.31	0.41	0.	35 0.2	29 0.	.29	0.46	0.54	0.51	0.39
K	0.15	0.15	0.17	0.	21 0.2	21 0.	.19	0.0	0.0	0.0	0.0
Ti	0.06	0.06	0.06	0.	.07 0.0	06 0.	.06	0.04	0.04	0.04	0.04
Okt	5.07	5.08	5.14	5.	.02 5.1	1 5.	.02	5.22	5.12	5.20	5.19
Alk	2.52	2.42	2.53	2	.65 2.4	51 2	.51	2.15	2.20	2.12	2.05
R^{+3}	14.8	15.9	14.6	12	.9 15.	1 14	.2	26.2	28.5	27.7	27.4
Fe^{+2}	43.2	42.0	43.3	45	.6 44.4	44	.9	43.7	43.4	44.1	43.6
Mg	42.0	42.1	42.1	41.	5 40.8	5 40.	.9	30.1	28.1	28.2	29.0
0											

	Trei	m. 6		Trem. 8			Γrem. 9		W	urz 121	e
SiO_2	41.4	41.0	43.6	43.3	44. 0	43.8	43.4	43.2	42.8	40.9	42.1
$Al_2 \tilde{O}_3$	17.8	18.0	15.7	15.8	16.7	17.3	17.2	17.2	13.0	13.1	13.3
FeO	24.6	24.3	16.2	17.8	17.7	20.3	20.5	21.1	15.1	16.1	15.7
MgO	3.6	3.7	8.0	8.0	8.0	6.1	6.5	5.6	9.6	10.0	9.9
CaO	10.0	10.0	10.2	10.5	11.3	10.0	10.2	9.9	12.3	11.8	11.9
Na_2O	1.4	1.4	1.8	1.9	1.7	1.9	1.8	1.9	1.7	1.7	1.5
K_2O	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	1.4	1.5	1.4
TiO_2	0.3	0.3	0.5	0.5	0.4	0.3	0.3	0.4	1.2	1.4	1.3
	99.1	98.7	96.0	97.8	99.8	99.7	99.9	99.3	97.1	96.5	97.1
Si	6.23	6.19	6.51	6.41	6.37	6.40	6.35	6.37	6.43	6.24	6.34
Aliv	1.77	1.81	1.49	1.59	1.63	1.60	1.65	1.63	1.57	1.76	1.66
Alvi	1.38	1.40	1.27	1.17	1.23	1.38	1.31	1.36	0.74	0.60	0.71
$\mathrm{Fe^{+2}}$	3.09	3.10	2.02	2.20	2.14	2.48	2.51	2.60	1.90	2.05	1.98
Mg	0.81	0.83	1.78	1.76	1.73	1.33	1.42	1.23	2.15	2.27	2.22
Ca	1.61	1.62	1.63	1.67	1.75	1.57	1.60	1.56	1.98	1.93	1.92
Na	0.41	0.41	0.52	0.55	0.48	0.54	0.51	0.54	0.50	0.50	0.44
K	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.27	0.29	0.27
Ti	0.03	0.03	0.06	0.06	0.04	0.03	0.03	0.04	0.14	0.16	0.15
Okt.	5.34	5.35	5.16	5.22	5.16	5.24	5.28	5.25	4.92	5.09	5.05
Alk.	2.02	2.03	2.15	2.21	2.23	2.10	2.11	2.11	2.74	2.72	2.63
\mathbf{R}^{+3}	26.5	26.8	25.8	23.5	24.6	27.0	25.5	26.8	17.7	15.0	16.9
${ m Fe^{+2}}$	58.4	57.6	39.7	42.7	42.0	47.6	47.7	49.8	38.6	40.4	39.1
Mg	15.1	15.6	34.5	33.8	33.4	25.4	26.8	23.4	43.7	44.6	44.0

								C	Cal	lcic	A	mj	phi	ibc	ole	s a	n	1 A	٩m	npł	nib	oli	ite	s f	ro	m t	he		ep	on	tir	ıe	Al	\mathbf{ps}					13	9					
	\mathbf{Cr}	500 900	2002	$< 10^{-10}$	$<\!20$	170	280	400	130	130	875	500	500	500	500	500	0002	000	00	270	380	70	550	34	15	00	02	80	120	450	350	200	10	16	650	850	100	300	500	160	170	009	006	080	500 500
	\mathbf{Mn}				300			000	\200 \2000	<200	950														340							0 A F	001									3200	2400	2800	54UU 2000
	\mathbf{Sr}	14	30	40	09	20	18	25	35	35 20	245	10	13.5	22	15.5	18.5	2 87 7	10.0	21 305	20	15	25	20	24	25	55 9 5	21	24	12	$<\!10$	500	360	35 000	23 23	15	13	19	11	11	400		50	130	190	40 30
	Co	< 50	150	- 00 14	10	120	65	30		<10	45	40	52	40	50	48 8 %	0 Q	48	41 19	200	< 50	< 50	100	25	10	70	08	>200	75		< 50	140	01 V	ور 11	25	80	25	200	200	< 50	< 50	55	50	5 X	90 20
	Ni	380	100 60	n I	$<\!10$	300	96	80	15	20 650	240	500	500	500	500	500	0/1	000	07	77	20	55	145	61	10	09 /	200 722	300	230	140	250		10	13	250	250	190	130	450	130	40	430	165	200	150 30
nts	\mathbf{Zr}	105	100	88 88	< 50	180	140	100	0° 20	CB 0	са. 00 65	60	100	140	80	150	150	145	105 295	110	100	80	170	80	< 50	200	130	230	140	< 100	80	130		48	110	78	< 100	50	220	са. 45	са. 50	170	110	170	240 120
vce eleme	Υ				$<\!10$			č	20 2	15	15) ŧ													<10							C F	<10										50	11	20 70
le 7. Tr_{d}	Cu				$<\!10$			1	15	I 4	29	Ì													9							c	x									58	21	54	48 15
Tab	Λ	300	350 960	140	400	250	360	350	> 1000	>1000	245	50	110	150	45	120	300	0/1	010 960	370	350	450	350	360	40	350	500 500	400	350	250	250	330 3	30	2 2	100	160	110	250	250	>500	150	250	300	250	250 360
	\mathbf{Pb}				120				100	60	10	9													110							•	<10									œ	65	0	$52 \times \bigcirc $
	В	19	20	29	< 10	250	28	18	$\frac{40}{2}$	25 60	01 V	0	67	20	5	10	10	2 2	202	30 230	61	30	20	50	< 80	26 90	280	450	230	18	10	450	08⊗ ⊗	и 	2.2	11	4	>500	250	19	14				≪80
	\mathbf{Be}	$\overset{\mathbf{c}}{\vee}$	ы к V	17	က	7 7	1	01 k V	25 25	30 A	~ ~		I	I		c ≀ ∘	о -		- 0	1 C V	a ∕ ∕	, eo	က	4		നറ	° . ∕	°. ∕ ∕	; ; ; ;	51 V	< 3	;; √	-] c ∨	- v	\sim	, vo	7 7	\sim 1	\sim	$\stackrel{<}{\sim}$	$\overset{\circ}{\sim}$		4		$\overline{\vee}$
	Ba	ca. 100	1001	001 ×	<100	100	<100	< 30		120	30	1200	0	0	0	0	0	197	105	001 >	ca. 100	350	$<\!30$	100	< 30	100	140	< 100	< 100	100	150	140	06 V	001/	303030	< 100	def.	120	< 100	$<\!100$		0 6	06 $>$	06 V	06 ∨ ∨
	Sample	AS 912	AS 941 Blon 48h	Bni 176	$\operatorname{Brg} 2$	Cal 25	Cal 29	Cal 30		Ct 19 «S» FK 644	HoVo	Hu 1015	Hu 1026 I	Hu 1026e I	Hu 1026 II	Hu 1026e II	Hu 1164d	0/11 nH	пи 11/08 ₃ Н., 1909 с	Luk 24	Mas 9z	Mas 19	Mera 33a	Mera 34	Mera 40	Mto 798a Mto 708b	Mto 823	Pior 9	PK 834	Riv 17	ShLz A 22a	Spl 125	TV 1160	Varzo 7	Vz 194	Vz 394	Vz 428	Vz 631	Vz 673	Wurz 138	Wurz 150	Wurz 175	Wurz 176	Wurz 177b	20.7.69.01 22.7.69.02

.

Sample		ŭ	ĸ			d'u	ŝ			u	×		$2 V_x$
AS 912	1.653	1.651	1.654		1.663	1.661	1.665		1.673	1.669	1.673		761/2-78
AS 941	1.663	1.664	1.664		1.675	1.675	1.675		1.683	1.684	1.684	1.604	72^{-75}
Blen 48b	1.655	1.655	1.651		1.665	1.665	1.659		1.675	1.675	1.669		83 - 86
$\operatorname{Brg}75$	1.656	1.656	1.628		1.665	1.666	1.667	1.641	1.673	1.674		1.651	78 - 83
Cal 25	1.655	1.655	1.655		1.665	1.667	1.667		1.672	1.673	1.675		72 1/2 - 74 1/2
Cal 29	1.665	1.663	1.663		1.667	1.667	1.667		1.683	1.685	1.683		$67 - 69 \frac{1}{12}$
Cal 30	1.645	1.637	1.651		1.657	1.653	1.665		1.672	1.659	1.675		$80 - 81\frac{1}{2}$
FK 644	1.683		1.681		1.706	1.707	1.699		1.7115		1.706		$46 \frac{1}{2} - 48$
Frod 1	1.656	1.652	1.653	1.652	1.669	1.664	1.664		1.681	1.673	1.674	1.673	80^{-87}
H_{0V0}	1.638	1.640	1.640		1.646	1.646			1.660	1.660			102
Hu 1164d	1.638	1.645	1.645	1.639	1.650	1.657	1.656	1.651	1.660	1.667	1.667	1.662	$82 - 83 \frac{1}{2}$
Hu 1170		1.645	1.639	1.643		1.655	1.651	1.653	1.660	1.665	1.661	1.661	$79 - 81^{-2}$
Hu 1170a ₃		1.665	1.667	1.666		1.678	1.679	1.679	1.680	1.686	1.689	1.687	60 - 65
Hu 1202a		1.678	1.676			1.690	1.671	1.691		1.696	1.697	1.697	53 - 56
Luk 24	1.635				1.647	1.651			1.657				81 - 82%
Mas 9z	1.664	1.658	1.660	1.644	1.674	1.672	1.672	1.677	1.672	1.680	1.680	1.683	64
Mas 19	1.671	1.665	1.665		1.685	1.681	1.681		1.691	1.685	1.685		$58 \frac{1}{2} - 60 \frac{1}{2}$
Mera 33a	1.655	1.647	1.653		1.667	1.663	1.663		1.673	1.671	1.675		$72 \frac{1}{2} - 80$
Mera 34	1.667	1.673	1.680	1.671	1.685	1.683	1.689		1.687	1.691	1.693		$60^{\circ}-63$
Mis 52c	1.624	1.622	1.624	1.628	1.635	1.642	1.634	1.642	1.642	1.652	1.642	1.651	61 - 77
Mto 798a	1.657	1.657	1.655		1.669	1.669	1.667		1.677	1.679	1.677		$74 \frac{1}{2} - 76$
Mto 798b	1.657	1.657	1.655		1.670	1.669	1.667		1.679	1.677	1.675		$74^{-}-74^{-}_{8}$
Mto 823	1.658	1.653	1.653	1.652	1.669	1.667	1.667	1.663	1.679	1.675	1.673	1.673	71
Pior 9	1.657	1.651	1.655		1.667	1.663	1.667		1.679	1.675	1.677		82 - 85
PK 834	1.645	1.643	1.645		1.659	1.657	1.658		1.667	1.667	1.667		$79 \frac{1}{2}$
Riv 17	1.655	1.657	1.658		1.669	1.667	1.668		1.677	1.673	1.677		$75 \overline{1_2} - 77$
Splug 11d	1.653	1.653			1.666	1.663			1.674	1.676			$71^{-}-72$
Trem 4	1.664	1.662	1.666	1.666	1.674	1.677	1.678	1.678	1.682	1.684	1.684	1.685	75 - 79
Trem 6	1.667	1.668	1.672	1.668	1.676	1.678	1.686	1.681	1.687	1.689	1.694	1.689	72 - 76
Trem 8	1.659	1.658	1.658	1.659	1.667	1.668	1.670	1.671	1.679	1.680	1.680	1.681	77 - 78
Trem 9	1.663	1.664	1.663	1.665	1.673	1.674	1.675	1.673	1.682	1.683	1.684	1.683	69 - 76
Vz 631	1.639	1.638	1.639		1.653	1.651	1.649		1.663	1.661	1.659		$82 \frac{1}{2} - 84 \frac{1}{2}$
VZ 673	1.623				1.645				1.659				84 - 87
Wurz 138	1.643	1.649	1.647	1.649	1.655	1.659	1.656	1.657	1.667	1.671	1.656	1.667	$81\frac{1}{2}-83$
Wurz 150	1.655	1.643			1.662	1.663?			1.676	1.671			$73\frac{1}{2} - 76\frac{1}{2}$

Table 8a. Optical data of calcic amphiboles

140

E. Wenk, H. Schwander and W. Stern

$2 V_x$	$-841/_{2}$	- 81	- 85	$\frac{1}{2}-83$	1^{1}
	8 8 8	08 08 08	80	81 80 80	82 84 25
	1.649		1.659		
Ś	1.641	1.643 1.644	1.647	$1.641 \\ 1.639$	1.639 1.651 1.643
u	1.643	1.643 1.642	1.653 1.628	1.643 1.639	1.639 1.653 1.647
	$\begin{array}{c} 1.630 \\ 1.627 \\ 1.648 \\ 1.653 \end{array}$	1.644	1.650 1.629	1.641 1.639	1.638 1.652 1.642
	1.639		1.651		
ŝ	1.632	$1.631 \\ 1.628$	1.638	$1.631 \\ 1.627$	1.627 1.640 1.631
ľu	1.633	$1.630 \\ 1.633$	$1.642 \\ 1.616$	$1.631 \\ 1.627$	1.627 1.641 1.635
	$1.616 \\ 1.616 \\ 1.634 \\ 1.634 \\ 1.641$	1.632	1.637 1.616	1.631 1.629	$1.627 \\ 1.639 \\ 1.623$
	1.623		1.640		
×	1.616	1.617 1.617	1.619	1.616 1.612	1.611 1.628 1.619
й	1.617	1.617 1.616	$1.628 \\ 1.602$	1.615 1.612	$1.611 \\ 1.630 \\ 1.621$
	$\begin{array}{c} 1.607 \\ 1.605 \\ 1.622 \\ 1.630 \end{array}$	1.620	$1.626 \\ 1.601$	1.615 1.6125	1.611 1.627 1.617
Sample	CT 19 «L» CT 19 «S» Hu 1015 Hu 1026 I	Hu 1026e I Hu 1026 II	Hu 1026e II Mera 40	Varzo 6 Varzo 7	Vz 194 Vz 394 Vz 428

Table 8b. Optical data of tremolitic/actinolitic hornblendes

.

${f Hornblendenden}$	+-	++	+		+ -	-+-	+	+	+ -	+	+ -}	+-	+	+	+	+		- +	+	+	+	+ -	┝	+	+	+	+	++
Коск	+ +	⊢		+	+	+				-	 -					+					+	+	-+		+			
\mathbf{B} iotite						+		+		-	+-	-+-						+		+		+					+	
Coordinates	658.6/135.6 650 2/115 1	717.7/157.3	775.4/143.9		701.3/146.6	729.8/143.5	729.3/134.8	730.7/123.7	698.8/147.4	691.0/142.7		675.2/123.4	678.7/125.2	678.1/124.9	678.8/125.8	674.7/123.4 631 A/ OR 9	631 4/ 96 2	714.9/151.5	766.1/124.2	767.2/123.5	750.6/129.1	754.9/116.6	734 6/145 1	734.6/145.1	734.6/145.1	687.3/147.6	687.3/147.6	694.8/145.4 $694. \ /153.5$
malysed samples (Tables 2–4) Location	Giessen, Binntal Simnlon Alte Kasama	Loderio	ESE Grevasalvas	Muretto	Muretto A. Cadonigo	A. Alögna	W Sta. Domenica	New Calanca-road	A. Campolungo	W F. Mascarpino	R. del Motto	A. Bosa, V. di Campo	N Cma. di Tramolino, V. di Campo	S Motta del Termine, V. di Campo	SE Mattignello, V. di Campo	SE A. Bosa, V. dı Campo Pfulwe	Pfilwe	NNW Aquila	Corte Vecchia, Bagni del Masino	Quarry E Bagni del Masino	Berzo S Prato/Chiavenna	S Motta dei Cecchi M40 Decebione	S Bernardino	S Bernardino	S Bernardino	Campo la Torba	Campo la Torba	W Colla near Fusio W Staz. Piora
Table 9a. <i>List of a</i> Rock	Oligoclase-amphibolite Oligoclase-amphibolite	Epidote-andesine-amphibolite	Albite-crossite-schist	Andesine-amphibolite	Chuorue-ongociase-ampinonue Green tremolite in dolomite-marble	Biotite-albite/oligoclase-amphibolite	Epidote-oligoclase-amphibolite	Biotite-oligoclase-andesine-amphibolite	Light-green tremolite in dolomite-marble	Wuaruz-calche-magnente-normblende-rock Chlorita hiotita remot emoticalita	Hornblendite	Inclusion in dolomite-marble	Inclusion in hornblende-plagioclase-gneiss	Epidote-oligoclase-amphibolite	Inclusion in oligoclase-amphibolite	Uligociase-amphibolite Prasinite	Hornhlende-albite vein	Biotite-oligoclase/andesine-amphibolite	Pyroxene-andesine/labradorite-amphibolite	Biotite-hornblende-andesine/labradorite-gneiss	Andesine/labradorite-amphibolite	Epidote-biotite-hornblende-andesine-gneiss Tramolita dioneida marblo	Calcite-clinozoisite-biotite-chlorite-amphibolite	Garnet-biotite-chlorite-amphibolite	Garnet-muscovite-amphibolite	Oligoclase-amphibolite	Biotute-oligoclase-amphibolite	Uligoclase-amphibolite Epidote-oligoclase-amphibolite
Sample	AS 912 AS 941	Blen 48 b	Bni 258	$\operatorname{Brg} 73$	Drg /0 Cadonigo	Cal 25	Cal 29	Cal 30	CT 19 EV 644	г. 044 С. Т. 580	HoVo	Hu 1015	Hu 1026	Hu 1164d	Hu 1170	ни 1202а КАW 657	KAW 658	Luk 24	Mas 9z	Mas 19 11 80	Mera 33a	Mera 34 Were 40	Mis 52	Mis 52b	Mis $52c$	Mto 798a	Mto 798b	Mto 823 Pior 9

Table 9a. List of analysed samples (Tables 2-4)

142

E. Wenk, H. Schwander and W. Stern

+	+				+	+	+	╋	+	+	+	+	+	+	+		+	+		+	+	+			+				+-	
	+	+	+	+	+		+-						+	+		+			+	+	+	+	+.	+	+	+	+	+	+	+
									+	+	+	+	+				+													
683.1/113.5	722.7/119.9	762.7/131.3	764.0/132.3	774.2/121.9	707.8/124.1	646.4/124.1	749.4/132.2	697.3/146.8	660.2/119.9	660.2/119.9	709.4/125.2	702.6/129.9	709.7/118.6	707.1/122.3	699.8/133.8	707.4/135.6	700.1/119.5	703.3/116.9	699.4/116.9	718.3/116.1	717.5/115.7	717.7/116.4	712.26/115.72	729.9/117.6	729.9/119.9	731.1/120.4	734.0/125.5	736.9/130.3	736.9/130.3	735.1/157.4
S Cma. del Sassone. Centovalli	NNW Gorduno, Riviera	Denc dal Luf	Bondasca, SW Gerp	Preda Rossa	Lavertezzo	Quarry Vogelsang	Liro W Chiavenna	Campolungo	E Ciöina di dentro	E Ciöina di dentro	Mte. Sambuco	NW A Tencio, V. Verzasca	Fontöbbia	Cargello near Corippo	A. Mugaglia, V. Redorta	Madone Grosso	Aurigeno-Dunzio	Brè-Monteggia (Locarno)	Riei, Verscio	W Sementina	San Defendente	N S. Defendente	Mti di Ditto	Mti. Laura	Mti. Laura	V. Traversagna	Quarry Cama	A. Scimetta, V. Mesocco	A. Scimetta, V. Mesocco	Valserberg
Homhlanda-andasina-maiss	Rnidote-dionside-andesine-amphibolite	Andesine/labradorite-amphibolite	Hornblendite	Enidote-biotite-hornblende-andesine-gneiss	Bytownite-amphibolite, inclusion in gneiss	Oligoclase-amphibolite	Clinozoisite-andesine-biotite-hornblende-schist	Tremolite in white dolomite-marble	Actinolite-phlogopite-calcite-oligoclase-gneiss	Actinolite-phlogopite-calcite-oligoclase-gneiss	Biotite-schist	Biotite-actinolite-schist	Actinolite-biotite-schist	Labradorite/bytownite-amphibolite	Hornblendić inclusion in oligoclase/andesine amphibolite	Andesine-amphibolite	Biotite-andesine/labradorite-amphibolite	Pyroxen-andesine-amphibolite	${f E}$ pidote-andesine-amphibolite	Titanite-quartz-andesine-amphibolite	Andesine-amphibolite	Andesine/labradorite-amphibolite	Pyroxene-bytownite-amphibolite	Andesine/labradorite-amphibolite	Andesine/labradorite-amphibolite	Andesine-amphibolite	Oligoclase-quartz-chlorite-amphibolite	Biotite-quartz-andesine-amphibolite	Biotite-andesine-amphibolite	Calcite-chlorite-epidôte-albite-schist
DK 834	LIX 001 Riv 17	Sci 171	Sci 186	Sci 244	Sh Lz A 22a	Spl 125	Spluga 11d	TV 116e	Varzo 6	Varzo 7	Vz 194	Vz 394	Vz 428	Vz 631	Vz 673	Vz 703	Wurz 138	Wurz 150	Wurz 166	Wurz 175	Wurz 176	Wurz 177b	Wurz 179	19.7.69.01	20.7.69.01	20.7.69.02	20.7.69.03	22.7.69.01	22.7.69.02	29.9.69.01

Calcic Amphiboles and Amphibolites from the Lepontine Alps

143

Sample	Rock type	Locality	Coordinates
Ad 36e	albite amphibolite, chlor. biot. musc. rut. ore bearing	Hinterrhein	728.2 /151.1
AS 907	clinoz. oligocl. amphibolite, tit. biot. bearing	Merezenbachtal	$666.4 \ / 145.8$
Bi 22	biot. epid. oligocl. amphibolite, rut. bearing	Alp Freiche	662.7 $/136.45$
Cal 21	andesine, amphibolite, epid. tit. bearing	W Santa Domenica	$728.7 \ / 134.8$
Frod 1	kyanite biotite garnet hornblende-schist	Frodalera	$706.7 \ /154.2$
Frod 2	hornblende garnet biot. schist	Frodalera	706.7 /154.2
Jo 545	biot. andes./labrad. amphibolite, tit. rut. epid. bearing	N Lago Devero	671.27/134.75
Sci 123	andesine amphibolite, ore bearing	Forno	774.45/136.65
Spl 104 b	chlor. epid. albite/oligocl. amphibolite, tit. bearing	S Alpe Straciugo	652.6 /108.0
Spl 119	albite/oligocl. epid. amphibolite, biot. tit. bearing	below Lago di Paione	$658.0 \ / 113.2$
Toce 82a	oligoel. amphibolite, tit. bearing	quarry SE Croppo	667.8 /106.8
Trem 4	garnet biotite chlor. hornblende-schist	Motto Bartola	$687.9 \ / 154.4$
Trem 6	biot. chlor. garnet hornblende-schist, staurolite bearing	Motto Bartola	687.9 /154.4
Trem 8	chlor. hornblende plagiocl. gneiss epid. bearing	Motto Bartola	687.9 /154.4
Trem 9	garnet hornblende schist, carbonate bearing	Lago Ritom	$695.5 \ / 155.3$

Table 9b. Amphiboles, analyzed by microprobe (Table 6)

Table 10. Grouping according to metamorphic grade

		Ana	lyses
Type	An content	\mathbf{Rock}	Mineral
greenstones (diabase a. o.)		29	
hornblende-bearing greenschists, prasinites		10	6
albite-amphibolites	0 - 5	9	8
albite-plus oligoclase-, or oligoclase-amphibolites	(0-5)+(10-25) or $17-30$	17/37	18
andesine-amphibolites	30 - 50	25	18
labradorite/bytownite-amphibolites	> 50	8	6
Garbenschiefer and oligoclase-amphibolites (Gotthard, Frodalera)	20 - 30	8	30
	Type greenstones (diabase a. o.) hornblende-bearing greenschists, prasinites albite-amphibolites albite-plus oligoclase-, or oligoclase-amphibolites andesine-amphibolites labradorite/bytownite-amphibolites Garbenschiefer and oligoclase-amphibolites (Gotthard, Frodalera)	TypeAn contentgreenstones (diabase a. o.) hornblende-bearing greenschists, prasinites albite-amphibolites $0-5$ albite-amphibolites $0-5$ albite-plus oligoclase-, or oligoclase-amphibolites $(0-5)+(10-25)$ or $17-30$ andesine-amphibolites $30-50$ labradorite/bytownite-amphibolites >50 Garbenschiefer and oligoclase-amphibolites $20-30$	TypeAn contentAna. Rockgreenstones (diabase a. o.)29hornblende-bearing greenschists, prasinites10albite-amphibolites $0-5$ 9albite-plus oligoclase-, or oligoclase-amphibolites $(0-5)+(10-25)$ or $17-30$ $17/37$ andesine-amphibolites $30-50$ 25labradorite/bytownite-amphibolites >50 8Garbenschiefer and oligoclase-amphibolites $20-30$ 8

le.
rac
g
hic
rp
n0
an
nei
1 2
ini
as
re
ine
\dot{q}
an
a.
era
rit
10
ca
hi
dκ
gre
roi
et:
t c
1 te
nG.
p
<i>or</i>
ncc
g c
i.
d
<i>10</i> ,
G
8.
ole
\dot{q}
hc
lu
a
ic
ulc
3
fo
68
ци
va
al
tic
isi
tat
\mathcal{O}
а.
11
0
Įď
T_{a}

Weight percentages

					n evgn	n percenu	nges				
Group	Rock type		SiO_2	$\mathrm{Al}_{2}\mathrm{O}_{3}$	${\rm Fe_2O_3}$	FeO	MnO	MgO	CaO	Na_2O	$\mathbf{K_2O}$
I	Greenschists (prasinites)	$\mathbf{V}_{\mathbf{x}}^{\mathbf{X}}$	6 48.18 2.05 4.3	$6 \\ 8.20 \\ 2.26 \\ 27.6$	$6 \\ 3.83 \\ 1.14 \\ 29.7$	$6 \\ 8.88 \\ 1.50 \\ 16.9$		$6\\15.10\\2.87\\19.0$	$6 \\ 10.82 \\ 1.10 \\ 10.2$	$6 \\ 1.95 \\ 0.50 \\ 25.9$	$\begin{array}{c} 6 \\ 0.15 \\ 0.14 \\ 0.14 \\ 91.9 \end{array}$
П	Albite-amphibolites	$\mathbf{V}_{\mathbf{x}}^{\mathbf{S}} \mathbf{N}_{\mathbf{x}}$	8 45.65 4.03 8.8	$8 \\ 3.62 \\ 32.0 \\ 32.$				$8 \\ 9.98 \\ 2.72 \\ 27.3$	$8 \\ 0.52 \\ 4.49$	$\begin{array}{c} 8\\ 2.24\\ 0.80\\ 35.8 \end{array}$	
Η	Hornblende rocks of the zone Tremola-Frodalera	$\mathbf{V}_{\mathbf{x}}^{\mathbf{S}}\mathbf{N}_{\mathbf{x}}$	$30 \\ 42.85 \\ 1.02 \\ 2.4$	$30 \\ 17.09 \\ 0.99 \\ 5.8 $	$13 \\ 4.09 \\ 1.08 \\ 26.3$	$13 \\ 12.85 \\ 1.44 \\ 11.2$	$30 \\ 0.18 \\ 0.11 \\ 0.11 \\ 63.4$	$30 \\ 7.60 \\ 1.60 \\ 21.1$	$30 \\ 10.27 \\ 0.43 \\ 4.2$	$\begin{array}{c} 30 \\ 1.64 \\ 0.28 \\ 16.9 \end{array}$	$egin{array}{c} 13 \\ 0.36 \\ 0.03 \\ 21.2 \end{array}$
IV	Oligoclase-amphi- bolites±albite	$\mathbf{V}_{\mathbf{x}}^{\mathbf{N}}$	$18 \\ 43.79 \\ 2.38 \\ 5.4$	$18\\12.52\\1.18\\9.4$	$18 \\ 4.82 \\ 1.56 \\ 32.4$	$18 \\ 11.46 \\ 2.62 \\ 22.9$	$\begin{array}{c} 18 \\ 0.26 \\ 0.13 \\ 50.9 \end{array}$	$18 \\ 10.66 \\ 2.06 \\ 19.4 $	$18 \\ 10.90 \\ 0.99 \\ 9.1$	$18 \\ 1.59 \\ 0.73 \\ 45.8$	$18 \\ 0.53 \\ 0.26 \\ 49.0$
Λ	Andesine- amphibolite	$\mathbf{V}_{\mathbf{x}}^{\mathbf{N}}$	18 44.77 2.19 4.9	$18\\11.72\\1.16\\9.8$	$18 \\ 3.98 \\ 1.43 \\ 35.8 $	$18 \\ 11.10 \\ 2.77 \\ 24.9 $	$18 \\ 0.32 \\ 0.14 \\ 44.8$	$18 \\ 11.63 \\ 2.46 \\ 21.2$	$18 \\ 11.91 \\ 0.52 \\ 4.3$	$18 \\ 1.19 \\ 0.37 \\ 31.0$	$18 \\ 0.50 \\ 0.25 \\ 50.9$
IA	Labradorite/ bytownite- amphibolite	$\mathbf{v}_{\mathbf{x}}^{\mathbf{x}}$ $\mathbf{v}_{\mathbf{x}}^{\mathbf{x}}$	$6 \\ 44.75 \\ 1.93 \\ 4.3$	$6 \\ 11.63 \\ 1.02 \\ 8.8$	$\begin{array}{c} 6\\ 3.62\\ 2.25\\ 62.1 \end{array}$	$6 \\ 11.27 \\ 3.03 \\ 26.9$	$\begin{array}{c} 6 \\ 0.27 \\ 0.08 \\ 30.5 \end{array}$	$6 \\ 11.98 \\ 1.61 \\ 13.4$	$\begin{array}{c} 6\\ 12.05\\ 0.65\\ 5.4\end{array}$	$\begin{array}{c} 6 \\ 1.10 \\ 0.50 \\ 45.3 \end{array}$	$\begin{array}{c} 6\\ 0.53\\ 0.35\\ 65.6\end{array}$

Calcic Amphiboles and Amphibolites from the Lepontine Alps

 H_2O^+ 6 0.58 0.58 24.2

TiO₂ 6 0.42 97.7 $\begin{array}{c} 13\\ 2.22\\ 0.09\\ 4.0\\ 1.8\\ 2.41\\ 1.12\\ 1.12\\ 1.97\\ 1.97\\ 1.97\\ 0.29\\ 0.29\\ 0.32\\ 0.32\\ 16.1\\ 1.98\end{array}$

 $\begin{array}{c} 28\\ 0.45\\ 0.45\\ 0.16\\ 35.4\\ 18\\ 0.96\\ 0.38\\ 39.7\\ 39.7\\ 0.88\\ 0.88\\ 0.88\\ 0.40\\ 0.40\\ 0.45\\ 0.45\\ 0.45\end{array}$

145

				4	\$		•	\$					
Group	Rock type		SiO_2	Al_2O_3	${\rm Fe_2O_3}$	FeO	MnO	MgO	CaO	Na_2O	${ m K_2O}$	TiO_2	$\mathrm{H_2O^+}$
0	Greenstones (diabase)	$\mathbf{v}_{\mathbf{x}}^{\mathbf{x}}$	$\begin{array}{c} 29 \\ 46.47 \\ 5.22 \\ 11.2 \end{array}$	$\begin{array}{c} 29\\ 17.02\\ 1.48\\ 8.7\end{array}$	$\begin{array}{c} 29 \\ 3.67 \\ 1.84 \\ 50.1 \end{array}$	$\begin{array}{c} 29\\ 5.54\\ 2.27\\ 40.9\end{array}$	$\begin{array}{c} 29 \\ 0.17 \\ 0.06 \\ 36.9 \end{array}$	$\begin{array}{c} 29 \\ 6.66 \\ 2.33 \\ 35.0 \end{array}$	$\begin{array}{c} 29 \\ 8.21 \\ 4.27 \\ 52.0 \end{array}$	$\begin{array}{c} 29 \\ 3.99 \\ 1.94 \\ 48.8 \end{array}$	$\begin{array}{c} 29 \\ 0.51 \\ 0.52 \\ 102.5 \end{array}$	$\begin{array}{c} 29 \\ 1.73 \\ 0.59 \\ 34.2 \end{array}$	$\begin{array}{c} 29 \\ 4.72 \\ 1.81 \\ 38.4 \end{array}$
I	Greenschists (prasinites)	$\mathbf{v}_{\mathbf{x}}^{\mathbf{x}}$	$10 \\ 47.45 \\ 3.09 \\ 6.5$	$10\\16.61\\2.20\\13.2$	$10 \\ 3.25 \\ 1.45 \\ 13.2$	$10 \\ 6.80 \\ 2.11 \\ 30.9$	$\begin{array}{c} 10 \\ 0.16 \\ 0.06 \\ 37.5 \end{array}$	$10 \\ 7.60 \\ 2.18 \\ 28.7$	$10 \\ 9.01 \\ 2.66 \\ 29.5$	$10 \\ 3.10 \\ 1.10 \\ 35.5$	$\begin{array}{c} 10 \\ 0.46 \\ 0.45 \\ 98.3 \end{array}$	$10 \\ 2.04 \\ 0.80 \\ 39.5$	$10 \\ 3.08 \\ 1.26 \\ 40.9$
П	Albite-amphibolites	$\mathbf{V}_{\mathbf{x}}^{\mathbf{N}}$ $\mathbf{N}_{\mathbf{x}}^{\mathbf{N}}$	$\begin{array}{c} 9 \\ 46.85 \\ 6.21 \\ 13.3 \end{array}$	$\begin{array}{c} 9 \\ 15.18 \\ 4.11 \\ 27.1 \end{array}$	$9\\3.10\\1.77$ 56.9	$\begin{array}{c} 9\\ 6.85\\ 2.31\\ 33.7\end{array}$	$\begin{array}{c} 9\\ 0.16\\ 0.09\\ 57.1 \end{array}$	$9 \\ 8.11 \\ 3.17 \\ 39.0 $	$\begin{array}{c} 9\\ 10.12\\ 3.13\\ 30.9\end{array}$	$9\\2.68\\0.72\\26.9$	$\begin{array}{c} 9 \\ 1.01 \\ 0.97 \\ 95.9 \end{array}$	$9 \\ 1.51 \\ 1.06 \\ 70.2$	$\begin{array}{c} 9\\1.91\\0.37\\19.5\end{array}$
H	Hornblende rocks of the zone Tremola-Frodalera	$\mathbf{v}_{\mathbf{x}}^{\mathbf{x}}$	8 53.15 3.58 6.7	$8 \\ 3.19 \\ 3.19 \\ 17.2$	$8 \\ 4.21 \\ 2.52 \\ 59.8 $	8 6.77 2.69 39.8	$\begin{array}{c} 8 \\ 0.11 \\ 0.12 \\ 0.12 \end{array}$	8 5.74 1.48 25.8	$8 \\ 6.23 \\ 2.33 \\ 37.5$	$8 \\ 0.58 \\ 32.4$	$8 \\ 1.65 \\ 0.84 \\ 51.0$	$8 \\ 1.42 \\ 0.87 \\ 61.1$	$\begin{array}{c} 8\\ 1.86\\ 0.87\\ 46.5 \end{array}$
IV	Oligoclase-amphi- bolites±albite	$\mathbf{N}_{\mathbf{x}}$ $\mathbf{N}_{\mathbf{x}}$ $\mathbf{N}_{\mathbf{x}}$	$17 \\ 49.01 \\ 2.94 \\ 6.0$	$17 \\ 15.30 \\ 1.24 \\ 8.1$	$17 \\ 2.86 \\ 1.16 \\ 40.7$	$17 \\ 8.62 \\ 3.12 \\ 36.2$	$16 \\ 0.20 \\ 0.12 \\ 59.3$	$37 \\ 7.43 \\ 2.40 \\ 32.4$	$37 \\ 9.20 \\ 1.71 \\ 18.6$	$37 \\ 2.82 \\ 1.12 \\ 39.6$	$37 \\ 0.84 \\ 0.82 \\ 0.7.2 \\ 0.12 \\ 0$	$17 \\ 1.76 \\ 0.70 \\ 39.4$	$17 \\ 2.40 \\ 0.71 \\ 29.6$
Δ	Andesine- amphibolite	$\mathbf{v}_{\mathbf{x}}^{\mathbf{N}}$	25 50.93 4.02 7.9	25 15.37 2.74 17.9	$25 \\ 2.60 \\ 1.39 \\ 53.6$	25 6.46 1.60 24.8	25 0.16 0.06 37.6	$25 \\ 7.97 \\ 3.11 \\ 39.0$	$25 \\ 10.06 \\ 1.91 \\ 19.0$	$25 \\ 2.58 \\ 0.72 \\ 27.9$	$25 \\ 0.84 \\ 0.74 \\ 87.7$	$25 \\ 1.17 \\ 0.48 \\ 0.6$	$25 \\ 1.62 \\ 0.55 \\ 33.9 \\ 33.9 \\ 35$
IV	Labradorite/ bytownite- amphibolite	$\mathbf{V}_{\mathbf{x}}^{\mathbf{X}}$	8 48.69 1.77 3.6	$8 \\ 16.26 \\ 1.46 \\ 9.0$	8 2.38 1.07 45.0	$8 \\ 6.65 \\ 1.43 \\ 21.5$	$8 \\ 0.12 \\ 0.06 \\ 52.4$	$8 \\ 8.50 \\ 2.46 \\ 28.9$	$8 \\ 12.12 \\ 1.41 \\ 11.6$	$\begin{array}{c} 8\\ 1.89\\ 0.72\\ 37.9\end{array}$	$\begin{array}{c} 8\\ 0.53\\ 0.55\\ 103.2 \end{array}$	8 1.47 0.57 38.6	$8 \\ 0.37 \\ 0.37 \\ 29.2$

Table 11b. Statistical values of amphibolites and hornblende bearing rocks. Grouping according to petrographical criteria and increasing metamorphic grade. Base: Weight percentages

146

•

E. Wenk, H. Schwander and W. Stern

Table 12. Average modes of plagioclase-amphibolites with changing An-content, deduced fromthe modes of 85 amphibolites from the Central Alps

An %	0	10	20	25	30	35	4 0	45	50	55	60	65	70	75	80	90
Vol-%												data	less 1	reliab	le	
Plagioclase	17	18	23	31	32	33	34	35	36.5	38	38	34	30	30.5	29	28.5
Hornblende	60	60	60.5	60	60	59.5	58.5	55.5	53.5	50	50	55	57	57	56	56.5
Clinopyroxene						0.5	1.5	4	6	7.5	8	7.5	7	6	8.5	5
Chlorite,																
Biotite	7	6	5	4	3.5	3	2	1	1	0.5					-	
$\mathbf{Klinozoisit}/$																
Epidote	13	11	7	3.5	2.5	1	1	1	0.5	0.5	0.5	0.5	1.5	3.5	2.5	3
Quartz	1	2	2.5	0.5		1	1	1	0.5	1	1	1	1.5	1	1.5	4
Accessories	2	3	2	1	2	2	2	2.5	2	2.5	2.5	2	3	2	2.5	3

Table 13. Approximate modal variation with rising metamorphic grade, calculated in cation %for 3 different amphibolites

1. Si 45.8, Al 15.7, Fe¹¹¹ 1.6, Fe¹¹ 7.4, Mg 13.6, Ca 10.2, Na 4.6, K 0.2, Ti 0.9 (approximate composition of average albite-amphibolite in Table 12).

	Metamorphic grade				
An Plag.	0	33	50		
% Plag.	17.5	26.5	35		
Ho	60	61	38		
Chlor.	7				
Biot.		3.5			
\mathbf{Clzt}	12.5	2			
Clpx			5		
Opx			19		
\mathbf{Ru}	1				
\mathbf{Tit}		1	2		
Ore	1	2	1		
$\mathbf{Q}\mathbf{z}$	1	4			

2. Si 45.3, Al 19.0, Fe¹¹¹ 1.6, Fe¹¹ 6.1, Mg 10.8, Ca 12.4, Na 3.1, K 0.5, Ti 1.2 (sample Wurz. 177b)

An Plag.	0	26	50	60	68
% Plag.	9.5	13.5	19	30	56.5
Ho	66.5	66.5	66.5	55	
\mathbf{Clzt}	18.5	14 [°]	9		
Clpx				9	20
Opx					18.5
\mathbf{Ru}	1	1	1	1	1
Ore	0.5	0.5	0.5	1	4
$\mathbf{Q}\mathbf{z}$	4	4.5	4	4	

3. Si 45.0, Al 19.0, Fe^{III} 3.0, Fe^{II} 4.9, Mg 9.8, Ca 14.1, Na 3.0, K 0.2, Ti 1.0 (sample Wurz. 179)

An Plag.	0	50	67	71
% Plag.	11.5	23	36.5	55.5
Ho	60.5	60	56	
\mathbf{Clzt}	21.5	13		_
Clpx	_		4	21
Opx				16
\mathbf{Tit}	2.5	2.5	2.5	3
Ore	1	0.5	1	4.5
$\mathbf{Q}\mathbf{z}$	3	1		

Si	\mathbf{Al}	\mathbf{Ca}	Na	\sum cat. %	An %
6	2		2	10	0
1	1	0.5			
6	2		2	12.5	20
1.33	1.33	0.67			
6	2		2	13.3	25
2.67	2.67	1.33			
6	2		2	16.7	40
4	4	2			
6	2		2	20	50
6	6	3			
6	2		2	25	60
12	12	6			
6	2		2	40	75
16	16	8			
6	2		2	50	80
36	36	18			
6	2	—	2	100	90

Table 14. Quantitative variation of plagioclase with constant Na content, but changing anorthite proportion. Calculations in cation percentages. Na = 2 cation %

Bibliography

- ANGEL, F. (1967): Über Mineralzonen, Tiefenzonen und Mineralfazies. Rückblicke und Ausblicke. Fortschr. Mineral. 44, 288.
- BINNS, R. A. (1965): The mineralogy of metamorphosed basic rocks from the Willyams Complex, Broken Hill District, New South Wales. Min. Mag. 35, 306.
- DEER, W. A., HOWIE, R. A. and ZUSSMAN, J. (1962): Rock forming Minerals. London.
- DIETRICH, V. (1969): Die Ophiolithe des Oberhalbsteins (Graubünden) und das Ophiolitmaterial der ostschweizerischen Molasseablagerungen, ein petrographischer Vergleich. Bern.
- DE QUERVAIN, F. und FRIEDLÄNDER, C. (1942): Chemismus schweizerischer Gesteine. 1. Nachtrag. Bern.
- DE QUERVAIN, F. und JENNY, V. (1956): Chemismus schweizerischer Gesteine. 2. Nachtrag. Bern.
- FREY, M. (1969): Die Metamorphose des Keupers vom Tafeljura bis zum Lukmanier-Gebiet. Bern. Beitr geol. Karte Schweiz N.F. 137.
- HASLER, P. (1949): Geologie und Petrographie der Sambuco-Massari-Gebirgsgruppe zwischen der oberen Leventina und Valle Maggia im nördlichen Tessin. SMPM 29, 50.
- HUNZIKER, J. C. (1966): Zur Geologie und Geochemie des Gebietes zwischen Valle Antigorio und Valle di Campo. SMPM 46, 474.
- KALKOWSKY, E. (1886): Elemente der Lithologie. Heidelberg.
- KNUP, P. (1958): Geologie und Petrographie des Gebietes zwischen Centovalli-Valle Vigezzo und Onsernone. Diss. SMPM 38, 83.
- Kuno, H. (1968): Differentiation of Basalt Magmas. In: Basalts, vol. 2, New York. John Wiley & Sons, New York, London, Sydney.
- LEAKE, B. E. (1969): The relationship between composition of calciferous amphibole and grade of metamorphism, in: Controls of Metamorphism, 299, Olivers Boyd, Edinburg.
- --- (1968): A Catalog of Analyzed Calciferous and Subcalciferous Amphiboles Together with Their Nomenclature and Associated Minerals. Geol. Soc. Am. Spec. Paper 98.

- NIGGLI, P. (1936): Über Molekularnormen zur Gesteinsberechnung. SMPM 16, 295.
- (1936a): Die Magmentypen. SMPM 16, 335.
- SCHWANDER, H., HUNZIKER, J. und STERN, W. (1968): Zur Mineralchemie von Hellglimmern in den Tessineralpen. SMPM 48, 357.
- STECK, A. und GLAUSER, E. (1968): Universal Drehtisch für optische Untersuchungen von Mineralkörnern. SMPM 48, 815.
- STEIGER, R. (1961): Die Hornblenden der Tremolaserie. Teil I. Chemismus und Dichte der Hornblenden. SMPM 41, 127.
- STERN, W. B. (1966): Zur Mineralchemie von Glimmern aus Tessiner Pegmatiten. SMPM 46, 138.
- --- (1972): Zur röntgenspektrometrischen Analyse von silikatischen Gesteinen und Mineralien. SMPM 52, 1.
- WEIBEL, M. und LOCHER, TH. (1964): Die Kontaktgesteine im Albigna- und Fornostollen (nördliches Bergeller Massiv). SMPM 44, 159.
- WENK, E., SCHWANDER, H., HUNZIKER, J. und STERN, W. (1963): Zur Mineralchemie von Biotit in den Tessineralpen. SMPM 43, 435.
- WENK, E. und KELLER, F. (1969): Isograde in Amphibolitserien der Zentralalpen. SMPM 49, 157.
- WENK, H. R. (1971): Variations of lattice constants in clinoamphiboles. Zschr. Kristallogr. 133, 341.
- WETZEL, R., SCHWANDER, H. und STERN, W. (1974): Hornblenden aus der Albit- bis Albitoligoklaszone zwischen Zermatt und Domodossola. SMPM 54, 151.

Manuscript received January 23, 1974.