

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 55 (1975)

Heft: 3

Artikel: On acid oligoclase in vitrophyre from Alftavik, NE-Iceland

Autor: Wenk, E. / Schwander, H. / Stern, W.

DOI: <https://doi.org/10.5169/seals-43077>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On Acid Oligoclase in Vitrophyre from Alftavik, NE-Iceland

By *E. Wenk, H. Schwander, W. Stern and R. Wetzel**)

With 1 figure

Abstract

Oligoclase phenocrysts Or 6 Ab 79 An 15 from the glassy margin (chemical analysis) of a rhyolitic dyke near Ytri Alftavik in NE-Iceland are described. The optic orientation expressed in Euler angles Φ 95.5°, Θ 69.5°, Ψ 84°, $2V_\gamma$ 126.5° fits high-temperature curves of BURRI-PARKER-WENK. Microlites have similar, slightly more calcic composition (An 17–18) and the residual glass is enriched in potassium and approaches proportions near the alkali-feldspar minimum. Similar vitrophyric rocks from E-Iceland, bearing only one type of slightly zoned feldspar, have been discussed by CARMICHAEL (1960, 1963).

The sample described was collected on the easternmost promontory of Ytri Alftavik in NE-Iceland. The rock forms the glassy margin of a rhyolitic dyke in the basaltic pyroclastic series. The locality is of difficult access and was reached following the itinerary described by THORODDSEN (1906, p. 275).

DESCRIPTION OF ROCK ISL 253: PERLITIC OLIGOCLASE-VITROPHYRE

8 vol % phenocrysts of glomeroporphritic, idiomorphic, well twinned oligoclase An 14–17, up to 2 mm long, with weak zonal structure. 2% phenocrysts of greenish, slightly pleochroic clinopyroxene, $c \wedge n_\gamma$ up to 53°, $2V_\gamma$ 58–60°, $n_\gamma - n_\alpha$ 0.029–0.036, and of Fe-rich olivine, partially or completely replaced by brownish sheet silicates of the chlorite group and of titanomagnetite; zircon is a common accessory.

90% glassy matrix and microlites. Glass brownish, perlitic with incipient devitrification along cracks, n 1.500 ± 0.001. Slightly flow-textured microlites

*) Mineralogisch-Petrographisches Institut der Universität, Bernoullistrasse 30, 4056 Basel.

<0.1 mm of polysynthetically twinned feldspar, brownish clinopyroxene (pigeonite in part) with small optic axial angle, $n_{\gamma}-n_{\alpha}$ 0.029–0.036, axial plane \perp (010) and \parallel (010), and ores. Proportion glass: microlites 6 : 1. Microprobe analyses¹⁾ proved that three selected feldspar microlites have oligoclase composition:

Or	3.3	4.1	1.8
Ab	78.9	79.0	80.4
An	17.8	16.9	17.8

The composition of the glass is difficult to assess. Compared with the total rock it is enriched in K₂O (3.3 wt-%) and impoverished in Na₂O (3.6–4.0) and CaO (0.3–0.5). Silica and alumina values are not reliable.

The XF-analysis of the total rock gave the following result:

	wt-%
SiO ₂	69.8
Al ₂ O ₃	13.0
Fe ₂ O ₃	0.9
FeO	1.9
MnO	0.1
MgO	0.1
CaO	1.3
Na ₂ O	4.9
K ₂ O	2.7
TiO ₂	0.2
P ₂ O ₅	0.1
H ₂ O	5.6
sum	<hr/> 100.6

The rock has granitic composition and represents a rhyolitic vitrophyre. From the analysis the norm Q 26.4, Or 17.0, Ab 46.5, An 5.8, M 4.3 (cation-%) is derived.

OPTICAL AND CHEMICAL PROPERTIES OF THE OLIGOCLASE PHENOCRYSTS

Oligoclase is well twinned after the albite-, Karlsbad-, complex albite-Karlsbad- and acline-laws (in order of decreasing frequency). From U-stage measurements on 8 twin-groups the An-range 14–17 resulted and an average optic axial angle $2V_{\gamma}$ 126.5°. One of the phenocrysts, composed of 4 individuals proved to be well suited for a detailed study. Repeated measurement gave the following mean Köhler-angles:

¹⁾ SEMQ-ARL-probe, thin section coated with Au, 15 KV, 20 μ A sample current (integrated), \approx 2 sec, beam \varnothing minimal.

Köhler-angles			
twin-axis	$\alpha \alpha'$	$\beta \beta'$	$\gamma \gamma'$
albite-law	169.5°	164°	19°
Karlsbad-law	138.5°	43.5°	169°
complex albite-Karlsbad-law	43°	139°	165°

From the original data, projected on a WULFF net with diametre 40 cm, the following average Euler-angles were read:

95.5° 69.5° 84°

Independant measurements and constructions by a second operator (A. GLAUSER) on the same twin-group gave the values:

95° 70° 84.5°

Interpreted on the high-temperature curves of BURRI-PARKER-WENK (1967), both sets of data indicate an An-content 14 to 17.

The same twin group was tested on the electron microprobe by serial point analyses in the ranges measured optically. The resulting mean values (in bracket one single higher An value, Ab values by difference) are:

Individual	Or	Ab	An
A	5	79	16 (24)
B	7	81	12
C	6	82	12
D	6	76	18
\bar{x}_1	6	79.5	14.5

A neighbouring phenocryst, composed also of 4 individuals, gave a similar average:

\bar{x}_2	6	78.5	15.5
accepted			
mean value	6	79	15

The data for Alftavik oligoclase agree very well with those obtained by I. S. E. CARMICHAEL (1960) for the plagioclase of pitchstone from Bondelfur (F. 8) and by R. SCHEDLER (1971) for samples from Lambadalur, all from the East coast of Iceland, and they fit also those of an oligoclase annealed at high pressure (H. R. WENK, 1969).

	Or	Ab	An	Φ	Θ	Ψ	$2V_\gamma$
Alftavik	6	79	15	95.5°	69.5°	84°	126°
Bondelfur	5.0	78.7	16.3	95.3°	68.7°	84.6°	116°
Lambadalur	7.0	78.8	14.2	91.6°	68.4°	83.6°	—
annealed	4.1	82.4	13.5	95.5°	69.2°	86.8°	128°

Our check shows that there is no need for a revision of the optic curves in the range of acid volcanic oligoclase.

DISCUSSION

In Fig. 1 the projections of feldspar phenocrysts (solid circles), microlites (dots), residual glasses (open circles) and rocks (squares) of vitrophyres from eastern and central Iceland are shown in the ternary feldspar diagram. It gives good information on the variation of the feldspar composition, but ad-




Fig. 1. Ternary feldspar diagram (mol %) showing feldspar phenocrysts (solid circles), microlites (dots), residual glasses (open circles) and rocks (squares) of vitrophyres from eastern and central Iceland. A = Alftavik. Published data from CARMICHAEL (1963, with tie-lines between feldspar and glass), SCHEDLER 1971, WENK et al. 1972. Curve with arrows connects zones observed in one and the same phenocryst from Domadalshraun. Lower curve shows general variation of feldspars studied by CARMICHAEL. Solvus for 1000°C after TUTTLE and BOWEN (1958), for 650°C after SECK (1971).

mittedly, the chemical analysis of residual glasses as well as the calculation of the normative proportion Or Ab An of glasses and vitrophyres are problematic, as the amount of alumina present could be partitioned in different way on K, Na and Ca.

The Alftavik oligoclases are situated within the range of feldspars studied by Carmichael and the residual Alftavik glass projects within the cluster of residual glasses in his diagram, not far from the alkali-feldspar minimum.

Unlike the acid two-feldspar lavas of Domadal in central Iceland, described in a previous paper (WENK et al. 1972), the sodic vitrophyric dyke rock studied here contains only one, slightly zoned feldspar, similar to Icelandic pitchstones discussed by CARMICHAEL (1963). Phenocrysts and microlites have oligoclase composition and the residual glass approaches the alkali-feldspar minimum.

Acknowledgement: We are obliged to A. GLAUSER for the check of optical measurements, to H.-H. KLEIN who drew Fig. 1 and to H. HÄNNI for the preparation of the thin-sections.

REFERENCES

- BURRI, C., PARKER, R. L., WENK, E. and WENK, H. R. (1967): Die optische Orientierung der Plagioklase. Birkhäuser, Basel.
- CARMICHAEL, I. S. E. (1960): The feldspar phenocrysts of some Tertiary acid glasses. *Miner. Mag.* 32, 587–608.
- (1963): The crystallization of feldspar in volcanic acid liquids. *Quart. J. geol. Soc. Lond.* 119, 95–131.
- SCHEDLER, R. (1971): Optical, chemical and structural determinations of volcanic, acid Icelandic plagioclase feldspars. *Schweiz. mineral. petrogr. Mitt.* 51, 463–509.
- THORODDSEN, TH. (1906): Island. Peterm. Mitt. Erg.heft, Nr. 153.
- WENK, E., SCHWANDER, H. and WETZEL, R. (1972): On a zoned ternary feldspar from Domadalshraun, Iceland. *SMPM* 52, 373–380.
- WENK, H. R. (1969): Annealing of oligoclase at high pressure. *Amer. Miner.* 54, 95–100.

Manuscript received June 30, 1975.