

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen = Bulletin suisse de minéralogie et pétrographie
Band: 58 (1978)
Heft: 1-2

Artikel: Tremolite- and diopside-bearing serpentine assemblages in the CaO-MgO-SiO₂-H₂O multisystem
Autor: Oterdoom, W.H.
DOI: <https://doi.org/10.5169/seals-45194>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Tremolite- and Diopside-bearing Serpentine Assemblages in the CaO-MgO-SiO₂-H₂O Multisystem

By *W. H. Oterdoom*, Zürich *)

Abstract

In low grade metamorphic ultramafics, up to the upper pumpellyitefacies, the assemblage chrysotile + tremolite has been described (ROST, 1949; PETERS, 1963; DIETRICH and PETERS, 1971). Also the assemblage chrysotile + diopside has been observed at the same and higher grades up to the reaction of chrysotile to antigorite + brucite where at the same time pumpellyite disappears (DIETRICH and PETERS, 1971). Tremolite with serpentine occurs at low grade while the assemblage tremolite + forsterite is confined to higher metamorphic grades. From these observations the stable P-T topology can be deduced in the CaO-MgO-SiO₂-H₂O multisystem among the phases brucite, chrysotile, antigorite, talc, forsterite, diopside, tremolite and H₂O.

INTRODUCTION

This work is an extension of the serpentine-multisystem (EVANS *et al.*, 1976), in that CaO appears as new component besides MgO-SiO₂-H₂O. The phases diopside and tremolite contain this component. We consider a system with 8 phases and 4 components with a variance of minus two. The unique P-T net will be constructed to satisfy some field observations, by combining two partial nets, each with a variance of minus one. For this reason the thermodynamic data from EVANS *et al.* (1976) and TROMMSDORFF and EVANS (1977a) plus some field observations will be used.

CHEMOGRAPHY AND TOPOLOGY OF THE PARTICIPATING PHASES

The reasons that antigorite (A) and chrysotile (C, abbreviations see table 1) are not considered as polymorphic minerals, are only explained here briefly, for an exhaustive discussion see WICKS and WHITTAKER (1975). Because of

*) Institut für Kristallographie und Petrographie, ETH-Zentrum, CH-8092 Zürich.

the sinelike double half-wave structure with variable curvature, antigorite has a variable chemical composition (KUNZE, 1961). The stronger the curvature of the wave, the less MgO and H₂O has antigorite in comparison to the ideal serpentine composition $\text{Mg}_3\text{Si}_2\text{O}_5(\text{OH})_4$. Theoretically the composition of antigorite can vary between those of chrysotile and talc and does not change the *relative* position of the phases in the chemographic projection (fig. 1). KUNZE




Fig. 1. Chemography of brucite B, chrysotile C, antigorite A, talc T, forsterite F, diopside Di and tremolite Tr in the system CaO-MgO-SiO₂-H₂O.

(1958, p. 316) made the connexion between the formula of the antigorite varieties and their associated layer wave-lengths (superperiods). The antigorite formula is $m \text{Mg}_{3(1-1/m)}\text{Si}_2\text{O}_5(\text{OH})_{1+3(1-2/m)}$. From X-ray studies a superperiod series of $m = 17$ tetrahedra per unit cell is established with $a_0 = 43.3 \text{ \AA}$. The antigorite formula consequently becomes $\text{Mg}_{48}\text{Si}_{34}\text{O}_{85}(\text{OH})_{62}$. The hypothesis of KUNZE (1961, p. 317) that the value of m in the superperiod series of antigorite is a function of metamorphic grade and chemical composition has not yet been investigated.

Until now no systematic derivation exists for a fully closed net of a 4 component system with 8 phases. The P-T net is constructed by means of combining two partial nets, each with a degree of freedom of minus one. We assume, that the pure phases chrysotile and forsterite can not occur together under normal geological conditions (EVANS *et al.*, 1976; HEMLEY *et al.*, 1977b; DUNGAN, 1977). With this assumption it is possible to define two partial nets with forsterite or chrysotile absent. The six phases brucite, chrysotile, antigorite, talc, diopside, tremolite and the three projected components CaO-MgO-SiO₂ from H₂O form the first partial net (a pseudoternary multisystem involving only solid-solid reactions). The chemography is a triangle with all of the phases along the edges, compare with figure 9, number 49 in DAY (1972). Diopside and tremolite lie on the same site of the 4 phase colinearity along MgO-SiO₂ in projection from H₂O, implying that the invariant points [Di] and

[Tr] coincide at a different level of stability: stable to metastable (DAY, rule 4, 1972). There are 8 possible univariant reactions and 6 possible invariant points. In the absent phase notation of DAY (1972) there are 5 possible nets with maximal closure: the triangular nets ((C) (Di)), ((Tr) (B)) and ((Tr) (A)), the quadrilateral nets ((Tr) (C)) and ((Di) (A)). Only ((Tr) (A)) satisfies the assemblages observed in the field (fig. 2a). The corresponding inverse net to that shown in fig. 2a is presumed not to be correct because it cannot be constructed such that all of the appropriate reactions may be oriented with respect to the P-T axis. The nets have been constructed such, that two thermodynamic conditions are satisfied: 1) with increasing temperature at constant pressure the high-entropy side of the reaction is stabilised, 2) with increasing pressure at constant temperature the low-volume side of the reaction is stabilised. Entropy- and volume-data are summarized in table 1.




Fig. 2a and 2b. The unique P-T nets with maximum closure satisfying field data. Fig. 2a for the six phases A, B, C, T, Di, Tr and fig. 2b for A, B, F, T, Di and Tr. Mineral abbreviations, see table 1.


In the second partial system chrysotile is absent and forsterite (and H_2O) are present. The fundamental chemography is preserved (fig. 9, number 49, DAY, 1972). Also in this partial system there are 8 possible univariant reactions with 6 possible invariant points (of which, as mentioned above, [Di] and [Tr] coincide at a different level), therefore [F] and [W] coincide at the same level of stability. Of the 5 possible nets only ((Tr) (A)) satisfies the geological observations (fig. 2b). The corresponding inverse net to fig. 2b is also not possible here for the same reasons as above.

Both partial nets have three univariant reactions ($Tr = Di + T$, $A = T + B$ and $Di + A = Tr + B$) and one invariant point, [C, F, W], in common. From this

framework the whole net can be built up, remembering that in a non-degenerate system with the variance of minus two, three invariant points lie on a single univariant reaction curve (KORZHINSKII, 1959, p. 128, table 2). The stable part of the net that results in this manner has been drawn in fig. 3. The corresponding inverse net to fig. 3 is not considered possible for geological reasons (EVANS *et al.*, 1976, p. 83).

FIELD- AND PETROGRAPHIC-DATA

Chrysotile + tremolite, antigorite + diopside and forsterite + tremolite are the relevant natural serpentine assemblages, formed by the addition of CaO to the system $\text{MgO-SiO}_2\text{-H}_2\text{O}$. Chrysotile + tremolite are found at the lowest metamorphic grades. The assemblages antigorite + diopside and forsterite + tremolite have formed during the progressive metamorphism and are related through reaction (BCT). This reaction (BCT) takes place between the forsterite-forming reaction from antigorite + brucite and the antigorite breakdown reaction to forsterite + talc (EVANS and TROMMSDORFF, 1970; TROMMSDORFF and EVANS, 1972). Antigorite + diopside have formed from chrysotile + tremolite and both assemblages are often described (ROST, 1949; DIETRICH and PETERS, 1971). For the assemblage chrysotile + tremolite we must assume a stability field, as this pair occurs in several serpentinites: Totalp (PETERS, 1963), Oberhalbstein (DIETRICH, 1969; DIETRICH and PETERS, 1971), Wurlitz in the Münchberger Gneissmass (ROST, 1949). In order that a stability field occurs in our multi-system for chrysotile + tremolite, the solid-solid reaction $\text{C} + \text{Tr} = \text{A} + \text{Di}$ (BTFW) has to be placed between the formation reaction of antigorite ($\text{C} + \text{T}$



(1) Totalp, (2) Tiefencastel, (3) Line P. Curver-Cunter, (4) Line P. Arlos-Rona (5) Marmorera, (6) Line Forcellina-Pass-Julier Pass, (7) Engadine Line (Ober-Engadin)

Fig. 4. Critical assemblages in ultramafics in the profile Totalp (Aroser Schuppenzone)–Oberhalbstein. Broken streaks mark local occurrence, the dotted line the occurrence of vein-antigorite. Data for Totalp were compiled after PETERS (1963), for Oberhalbstein after DIETRICH (1969) and DIETRICH and PETERS (1971). Section Totalp–Tiefencastel not on scale.

= A) and the breakdown reaction of chrysotile ($C = B + A$), see also the sequence of assemblages in the Oberhalbstein (fig. 4).

The P-T diagram for CaO-bearing serpentinites must contain the following relevant, divariant fields with progressive metamorphism: $C + Tr + T$, $A + C + Tr$, $A + C + Di$, $A + B + Di$, $A + F + Di$, $A + F + Tr$ and $F + T + Tr$. The assemblage brucite + tremolite has not been described although it would represent the upper pressure limit of antigorite + diopside.

Both tremolite and diopside occur in weakly metamorphosed (prehnite-pumpellyite facies) areas, in the Totalp and in the Oberhalbstein. However, in the Totalp, PETERS (1963) observed tremolite, but never diopside, as new epitaxial growth on primary diallage or as aggregates of amphibole fibres as pseudomorphs after pyroxenes. However, diopside appears in the Totalp in fissures in clinopyroxene serpentinites. Thus the rock-forming assemblage chrysotile + diopside does not occur in the Totalp, but appears first in the higher grade Oberhalbstein. The observed sequence of assemblages in progressive metamorphism is presented in fig. 4.

THE PHASE-DIAGRAM

The P, T and f_{H_2O} phase relations between the 8 phases brucite, chrysotile, antigorite, talc, forsterite, diopside, tremolite and H_2O can be calculated quantitatively in the system $CaO-MgO-SiO_2-H_2O$ from equilibrium data from only four linearly independent reactions, which together contain all 8 phases (SKIPPEN, 1971). Three of the four selected reactions have been determined experimentally and can describe the system $MgO-SiO_2-H_2O$ quantitatively, together with additional information from field data (EVANS *et al.*, 1976). The calculated location of the dehydration reactions, except for the breakdown of antigorite ($A = F + T + H_2O$, see Appendix) agrees to within $\pm 10^\circ C$ with the results of HEMLEY *et al.* (1977a, b). As the fourth linear independent reaction, $A + Di = F + Tr + H_2O$ was selected. TROMMSDORFF and EVANS (1977a) list the equilibrium data for this reaction. With this choice of data we have an internally consistent set, that can also be compared with the set for the ophicarbonate system of TROMMSDORFF and EVANS (1977a, b). For the equilibrium constants for all the univariant reactions, see table 2.

The reaction $C + Tr = A + Di$ (BTFW) takes place according to this data, at 2 kbars and $192^\circ C$. So that (BTFW) lies just above the reaction $C + T = A$ (BFWDiTr). Although absolute temperature calibration in low grade ultramafic rocks is very questionable, it is clear that field data from the Oberhalbstein, see fig. 4, suggest a slightly larger temperature interval than the calculated $7^\circ C$ at 2 kbars, as the first appearance of antigorite + diopside takes place for the first time in the southern Oberhalbstein, about 6 km south of the first

Table 2. *Univariant equilibria in CaO bearing serpentinites*

Absent	Stoichiometry ²⁾	Equilibrium constant ¹⁾			Linear dependency	
		A	B	C		
(C, B, Di, Tr)	A = 18F + 4T + 27H ₂ O	-210330	433.89	5.0019	1	-
(A, B, Di, Tr)	5C = 6F + T + 9H ₂ O	-76500	158.31	1.7259	-	1
(A, T, Di, Tr)	B + C = 2F + 3H ₂ O	-23319	51.57	0.5478	-	1
(C, B, T)	8Di + A = 18F + 4Tr + 27H ₂ O	-213995	445.04	4.7864	-	1
(B, F, W, Di, Tr)	15C + T = A	-19170	41.04	0.1758	-1	3
(B, T, F, W)	15C + Tr = A + 2Di	-18254	38.25	0.2297	-0.75	3
(T, F, W, Di, Tr)	17C = A + 3B	-25713	44.64	0.2583	-1	4
(B, T, Di, Tr)	20C = A + 6F + 9H ₂ O	-95670	199.35	1.9017	-1	4
(C, T, Di, Tr)	A + 20B = 34F + 51H ₂ O	-370710	832.05	9.0543	1	-4
(A, B, T)	5C + 2Di = 6F + Tr + 9H ₂ O	-77416	161.10	1.6720	-0.25	1
(C, A, B, F, W)	2Di + T = Tr	-916	2.79	-0.0539	-0.25	-
(C, F, W, Di, Tr)	2A = 45B + 17T	-59805	-28.08	0.8859	2	9
(A, F, W, Di, Tr)	2C = 3B + T	-6543	3.60	0.0825	-	1
(A, C, Di, Tr)	5B + T = 4F + 6H ₂ O	-40095	99.54	1.0131	-	-1
(C, T, F, W)	34Di + 2A = 17Tr + 45B	-75381	19.31	-0.0300	-2.25	9
(A, T, F, W)	2Di + 2C = Tr + 3B	-7459	6.39	0.0286	-0.25	1
(C, A, T)	Tr + 5B = 2Di + 4F + 6H ₂ O	-39179	96.75	1.0670	0.25	-1

¹⁾ $\ln K = \frac{A}{T} + B + \frac{C(P-1)}{T}$, where T is in °K and P is in bars.

²⁾ See Table 1 for abbreviations and mineral formulae. Fugacity data from BURNHAM et al. (1969).

appearance of antigorite (not vein-antigorite) in the univariant assemblage antigorite + chrysotile + talc and also about 6 km north of the first appearance of antigorite + brucite. This suggests that the reaction $C + Tr = A + Di$ (BTFW) lies somewhere half way between the reactions $C + T = A$ and $C = A + B$. Here one should bear in mind, that the solid-solution effect can be very large on the solid-solid reaction (BTFW), because ΔS of the reaction is small ($\Delta T = \frac{RT \ln K}{\Delta S_r}$).

The equilibrium of the reaction $A + Di = F + Tr + H_2O$ (BCT) can be calculated for 2 kbars at about $486^\circ C$ ($P_{H_2O} = P_{total}$). The reaction (BCT) lies above 1 kbar about $30^\circ C$ beneath the breakdown reaction of antigorite (BCDiTr). The measured widths of the tremolite + antigorite + forsterite and tremolite + talc + forsterite zones in the Bergell tonalite aureole (TROMMSDORFF and EVANS, 1972) could point to a little larger temperature interval between reaction (BCT) and reaction (BCDiTr).

With respect to the field observations both CaO-bearing reactions may be better located if the $\Delta G^\circ_{f,298}$ for diopside is decreased or if $\Delta G^\circ_{f,298}$ for tremolite is increased, resulting in new values for the equilibrium constants A and B of both CaO-bearing reactions: $A = -19597$, $B = 38.85$ for reaction (BTFW) and $A = -208621$, $B = 442.64$ for reaction (BCT). The effect of solid-solution on




Fig. 5. Temperature-pressure ($P_{H_2O} = P_{total}$) diagram for stable (continuous lines) and meta-stable (dashed lines) equilibria in the multisystem antigorite, brucite, chrysotile, talc, forsterite, diopside, tremolite and H_2O . The location of $C + Tr = A + Di$ is based on equilibrium constants A and B cited on page 134.

the location of (BCT) and (BCDiTr) is similar, thus the width of the temperature interval will not be changed by solid-solution. TROMMSDORFF and EVANS (1972) determined the partitioning of Mg between olivine with $x_{\text{Mg}}^{\text{Ol}} = 0.9$ and other phases to be $x_{\text{Mg}}^{\text{Tr}} = 0.976$, $x_{\text{Mg}}^{\text{Di}} = 0.969$, $x_{\text{Mg}}^{\text{Tr}} = 0.961$ and $x_{\text{Mg}}^{\text{A}} = 0.952$ for $P_s = 2$ kbars and $T = 525 \pm 50^\circ \text{C}$. Assuming $\gamma_{\text{Mg}} \approx 1$, the temperature shift at 2 kbars will be about -8°C for both reactions.

The locations of the internally consistent reaction boundaries in the $P_{\text{H}_2\text{O}} (= P_{\text{total}})$ - T diagram are shown in fig. 5. Only the position of (BTFW) is inferred from field data. The location of the dehydration reactions are probably good to at least $\pm 20^\circ \text{C}$ (EVANS *et al.*, 1976), whereas the position of the solid-solid reactions must be regarded as very approximate, as is explained by the example of $\text{C} + \text{Tr} = \text{A} + \text{Di}$ (BTFW). This problem is clear from the fact that the assemblage chrysotile + talc appears south of the Engadine Line (Upper Engadine), several kilometers south of the first appearance of antigorite. This same holds for chrysotile + tremolite relative to antigorite + diopside.

CONCLUSIONS

It is evident from field observations and in accordance with our phase diagram, that tremolite is the most important CaO -bearing mineral of low grade ultramafics. At slightly higher grades (from the upper pumpellyite facies) diopside will dominate until the lower amphibolite facies, where diopside plus antigorite will give forsterite + tremolite.

Acknowledgements

The manuscript was improved by the constructive reviews of A. B. Thompson and V. Trommsdorff. I would like to thank Kathy Riklin for making the drawings. The author wishes to acknowledge the support of the Schweizerischer Nationalfonds, No. 2.615-0.76. The analysis of 94-NZ-62-5 was kindly supplied by B. W. Evans.

Appendix

Problems with the antigorite breakdown equilibrium

For the reaction antigorite (Kunze-type) = forsterite + talc + H_2O , $\text{Mg}_{48}\text{Si}_{34}\text{O}_{85}(\text{OH})_{62} = 18\text{Mg}_2\text{SiO}_4 + 4\text{Mg}_3\text{Si}_4\text{O}_{10}(\text{OH})_2 + 27\text{H}_2\text{O}$, we can recalculate $\Delta G_{f,298}^0$ of antigorite (Kunze-type), using HEMLEY's (1977b) experimentally determined equilibrium of antigorite + talc + forsterite for 1 kbar at 514°C . With the following values for antigorite (Kunze-type): $V^0 = 102.89 \text{ cm}^3$ (KUNZE, 1961), $S_{298}^0 = 50.6 \text{ cal/deg} \cdot \text{gfw}$ (recomputed value of KING *et al.*, 1967; OTERDOOM, 1975) $\Delta S_{f,298}^0 = -249.196 \text{ cal/deg}$, c_p -constants with equation $c_p = a + b \cdot T + c \cdot T^{-2}$: $a = 72.793 \text{ cal/deg} \cdot \text{gfw}$, $b = 30.985 \cdot 10^3 \text{ cal/deg}^2 \cdot \text{gfw}$, $c = -16.975 \cdot 10^{-5} \text{ cal} \cdot \text{deg/gfw}$ (recomputed value of KING *et al.*, 1967; OTERDOOM, 1975) and thermodynamic data for talc and forsterite as listed in table 3 of HEMLEY *et al.* (1977b), $\Delta G_{f,298}^0 = -930533 \text{ cal/gfw}$ for antigorite (Kunze-type). With these data the breakdown reaction of antigorite lies just above the upper limit of Johannes experimentally deter-

mined brackets. The calculation scheme followed has been described by HEMLEY et al. (1977b) with their equation 2. The recalculation has been made because $\Delta S_{T, 298}^0$ of antigorite in HEMLEY's table 3, p. 360, is based on two different antigorite compositions (ideal- and Kunze-type) and V^0 was based on unitcell data given by PAGE and COLEMAN (1967, table 3), that yield an unreasonably large molar volume.

The still existing discrepancy of $\pm 20^\circ \text{C}$ between the locations of the antigorite breakdown reaction of HEMLEY et al. (recalculated) and EVANS et al. could be possibly explained by the different chemical compositions and thus the different antigorite varieties of Hemley's and Johannes samples. Johannes antigorite (EVANS et al., 1976, table 4) shows lower SiO_2 , higher MgO and H_2O content than the one of (HEMLEY et al., 1977b, table 1) and plots so more towards chrysotile whereas Hemley's antigorite plots more towards talc.

The recalculated antigorite breakdown based on HEMLEY et al. is consistent with the antigorite breakdown of Johannes, as supported by field evidence: Johannes antigorite shows less solid solution towards talc and his sample comes from an area (Piz Lunghin, Engadine) where antigorite \pm brucite is the stable assemblage without relict olivine (DIETRICH and PETERS, 1971). The first appearance of forsterite in this area is some kilometers to the south in Val Malenco. On the other hand, Hemley's antigorite, 94-NZ-62-5 supplied by R. COLEMAN, originates from an area with the assemblage antigorite + magnesite \pm talc \pm relict olivine (COLEMAN, 1966, p. 70).

Added in press:

A reinvestigation of 94-NZ-62-5 reveals an almost ideal Kunze-type formula. Compare however the USGS analysis (in PAGE and COLEMAN, 1967, tab. 1).

Composition of 94-NZ-62-5, from Griffin Range, New Zealand

	1) wt %	2) wt %	3)	
SiO_2	44.35	44.35	33.79	} 34.00
TiO_2	0.03	trace	—	
Al_2O_3	0.55	0.55	0.49	
			0.21 0.28	
Fe_2O_3	4.15*)	—	—	} 47.94
FeO	—	3.1**)	1.97	
MnO	0.07	0.03	0.02	
MgO	40.10	40.10	45.54	
CaO	0.04	0.03	0.02	
Na_2O	0.03	trace	—	
K_2O	<0.01	0	—	
P_2O_5	0.01	0	—	
Cr_2O_3	0.065	0.06	0.04	
NiO	0.25	0.11	0.07	
ZnO	0.005	trace	—	
$\text{H}_2\text{O}+$	10.01	11.9	60.48	
	99.65	100.23		

1) Gravimetric- and spectrographic analysis of B. Ayranci and M. Weibel (Zurich) corrected for CO_2 in magnesite.

2) "Best" antigorite-composition based on gravimetrical- and spectrographical analysis of the USGS (in PAGE and COLEMAN, 1967) and Ayranci-Weibel and on microprobe analysis of Evans (Seattle).

*) Total iron as Fe_2O_3 .

**) Total iron as FeO .

3) Numbers of ions on the basis of 116 anhydrous oxygens.

References

- BURNHAM, C. W., J. R. HOLLOWAY and N. F. DAVIS (1969): Thermodynamic properties of water to 1000° C and 10,000 bars. *Geol. Soc. Amer., Spec. Paper 132*, 96 p.
- COLEMAN, R. G. (1966): New Zealand serpentinites and associated metasomatic rocks. *New Zealand Geol. Survey Bull. 76*, 102 p.
- DAY, H. D. (1972): Geometrical analysis of phase equilibria in ternary systems of six phases. *Amer. J. Sci. 272*, 711–734.
- DIETRICH, V. (1969): Die Ophiolithe des Oberhalbsteins (Graubünden) und das Ophiolithmaterial der ostschweizerischen Molasseablagerungen, ein petrographischer Vergleich. Dissertation. Europäische Hochschulschriften, Reihe 17, Erdwissenschaften, Nr. 1, 180 S. Verlag Herbert Lang & Cie. AG, Bern.
- DIETRICH, V. und T. J. PETERS (1971): Regionale Verteilung der Mg-Phyllosilikate in den Serpentinitten des Oberhalbsteins. *Schweiz. mineral. petrogr. Mitt. 51*, 329–348.
- DUNGAN, M. A. (1977): Metastability in serpentine-olivine equilibria. *Am. Min. 62*, 1018–1029.
- EVANS, B. W. and V. TROMMSDORFF (1970): Regional metamorphism of ultramafic rocks in the Central Alps: Parageneses in the system CaO-MgO-SiO₂-H₂O. *Schweiz. mineral. petrogr. Mitt. 50*, 481–492.
- EVANS, B. W., W. JOHANNES, W. H. OTERDOOM, V. TROMMSDORFF (1976): Stability of chrysotile and antigorite in the serpentine multisystem. *Schweiz. mineral. petrogr. Mitt. 56*, 79–93.
- HEMLEY, J. J., J. W. MONTOYA, C. L. CHRIST, P. B. HOSTETLER (1977a): Mineral equilibria in the MgO-SiO₂-H₂O system: I, talc-chrysotile-forsterite-brucite stability relations. *Amer. J. Sci. 277*, 322–351.
- HEMLEY, J. J., J. W. MONTOYA, D. R. SHAW and R. W. LUCE (1977b): Mineral equilibria in the MgO-SiO₂-H₂O system: II, talc-antigorite-forsterite-anthophyllite-enstatite stability relations and some geologic implications in the system. *Amer. J. Sci. 277*, 353–383.
- KING, E. G., R. BARANY, W. W. WELLER and L. B. PANKRATZ (1967): Thermodynamic properties of forsterite and serpentine. U.S. Bureau of Mines, Report of Investigations 6962, 19 p.
- KORZHINSKII, D. S. (1957): Physicochemical basis of the analysis of the paragenesis of minerals. (Engl. transl., New York consultants Bureau, 1959.)
- KUNZE, G. (1958): "Die gewellte Struktur des Antigorits II." *Z. Krist. 110*, S. 282–320.
- KUNZE, G. (1961): Antigorit. Strukturtheoretische Grundlagen und ihre praktische Bedeutung für weitere Serpentin-Forschung. *Fortschr. Mineral. 39*, 206–324.
- OTERDOOM, W. H. (1975): Geologisch-petrographische Untersuchungen des Malencoserpentin in Val Scerscen. Diplomarbeit, ETH Zürich.
- PAGE, N. J. and R. G. COLEMAN (1967): Serpentine-mineral analyses and physical properties. U.S. Geol. Survey Prof. Paper, 575-B, B103–B107.
- PETERS, T. J. (1963): Mineralogie und Petrographie des Totalpsersperitins bei Davos. *Schweiz. mineral. petrogr. Mitt. 43*, 529–685.
- ROBIE, R. A. and D. R. WALDBAUM (1968): Thermodynamic properties of minerals and related substances at 298.15 K (25.0° C) and one atmosphere (1.013 bars) pressure and at higher temperatures. U.S. geol. Surv. Bull. 1259, 256 S.
- ROST, F. (1949): Das Serpentin-Gabbro-Vorkommen von Wurlitz und seine Mineralien. *Heidelberger Beiträge zur Mineralogie und Petrographie 1*, 626–688.
- SKIPPEN, G. B. (1971): Experimental data for reactions in siliceous marbles. *J. Geol. 79*, 457–481.

- TROMMSDORFF, V. and B. W. EVANS (1972): Progressive metamorphism of antigorite schists in the Bergell tonalite aureole (Italy). *Amer. J. Sci.* 272, 423-437.
- TROMMSDORFF, V. and B. W. EVANS (1974): Alpine metamorphism of peridotitic rocks. *Schweiz. mineral. petrogr. Mitt.* 54, 333-352.
- (1977a): Antigorite-ophicarbonates: Phase relations in a portion of the system CaO-MgO-SiO₂-H₂O-CO₂. *Contr. Mineral. Petrol.* 60, 39-56.
- (1977b): Antigorite-ophicarbonates: Contactmetamorphism in Val Malenco, Italy. *Contr. Mineral. Petrol.* 62, 301-312.
- WICKS, F. J. and E. J. W. WHITTAKER (1975): A reappraisal of the structures of the serpentine minerals. *Can. Mineral.* 13, 227-243.

Manuscript received May 25, 1978.