

Zeitschrift: Schweizerische mineralogische und petrographische Mitteilungen =
Bulletin suisse de minéralogie et pétrographie

Band: 59 (1979)

Heft: 1-2

Artikel: Geothermic and radiometric investigations

Autor: Rybach, L.

DOI: <https://doi.org/10.5169/seals-46050>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 26.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Geothermic and Radiometric Investigations

Report by *L. Rybach* *)

The present report summarizes the activity of Working Group No. 5 during the period 1971–79. Besides the activities reported below which relate closely to geodynamic problems a number of further sidelines of geothermics and radiometry have been followed up, mainly in view to practical applications: assessment of geothermal potential, underground heat storage, prospecting for uranium mineralisations, radioactive waste disposal. Computer codes as well as experimental techniques developed for these «applied» topics have been subsequently utilised with great benefit in the more «general» projects listed below.

Working Group Members are: Ph. Bodmer, P. Finckh, K.J. Hsü, E. Kissling, M. Parini, L. Rybach, D. Werner (Zurich), T. P. Labhart (Berne), G. Berset **) (Lucerne).

1. SUBSURFACE TEMPERATURES

A reliable data base is necessary for downward extrapolation of near-surface data in order to delineate the temperature field in the earth's crust. Temperature measurements have been performed in all accessible drill-holes, tunnels, shafts etc.: continuous logging as well as bottom-hole measurements (the latter corrected for effects of mud circulation) in drillholes, single-point measurements in the rock face in tunnels and shafts. The results are compiled in a uniform format (KRÜSI et al., 1978).

Computer programs have been developed to account for the effect of surface topography and its evolution on subsurface temperatures (BODMER et al., 1979). The availability of a topographic array, covering whole Switzerland with a distance between mesh points of 250 m, avoids digitizing by hand. In areas like the Swiss Alps where considerable relief exists the effect of topography on subsurface temperatures prevails down to depths ≥ 5 km.

*) Institute of Geophysics, ETH-Hoenggerberg, CH-8093 Zurich (Switzerland)

**) Working Group Chairman 1971–1974

2. HEAT FLOW DETERMINATIONS

Heat flow determinations have been carried out in drillholes, shafts and in lake-bottom sediments. The corrections applied account for topography, uplift/erosion, sedimentation and climatic changes. Special attention was given to the evaluation of suitable sites for heat flow determinations in lake bottom sediments (v. HERZEN et al. 1974, FINCKH 1976, FINCKH & KELTS 1976). The results are discussed in accordance with their geologic-tectonic settings in RYBACH & FINCKH (1979). In the Northern Foreland of the Swiss Alps, especially towards the Upper Rhine Graben, heat flow values up to 100 mW/m² have been observed, whereas the Swiss Alps are characterized by generally lower and less uniform heat flow, reflecting the complex present and past geodynamics of this area (RYBACH et al. 1977, WERNER 1977).

In order to summarize on-going geothermal research activities in Europe and to represent the heat flow field on a continental scale the Monograph «Terrestrial Heat Flow in Europe» has been prepared. Due to the manifold geodynamic implications of heat flow this Monograph appears as «*Inter-Union Commission of Geodynamics Scientific Report No. 58*» (ČERMÁK & RYBACH 1979).

3. HEAT PRODUCTION

Abundance and distribution of radiogenic heat sources greatly influence the temperature field in the continental crust (see e.g. RYBACH 1976 a). Experimental techniques (gamma ray spectrometry) have been developed for heat production determination (RYBACH, 1971). Revised heat production constants for the main radioelements uranium, thorium and potassium are given in Table 1.

Table 1 Revised heat production constants (RYBACH, 1976 b)

Element	A [cal/g.y] (Birch, 1954)	A [cal/g.y] (Rybach, 1976 b)	A [W/kg]
Uranium	0.73	0.718	$9.525 \cdot 10^{-5}$
Thorium	0.20	0.198	$2.561 \cdot 10^{-5}$
Potassium	$276 \cdot 10^{-6}$	$26.2 \cdot 10^{-6}$	$3.477 \cdot 10^{-9}$

Systematic laboratory studies have been carried out in order to assign characteristic heat production values to the main rock types of the Swiss Alps (RYBACH, 1973). Geochemical variation trends (heat production - bulk chemical composition of rocks) are summarized and discussed in RYBACH (1976c).




Fig. 1 The seismic velocity - heat production relationship for different pressures (from RYBACH, 1978).

The zonation pattern of heat production $\underline{A}$ in granitic batholiths (decreasing $\underline{A}$ towards the center of the body) has been studied in detail in the Rotondo granite (Gotthard massif, Central Alps): vertical decrease of $\underline{A}$ with depth has been found; the logarithmic decrement (for exponential decrease) is $H=8$ km (KISSLING et al. 1979).

An empirical relationship between heat production $\underline{A}$ and compressional wave velocity $\underline{v_p}$ has been found (RYBACH, 1973; 1976b): in the range characteristic of crustal rocks (5–8 km/sec) $\underline{A}$ decreases exponentially with increasing $\underline{v_p}$: $A(v_p) = a \cdot \exp(-b \cdot v_p)$; where the numerical factors $\underline{a}$ and $\underline{b}$ depend on in-situ pressure and temperature (Fig. 1).

4. GEOTHERMICS AND GEODYNAMICS

In order to shed light on the manifold interrelations of geothermics and geodynamics basic theoretical investigations have been carried out: dynamic

modeling of heat conduction in a moving system, interactions between heat and mass transfer (WERNER 1975).

Special studies have been devoted to

- a) gravity anomalies in the wake of drifting continents, caused partly by radiogenic heating of the overriding lithosphere (KAHLE & WERNER 1975)
- b) cooling history of the Lepontine area (Central Alps), considering time-dependent uplift/erosion (WERNER et al. 1976; see also Fig. 2)
- c) the geothermal anomaly of the Upper Rhine Graben: the regional part can be attributed to hot rising mantle material below this continental rift structure (WERNER & FUCHS, 1977; WERNER et al., 1978; KAHLE & WERNER, 1979; WERNER & KAHLE, 1979, cf. Fig. 3), whereas local anomalies like the Landau anomaly are interpreted as hydrothermic effects (WERNER & PARINI, in preparation). The anomalous temperature field can explain the observed discrepancy in the gravity anomalies (up to 160 mgal). The velocity and density inversions in the upper crust which are associated with pronounced lateral and vertical temperature gradients are also responsible for the relative vertical motions in the Rhine Graben (MÜLLER & RYBACH, 1974).

5. GEOTHERMAL MAPS

Recent activities in Switzerland focus on the preparation of maps representing the geothermal conditions at depth. This project is sponsored by the Swiss Geophysical Commission and by the Federal Office of Energy. Based on




Fig. 2 Time-dependent uplift/erosion in the Lepontine area (upper part of diagram) and corresponding cooling history (lower part). Calculation by D. WERNER, based on data from WERNER et al. (1976) and apatite fission track data from WAGNER et al. (1977). Orogenic activity was strongest 20 mio years ago; quiescent period during the Pliocene.




Fig. 3 Model of material and heat transport in the uppermost mantle below the Rhine Graben (from WERNER & KAHLE, 1979)

Left: The velocity field $v = (v_x, v_z)$ for the upwelling mantle material under the graben zone which produces a thermal anomaly;

Right: The temperature field after 45 mio years caused by convection and thermal conduction; a calculated boundary between lithosphere and asthenosphere (stippled area) depending on temperature and pressure is also indicated.

References

- BODMER, PH., ENGLAND, P. C., KISSLING, E. & RYBACH, L. (1979): On the correction of subsurface temperature measurements for the effect of topographic relief. Part II: Application to temperature measurements in the Central Alps. *In*: V. Čermák & L. Rybach (1979), p. 78–87.
- ČERMÁK, V. & RYBACH, L. (1979): *Terrestrial Heat Flow in Europe*. Springer Verlag, Heidelberg-Berlin-New York, 328 pp.
- FINCKH, P. G. (1976): *Wärmeflussbestimmungen in Randalpenseen*. Diss. ETH Zurich No. 5787, 105 pp.
- FINCKH, P. G. & KELTS, K. (1976): Geophysical investigations into the nature of pre-Holocene sediments of Lake Zurich. *Eclogae geol. Helv.* 69, 139–148.
- V. HERZEN, R. P., FINCKH, P. & HSÜ, K. J. (1974): Heat flow measurements in Swiss lakes, *J. Geophys.* 40, 141–172.
- KAHLE, H.-G. & WERNER, D. (1975): Gravity and temperature anomalies in the wake of drifting continents. *Tectonophysics* 29, 487–504.
- KAHLE, H.-G. & WERNER, D. (1979): A geophysical study of the Rhinegraben rift system. Part II: Gravity anomalies and geothermal implications. *Geophys. J. R. astr. Soc.* (in press).
- KRÜSI, H. R., BODMER, PH. & RYBACH, L. (1978): *Sammlung, Kompilation and Interpretation geothermischer Daten in der Schweiz*. Swiss Federal Office of Energy, Bern, 79 pp.
- KISSLING, E., LABHART, T. P. & RYBACH, L. (1979): Radiometrische Untersuchungen am Rotondogranit. *Schweiz. Min. Petr. Mittg.* (in press).
- LABHART, T. P. (1976): Die Radioaktivität alpiner Gesteine. *Schweizer Strahler* 4, 3–14.
- LABHART, T. P. & RYBACH, L. (1974): Granite und Uranvererzungen in den Schweizer Alpen. *Geol. Rundschau* 63, 135–147.
- MÜLLER, ST. & RYBACH, L. (1974): Crustal dynamics in the central part of the Rhinegraben. *In*: J. H. Illies & K. Fuchs (eds.): *Approaches to Taphrogenesis*, Schweizerbart, Stuttgart, p. 379–388.
- RYBACH, L. (1971): Radiometric Techniques. *In*: R. E. Wainerdi & E. A. Uken (eds.): *Modern Methods of Geochemical Analysis*, Plenum Publishing Co., New York, p. 271–318.
- RYBACH, L. (1973): *Wärmeproduktionsbestimmungen an Gesteinen der Schweizer Alpen*. Beitr. Geol. Schweiz, Geotechn. Ser. 51, 43 pp.
- RYBACH, L. (1976a): Die Gesteinsradioaktivität und ihr Einfluss auf das Temperaturfeld in der kontinentalen Kruste. *J. Geophys.* 42, 93–101.
- RYBACH, L. (1976b): Radioactive heat production and its relation to other petrophysical parameters. *Pure Appl. Geophys.* 114, 309–317.
- RYBACH, L. (1976c): Radioactive heat production, a physical property determined by the chemistry of rocks. *In*: R. G. J. Strens (ed.): *«The Physics and Chemistry of Minerals and Rocks»*, Wiley & Sons, London, p. 309–318.
- RYBACH, L. (1978): The relationship between seismic velocity and radioactive heat production in crustal rocks: An exponential law. *In*: L. Rybach & L. Stegena (eds.): *«Geothermics and Geothermal Energy»*, Special Issue *Pure Appl. Geophys.* 117, p. 75–82.
- RYBACH, L. & JAFFÉ, F. (1976): Geothermal potential in Switzerland. *In*: *Proc. 2nd United Nations Symposium on the Development and Use of Geothermal Resources*, San Francisco, Vol. I, p. 241–244.
- RYBACH, L., WERNER, D., MUELLER, ST. & BERSSET, G. (1977): Heat flow, heat production and crustal dynamics in the Central Alps, Switzerland. *In*: A. M. Jessop (ed.): *Heat Flow and Geodynamics*, Special Issue *Tectonophysics* 41, p. 113–126.
- RYBACH, L. & FINCKH, P. G. (1979): Heat flow data in Switzerland. *In*: V. Čermák & L. Rybach (1979), p. 278–282.
- WAGNER, G. A., REIMER, G. M. & JÄGER, E. (1977): Cooling ages derived by apatite fission track, mica Rb-Sr and K-Ar dating: The uplift and cooling history of the Central Alps. *Mem. Ist. Geol. Mineral. Univ. di Padova* Vol. XXX, 28 pp.

- WERNER, D. (1975): Probleme der Geothermik am Beispiel des Rheingrabens. Diss. Univ. Karlsruhe, 171 pp.
- WERNER, D. (1977): Erdwärme und Lithosphäre im Raum Südwestdeutschland/Schweiz. *Naturwiss.* 64, 626–631.
- WERNER, D. & DOEBL, F. (1974): Eine geothermische Karte des Rheingrabenuntergrundes. *In*: J. H. Illies & K. Fuchs (eds.): *Approaches to Taphrogenesis*, Schweizerbart, Stuttgart, 182–191.
- WERNER, D., KÖPPEL, V., HÄNNY, R. & RYBACH, L. (1976): Cooling models for the Lepontine area (Central Swiss Alps). *Schweiz. Min. Petr. Mittg.* 56, 661–667.
- WERNER, D. & FUCHS, K. (1977): Exploration and interpretation of the geothermal anomaly in the Rhine Graben. *EC Seminar on Geothermal Energy, Brussels, Vol. II*, p. 189–208.
- WERNER, D., FUCHS, K., PARINI, M., SATTEL, G., SCHROTH, G. & DOEBL, F. (1978): Exploration and interpretation of the strongly anomalous temperature field underneath the Rhinegraben. *Publ. European Communities, Brussels*.
- WERNER, D. & KAHLE, H.-G. (1979): A geophysical study of the Rhinegraben rift system. Part I: Kinematics and geothermics. *Geophys. J. R. astr. Soc.* (in press).
- WERNER, D., KAHLE, H.-G., ANSORGE, J. & MUELLER, ST. (1979): The Rhinegraben Rift System: A new model from geophysical results (in press).

Manuscript received July 23, 1979