

**Zeitschrift:** Schweizerische mineralogische und petrographische Mitteilungen =  
Bulletin suisse de minéralogie et pétrographie

**Band:** 77 (1997)

**Heft:** 1

**Artikel:** Blueschist relics in the Mesozoic cover series of the Menderes Massif  
and correlations with Samos Island, Cyclades

**Autor:** Candan, Osman / Dora, O. Özcan / Oberhänsli, Roland

**DOI:** <https://doi.org/10.5169/seals-58471>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 30.06.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

SHORT NOTE

## Blueschist relics in the Mesozoic cover series of the Menderes Massif and correlations with Samos Island, Cyclades

by Osman Candan<sup>1</sup>, O. Özcan Dora<sup>1</sup>, Roland Oberhänsli<sup>2</sup>, Frederike Oelsner<sup>2</sup> and Stefan Dürr<sup>3</sup>

### Abstract

In the western part of the Menderes Massif, Selçuk-Dilek Peninsula, relics of Tertiary HP metamorphism under epidote-blueschist facies conditions were recognized in the Mesozoic cover series. This subduction-related high-pressure metamorphism, with temperature of less than 470 °C and minimum pressure of about 10 kbar, was subsequently followed by a Barrovian-type overprint under greenschist facies conditions during Late Eocene / Early Oligocene. Based on the lithostratigraphy and the two-fold Tertiary metamorphism, the Mesozoic cover series in this area can be correlated with the Vourlites unit on Samos Island, belonging to the Cycladic Crystalline Complex.

*Keywords:* blueschist, eclogite, Menderes Massif, Samos Island, Cycladic Crystalline Complex.

### Introduction

The Menderes Massif (MM), composed of a large crystalline nucleus within the Alpidic orogenic belt, is exposed in western Anatolia. This crystalline complex is bordered to the north and west by the Izmir-Ankara Zone and to the south by the Lycian nappes of the Taurides. The MM is considered to be linked to the Cycladic Crystalline Complex to the west in the Central Aegean Sea, as a part of Median Crystalline Belt (DÜRR et al., 1978). The latter is one of the world's typical blueschist belts.

Some relic eclogites related to high-pressure metamorphism in the MM were recorded in the Precambrian "core series" (CANDAN et al., 1994; OBERHÄNSLI et al., 1995). In recent years, some evidence for high-pressure metamorphism, most probably not related to the HP event in the "core series", has been detected in the "cover series". The aim of this short note is to document the presence of blueschists as well as eclogites in the Mesozoic cover series of the MM, and to correlate the Selçuk region, in the western part of the MM, with the Island of Samos in the Cycladic island arc.

### Geology and phase relationships in the high-pressure rocks

The relics of Tertiary high-pressure metamorphism in the Mesozoic cover series were recognized in the areas of Selçuk, Kusadase and Dilek Peninsula (Fig. 1). The relevant rocks include

- i) blueschist metabasites and metapelites, and
- ii) high-pressure blocks in a metamorphosed olistostrome.

*Blueschist relics* were only detected in the Dilek Peninsula, where marble-schist successions with minor metabasite intercalations display predominantly greenschist facies assemblages. In three localities, well-preserved blue amphiboles were found in metabasites within thin epidote-crossite-rich layers and lense-shaped bodies, which are suggesting a protolith of mafic volcanic tuffs (Fig. 1). Furthermore, carbonate-rich metapelites, most probably mixed with volcanic material, contain blue amphibole porphyroblasts up to 2 cm in length. Some of the crossites rimmed by barroisitic green amphiboles are altered into actinolite and chlorite. Phyllitic metapelites, inter-

<sup>1</sup> Dokuz Eylül Üniversitesi Müh. Fak. Jeoloji Müh. Bölümü Bornova-Izmir, Turkey.

<sup>2</sup> Institut für Geowissenschaften Universität Potsdam – PF 601553, D-14415 Potsdam, Germany.

<sup>3</sup> Institut für Geowissenschaften Johannes Gutenberg Universität Mainz, D-55099 Mainz, Germany.




Fig. 1 Correlation of the main Mesozoic units between Samos Island, Cyclades, and the western part of Menderes Massif. Relic outcrops of Tertiary blueschist metamorphism are shown. The geological map of Samos is modified after PAPANIKOLAOU (1979) and OKRUSCH et al. (1985). The localities of metabauxites and rudists are after ÖNAY (1949) and ÖZER (1993).

Tab. 1 Representative microprobe analyses of Na and Na-Ca amphiboles from blueschist facies metabasites. H<sub>2</sub>O recalculated from stoichiometry.

	Na amphibole			Na-Ca amphibole		
	3-35	3-38	3-41	3-43	3-19	3-30
SiO <sub>2</sub>	56.24	56.65	56.36	56.21	50.89	50.35
TiO <sub>2</sub>	0.03	0.00	0.04	0.04	0.10	0.12
Al <sub>2</sub> O <sub>3</sub>	7.33	6.14	7.87	7.35	4.83	5.09
Cr <sub>2</sub> O <sub>3</sub>	0.03	0.02	0.02	0.00	0.05	0.00
Fe <sub>2</sub> O <sub>3</sub>	8.57	9.14	6.84	7.37	8.65	7.42
FeO	8.83	9.14	9.38	9.97	18.28	10.47
MnO	0.14	0.18	0.11	0.07	0.31	0.35
MgO	9.18	9.00	9.05	8.84	11.69	11.71
CaO	0.65	0.38	0.75	0.62	7.81	8.13
Na <sub>2</sub> O	6.91	6.85	6.73	6.99	3.38	3.13
K <sub>2</sub> O	0.00	0.00	0.03	0.00	0.17	0.19
H <sub>2</sub> O*	2.13	2.12	2.13	2.12	2.06	2.04
Total	100.04	99.61	99.30	99.57	100.20	98.98
Numbers of ions on the basis of 23 O and 2 OH						
Si	7.898	8.006	7.943	7.942	7.394	7.395
Ti	0.003	0.000	0.004	0.004	0.011	0.013
Al	1.213	1.023	1.306	1.224	0.827	0.880
Cr	0.003	0.002	0.002	0.000	0.005	0.000
Fe <sup>3+</sup>	0.906	0.972	0.726	0.783	0.946	0.820
Fe <sup>2+</sup>	1.037	1.080	1.106	1.178	1.249	1.286
Mn	0.016	0.021	0.013	0.008	0.038	0.043
Mg	1.923	1.896	1.902	1.861	2.532	2.563
Ca	0.097	0.057	0.114	0.093	1.215	1.279
Na	1.881	1.876	1.840	1.914	0.952	0.891
K	0.001	0.000	0.005	0.000	0.031	0.035
OH	2.000	2.000	2.000	2.000	2.000	2.000

bedded with marbles, exhibit a greenschist assemblage of chlorite + albite + chloritoid + white mica + quartz. Irrespective of the retrograde minerals, the mineral assemblages of blueschists can be summarized as:

*Metabasites*

crossite + epidote + albite + phengite  
crossite + epidote + albite

*Metapelites*

crossite + calcite + epidote + albite  
crossite + calcite + epidote + phengite

All assemblages also contain additional quartz.

*Blocks with high-pressure mineral assemblages* occurring in the metamorphosed olistostrome unit (Fig. 1) are mainly composed of eclogite, smaragdite-omphacite metagabbro and flaser metagabbro. The high-pressure relics, mostly associated with strongly foliated metaserpentinites,




Fig. 2 Compositional diagrams of (A) alkali amphiboles and (B) Na-Ca amphiboles from blueschist facies metabasites (after LEAKE, 1978).

are well defined around the Selçuk area. Transitional metamorphic stages from gabbro to eclogite are recognized in some blocks. Most of the eclogites display retrograde alteration to garnet amphibolites. Fresh eclogites with zoisite/clinozoisite-omphacite-rich layers, forming lenses and schlieren up to 10 meters, are characterized by the following assemblages:

*Eclogite*

omphacite + garnet + zoisite + epidote  
omphacite + garnet + zoisite + rutile

*Zoisite-omphacite-rich rocks*

omphacite + epidote + zoisite  
omphacite + zoisite

**Mineral geochemistry and P-T estimations**

Representative sodic and sodic-calcic amphibole analyses of the blueschist metabasites are given in table 1. Textural evidence reveal that sodic amphiboles which were generated by the high-pressure metamorphism are partly or completely altered into green amphibole and chlorite. Al<sub>2</sub>O<sub>3</sub> and Na<sub>2</sub>O contents in blue amphiboles range

between 6.1–7.9 and 6.3–7.0 wt% respectively and CaO is lower than 1.2 wt%. Sodic amphiboles plotted on the LEAKE (1978) diagram concentrate in the crossite field (Fig. 2a). Green amphiboles, which are partly scattered into the winchite field, generally display a barroisitic composition in the Na–Ca amphibole diagram (Fig. 2b).

Distribution of the blueschist relics coincides roughly with the diaspore-corundum isograd in the Dilek Peninsula, as well as on the Island of Samos. In the eastern part of Samos, metamorphic temperature and pressure for the HP metamorphic phase was estimated at 440 °C and 12 kbar (OKRUSCH et al., 1985). If such a temperature value is accepted for the Dilek Peninsula, the crossitic sodic amphiboles, which have a compositional scatter between groups 5 and 3 in the epidote-blueschist facies system after EVANS (1990), lead to assume minimum pressures between 7.5 and 9.5 kbar, respectively. At 9.5 kb pressure, a maximum temperature of 470 °C is given experimentally by HAAS (1972) for the diaspore-corundum isograd. Therefore, as a consequence of the absence of corundum in the metabauxites coexisting with blueschist assemblages on the Dilek Peninsula, the maximum temperature must have been less than 470 °C during high-pressure metamorphism.

### Discussion

The Attica-Cycladic crystalline complex in the central Aegean Sea is characterized by spectacular outcrops of eclogites and blueschists. This complex underwent high-pressure metamorphism at a grade corresponding to the transition between eclogite and epidote-blueschist facies metamorphism in Late Eocene / Early Oligocene time, and was subsequently overprinted by a medium-pressure Barrovian-type metamorphism under greenschist to amphibolite facies conditions at the Oligocene/Miocene boundary (ALTHERR et al., 1979).

The Island of Samos is a part of the Cycladic Blueschist Belt, and is situated two kilometers NW of the Dilek Peninsula. According to THEODOROPOULOS (1979) the crystalline series of Samos form a normal stratigraphic succession, whereas PAPANIKOLAOU (1979) and OKRUSCH et al. (1985) regard it as a nappe pile. The Ampelos nappe, which contains widespread blueschist relics, glaucophane metagabbro, glaucophanite and omphacitic rocks (MYPOSKOS and PERDIKATIS, 1984; OKRUSCH et al., 1985; CHEN, 1992), is made up of three tectonic units: Vourliutes, Ampelos and Aghios. The Vourliutes unit which is ex-

posed in the eastern part of Samos, consists of interbedded marbles and metapelites with minor ultramafic and mafic intercalations (OKRUSCH et al., 1985). This isoclinally folded succession (CHEN, 1992) with widespread blueschist relics is typified by metaclastic quartzitic phyllites with the striking assemblage of kyanite + chloritoid. One sample containing kyanite + chloritoid ± Mg-charpholite is reported (OKRUSCH et al., 1985). Metabauxite deposits in the marbles with coexisting diaspore + chloritoid, but without corundum, were documented by PAPANIKOLAOU (1979). The same authors also recognized traces of rudist fossils, suggesting an Upper Cretaceous age (ÖZER, 1993) for the metabauxite-bearing marbles (Fig. 1).

The age of the HP metamorphism, which affected the core series (CANDAN et al., 1994; OBERHÄNSLI et al., 1995) and most probably was accompanied by a HT-granulite facies event, is unknown. The blueschist and eclogite relics in the Mesozoic cover series are restricted to the Selçuk-Dilek Peninsula, in the western part of the MM (DORA et al., 1995). The rock succession of this region can be divided into two main units, from bottom to top, (i) interbedded marbles and schists, and (ii) a metamorphosed olistostrome unit. Although a gradational contact between these two units is suggested by ERDOGAN and GÜNGÖR (1992), the true character of this contact is controversial. An isoclinally folded marble-schist succession is dominated in its lowest part by metaclastites, metaconglomerates with quartzite pebbles, quartzites and quartz phyllites, which are characterized by assemblages of chloritoid + kyanite. The metaclastic units include blueschist bearing metabasites, and change gradually upward to thick metacarbonate series interbedded with minor carbonate-rich blueschists and chloritoid-chlorite phyllites. The marbles with well-preserved rudist fossils indicating an Upper Cretaceous age (ÖZER, 1993), are characterized by the widespread occurrence of metabauxites, diasporites and corundum-bearing rocks. In terms of lithostratigraphic succession and the presence of blueschist relics the marble-schist succession in particular can be correlated with the Vourliutes unit of Samos.

Although the metamorphosed olistostrome unit comprise spectacular high-pressure blocks, no high-pressure relics have been recognized in the matrix which is dominated by chlorite-albite schists. It is still open to discussion whether these high-pressure blocks are exotic or whether they suffered high-pressure metamorphism together with the matrix under transitional conditions from epidote-blueschist to eclogite facies as on

Syros and Syfnos islands (MATTHEWS and SCHLIESTEDT, 1984; RIDLEY and DIXON, 1984). As an alternative interpretation, we suggest that the matrix has been completely overprinted by the later greenschist assemblages, while the mafic blocks mostly preserved their HP assemblages. This olistostromal unit, which does not crop out on the Island of Samos, could be correlated with the metamorphosed olistostrome unit in Syros (RIDLEY and DIXON, 1984; OKRUSCH and BRÖCKER, 1990).

The blueschist relics from the Selçuk area reveal that a subduction-related high-pressure/low-temperature metamorphism affected the western end of the MM under epidote-blueschist to eclogite (?) facies conditions during Tertiary time. The HP/LT assemblages were subsequently overprinted by a Barrovian-type medium-pressure metamorphism under greenschist to amphibolite facies conditions during Late Eocene / Early Oligocene boundary (SATIR and FRIEDRICHSEN, 1986; DORA et al., 1995; HETZEL and REISCHMANN, 1996). The newly presented evidence of a HP event during the Tertiary metamorphic evolution provides strong support for the correlation of the Menderes Massif and the Cyclades crystalline complexes, as previously proposed by DÜRR et al. (1978) and OKRUSCH et al. (1985).

#### Acknowledgements

This work was supported by Volkswagen Stiftung/Germany and DFG-Project OB80/12. We thank R. Altherr for help with microprobe work, M. Çetinkaplan and E. Koralay for assistance during field work and O. Kaya for critically reading the manuscript.

#### References

- ALTHERR, R., SCHLIESTEDT, M., OKRUSCH, M., SEIDEL, E., KREUZER, H., HARRE, W., LENZ, H., WENDT, I. and WAGNER, G.A. (1979): Geochronology of high-pressure rocks on Sifnos (Cyclades, Greece). *Contr. Mineral. Petrol.*, 70, 245–255.
- CANDAN, O., DORA, O.Ö., DÜRR, ST. and OBERHÄNSLI, R. (1994): Erster Nachweis von Granulit- und Eklogit-Relikten im Menderes-Massiv / Türkei. *Göttinger. Abh. Geol. Paläont.*, 5. Symposium TSK, 217–220.
- CHEN, G. (1992): Evolution of the high- and medium-pressure metamorphic rocks on the island of Samos, Greece. Ph. D. Theses, Julius-Maximilians University, Würzburg, 170 pp.
- DORA, O.Ö., CANDAN, O., DÜRR, ST. and OBERHÄNSLI, R. (1995): New evidence concerning the geotectonic evolution of the Menderes Massif. *Internat. Earth Science Colloq. on the Aegean Region 1995, Izmir – Turkey. Proceedings.*
- DÜRR, ST., ALTHERR, R., KELLER, J., OKRUSCH, M. and SEIDEL, E. (1978): The Median Aegean crystalline belt: Stratigraphy, structure, metamorphism and magmatism. In: CLOSS, H., ROEDER., D.H. and SCHMIDT, K. (eds): *Alps, Apennines and Hellenides*, 455–478, Schweizerbart, Stuttgart.
- ERDOGAN, B. and GÜNGÖR, T. (1992): Menderes Masi-fi'nin kuzy kanadinin stratigrafisi ve tektonik evrimi. *TPJD Bülteni*, c. 2/1, S., 1–20.
- EVANS, B.W. (1990): Phase relations of epidote-blueschist. *Lithos*, special. T.I.E.C. issue., 25, 3–23.
- HAAS, H. (1972): Equilibria in the system  $Al_2O_3$ - $SiO_2$ - $H_2O$  involving the stability limits of diaspore and pyrophyllite, and thermodynamic data on these minerals. *Amer. Mineral.*, 57, 1375–1385.
- HETZEL, R. and REISCHMANN, T. (1996): Intrusion age of Pan-African augen gneisses in the southern Menderes Massif and the age of cooling after Alpine ductile extensional deformation. *Geol. Mag.*, 133, 5, 565–572.
- LEAKE, B.E. (1978): Nomenclature of amphiboles. *Amer. Mineral.*, 63, 1023–1053.
- MATTHEWS, A. and SCHLIESTEDT, M. (1984): Evolution of the blueschist and greenschist facies rocks of Sifnos, Cyclades, Greece. *Contrib. Mineral. Petrol.*, 88, 150–163.
- MPOSKOS, E. and PERDIKATIS, V. (1984): Petrology of glaucophane metagabbros and related rocks from Samos, Aegean Island (Greece). *N. Jb. Min. Abh.* 149, 1, 43–63.
- OBERHÄNSLI, R., CANDAN, O., MEZGER, K., DORA, O.Ö. and DÜRR, ST. (1995): Eclogites and granulites in the Menderes Massif, Western Turkey. *EUG. 8, Strasbourg, Terra Abstr.*, 18.
- OKRUSCH, M. and BRÖCKER, M. (1990): Eclogites associated with high grade blue schists in the Cyclades archipelago, Greece: A review. *Eur. J. Mineral.*, 2, 451–478.
- OKRUSCH, M., RICHTER, P. and KATSIKATIS, G. (1985): High pressure rocks of Samos, Greece. The geological evolution of Eastern Mediterranean. *Geol. Soc. London. Spec. Pub. 17*, (ed.) Dixon and Robertson., 529–536.
- ÖNAY, T. (1949): Über die Smirgelgesteine SW-Anatoliens. *Schweiz. Mineral. Petrogr. Mitt.*, 29, 359–484.
- ÖZER, S. (1993): Upper Cretaceous rudists from the Menderes Massif. 6th. Cong. of the Geol. Soc. of Greece, *Bull. of the Geol. Soc. of Greece*, XXVIII, 3, 55–73.
- PAPANIKOLAOU, D.J. (1979): Unités tectoniques et phases de déformation dans l'île de Samos, Mer Egée, Greece. *Bull. Soc. Géol. France*, XIX, 6, 745–752.
- RIDLEY, J. and DIXON, J.E. (1984): Reaction pathways during the progress deformation of a blueschist metabasite: the role of chemical disequilibrium and restricted range equilibrium. *J. metamorphic Geol.* 2, 115–128.
- SATIR, M. and FRIEDRICHSEN, H. (1986): The origin and evolution of the Menderes Massif, W-Turkey: Rubidium/Strontium and oxygen isotope study. *Geol. Rundsch.*, 75/3, 703–714.
- THEODOROPOULOS, D. (1979): Geological map of Greece 1:50.000, island of Samos, Athens, I.G.M.E.

Manuscript received December 30, 1996; manuscript accepted February 2, 1997.