

Nd, Sr, Pb isotope study of the Western Carpathians : implications for Palaeozoic evolution

Autor(en): **Poller, Ulrike / Todt, Wolfgang / Kohút, Milan**

Objektyp: **Article**

Zeitschrift: **Schweizerische mineralogische und petrographische Mitteilungen
= Bulletin suisse de minéralogie et pétrographie**

Band (Jahr): **81 (2001)**

Heft 2

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-61685>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Nd, Sr, Pb isotope study of the Western Carpathians: implications for Palaeozoic evolution

by Ulrike Poller¹, Wolfgang Todt¹, Milan Kohút² and Marian Janák³

Abstract

Nd, Sr and Pb–Pb whole rock isotope data from the Variscan basement of the Western Carpathians are presented in order to define the eastern continuation of the Variscides in Central Europe. Isotope analyses of migmatites, orthogneisses and granitoids from the Tatra Mountains (Upper unit) and two neighbouring areas (Velká Fatra and Lower Tatra Mountains) have been examined. The granites, migmatites and orthogneisses show mainly crustal characteristics with $\epsilon\text{Nd}_{(330)}$ values ranging from -2 to -9 and $\epsilon\text{Sr}_{(330)}$ values between 10 and 387. These results are in good agreement with the Pb–Pb whole rock data and document a crustal origin for the investigated rocks with only minor influence of an enriched mafic component such as recycled oceanic crust. REE and geochemical data are consistent with a collisional environment for the emplacement of the granitoids and the precursor of the orthogneisses. The isotope data confirm that the Western Carpathians own a quite similar isotopic composition than other Variscan regions in Europe.

On the basis of the data presented, in combination with our previous U–Pb zircon dating, we propose the following geological evolution for the Tatra Mountains. In Early Devonian time (~ 406 Ma) subduction-related magmatism led to the formation of the earliest granitoids, the precursors of the present-day orthogneisses. In Late Devonian and Carboniferous time (360–350 Ma), younger granitoids intruded, most probably as a result of collisional thickening. During this event the earlier granitoids were recrystallised and deformed, resulting in the formation of the orthogneiss fabric. The youngest granitic magmatism in the Carboniferous (315 Ma) is related to the Late Variscan orogenic collapse.

Keywords: Variscides, Western Carpathians, radiogenic isotopes, granitoids.

Introduction

The Variscides constitute the main orogenic belt of Europe resulting from the convergence between Gondwana and Laurasia during Palaeozoic time (e.g. MATTE, 1986; ZIEGLER, 1986; NEUGEBAUER, 1988; FRANKE 1989). The Carpathians are the key area for the study of the eastern and south-eastern continuation of the Variscan orogen in Central and Eastern Europe. Similar to the Alps, Variscan complexes form the pre-Mesozoic basement of the Carpathian belt (e.g. KRIST et al., 1992; NEUBAUER and VON RAUMER, 1993). However, the precise chronology and plate tectonic setting of the Western Carpathians are not well constrained.

This study is concerned with the Variscan rock complexes exposed in the Tatra Mountains, the Velká Fatra and the Lower Tatra Mountains (Fig. 1a), which are located in the northernmost sector of the Western Carpathians. They are representative of the so-called core mountains within the Tatric unit, a major tectonic unit of the Western Carpathians (ANDRUSOV, 1968; KRIST et al., 1992).

Nd, Sr and Pb–Pb isotopic data on orthogneisses, migmatites and granites are presented, most of them from the Western and the High Tatra Mountains (Fig. 1b). It is the aim of this paper to present these isotopic results, together with petrological and geochronological data (JANÁK et al., 1996, 1999, POLLER et al., 2000; POLLER and TODT, 2001), and to discuss them relevant to the

¹ Max-Planck-Institut für Chemie, Abt. Geochemie, Postfach 3060, D-55020 Mainz, Germany.
<poller@mpch-mainz.mpg.de>

² Dionyz Stúr Institute of Geology, Geological Survey of Slovakia, 817 04 Bratislava, Slovak Republic.

³ Geological Institute, Slovak Academy of Science, Dúbravská 9, 842 26 Bratislava, Slovak Republic.




Fig. 1 Simplified geological maps (after KOHÚT and JANÁK, 1994). (A) Western Carpathians with Fatra, Low Tatra, and Tatra Mountains. (B) Detailed view to the Western and High Tatra Mountains.

geodynamic evolution of the pre-Alpine basement of the Western Carpathians and its relation to other parts of the European Variscides.

Geological Setting

The crystalline basement of the Tatra Mountains (Fig. 1b) is composed of pre-Mesozoic metamorphic and granitoid rocks, overlain by Mesozoic and Cenozoic sedimentary cover sequences and nappes. Metamorphic rocks are abundant in the western part (Western Tatra Mountains), whereas in the eastern part (High Tatra Mountains) they occur only as xenoliths in granitoids. Within the basement, two superimposed tectonic units (lower and upper) have been distinguished, differing in lithology and metamorphic grade (JANÁK, 1994).

The *Lower Unit* is exposed in the Western Tatra in a tectonic window and is composed of micaschists. Kyanite-, staurolite-, fibrolitic sillimanite-, and garnet-bearing metapelites alternate with quartz-rich metapsammites, interpreted to represent former flysch sediments (KAHAN, 1969; GURK, 1999). Two metamorphic zones have been distinguished – the kyanite-staurolite and kyanite-sillimanite (fibrolite) zone (JANÁK, 1994).

The *Upper Unit* is composed of metapelitic paragneiss, amphibolite, and orthogneiss. At its base, kyanite indicates a pressure-dominated type of metamorphism. Metapelites contain kyanite and show migmatization with formation of granitic leucosomes. Orthogneisses are porphyritic and coarse grained, sometimes also leucocratic and medium to fine-grained.

The banded amphibolites include layers of mafic (amphibolitic) and felsic (tonalitic to trondhjemitic) composition and enclose lenses of retrograded eclogites containing garnet and clinopyroxene (JANÁK et al., 1996). Higher levels of the Upper Unit belonging to the sillimanite zone, exhibit widespread migmatization and are intruded by granitoids. Sillimanite, garnet, K-feldspar, and cordierite are diagnostic minerals in these high-grade metapelites. The granitoids comprise leucogranites to biotite tonalites as well as amphibole diorites, and form a sheet-like pluton (KOHÚT and JANÁK, 1994).

The crystalline basement of the Tatra Mountains was affected by Variscan and Alpine deformation under distinct P–T conditions and kinematics (KAHAN, 1969; FRITZ et al., 1992). The earliest Variscan deformation (D1) is related to south-east thrusting of the Upper Unit onto the Lower Unit. Subsequent deformation (D2) was the result of Variscan extension and mostly oblit-

erated the compressional, thrust-related deformation fabrics. Alpine deformation (D3) under brittle conditions is manifested by north-west shear, resulting from Late Cretaceous compression. The last major deformation (D4), related to Tertiary extension, produced updoming and normal faulting in a north-south to NW–SE direction.

Rb–Sr whole-rock isochron dating (BURCHART, 1968) indicates Early Palaeozoic tectonometamorphic activities between 380–420 Ma. Granitoid magmatism, according to Rb–Sr isochrons, took place at 290–310 Ma (BURCHART, 1968) or at 340–350 Ma (GAWEDA, 1995).

Recent U–Pb single zircon dating resulted in a detailed chronology for the different magmatic events in the Tatra Mountains (POLLER et al., 2000; POLLER and TÖDT, 2001). The oldest granites are the present-day porphyric orthogneisses. These intruded 406 Ma ago, followed some 50 Ma later by the main metamorphic overprint and intrusion of granites of the Western Tatra 355 Ma ago. For the High Tatra Mountains this Late Devonian age is reflected only by concordant zircons in the migmatites. Moreover, migmatites show a second concordant zircon age at 335 Ma that is interpreted as the final migmatization. The final granite magmatism at 315 Ma has been documented by zircons in the High Tatra Mountains only. Cooling ages from granitoids and migmatites range between 330 and 300 Ma according to ^{40}Ar – ^{39}Ar white mica dating (MALUSKI et al., 1993; JANÁK and ONSTOTT, 1993; JANÁK 1994) and reflect the late Variscan exhumation.

There is no geochronological indication for a high-temperature Alpine rejuvenation of the basement. Apatite fission track data record the final uplift during the Late Tertiary, at 15–10 Ma (KOVÁČ et al., 1994).

In this study, rocks from the Upper Unit – orthogneisses, migmatites, and granitoids – of the Tatra Mountains have been investigated using geochemical and radiogenic isotope methods, in order to constrain the nature of the source rocks involved and the geotectonic evolution of the Tatricum. Details of petrography and analytical procedures are given in the Appendices 1 and 2, respectively.

Results

ORTHOgneisses

The *porphyric, coarse-grained orthogneisses* show monzogranitic-granodioritic and peraluminous characteristics documented e.g. by the alumina




Fig. 2 Discrimination diagram using the alumina saturation index (ASI) versus SiO_2 for the characterisation of S- and I-type granitoids (after CHAPPELL and WHITE, 1974).

saturation index (ASI), which is above 1.12 for all investigated orthogneisses (Fig. 2). Their mineral (see Appendix 1) and chemical compositions reflect an S-type precursor. Considering the high ASI values, $\text{FeO}_{\text{tot}} / (\text{FeO}_{\text{tot}} + \text{MgO})$ ratios around 0.7, and the crustal-type $^{87}\text{Sr}/^{86}\text{Sr}$ values above 0.709 they are characterised as ‘‘MPG’’ granites after BARBARIN (1999). MPG stands for muscovite-bearing peraluminous granites and describes the precursor of the Tatra orthogneisses with respect to their petrographical and geochemical features. A continental collision is the typical geotectonic environment of these granitoids (BAR-




Fig. 3 Chondrite normalised rare earth element diagram for Western and High Tatra granitoids. Normalising data are from EVENSEN (1978).

BARIN, 1999). The main process responsible for the evolution of such a crustal-derived magma should be fractional crystallisation. The differentiation status of the Western Tatra orthogneisses is assumed to be rather evolved, visible in the high ASI values (BARBARIN, 1996) and the strong peraluminous character (CHAPPELL et al., 1998). Nevertheless, typical characteristics of highly fractionated granitoids such as increasing P, HREE, and Y contents are not always found. Analyses of the REE of the porphyric orthogneisses resulted in typical crustal chondrite-normalised patterns (Fig. 3), showing a negative Eu anomaly with Eu/Eu^* ratios between 0.4 and 0.9.

The *leucocratic orthogneisses* are syenogranitic, with clearly higher SiO_2 contents (75–79 wt%) than the porphyric type. The ASI in sample UP 1005 is 1.3, which indicates the protolith as an S-type granitoid, whereas sample UP 1012 has an ASI of 1.1, indicating a slightly less peraluminous character of the precursor.

Trace elements Nb, Y, Rb, and Zr as well as $\text{FeO}_{\text{tot}} / (\text{FeO}_{\text{tot}} + \text{MgO})$ ratios between 0.72 and 0.77 classify the precursors of these rocks as former MPGs that most probably intruded during the collision of two continents or continental fragments. The source material of the leucocratic orthogneisses again was crustal, but compared with the porphyric gneisses, the leucocratic gneisses contain less mafic minerals such as biotite. The leucocratic orthogneiss UP 1012 has an Eu/Eu^* ratio of 0.7 defining only a slight negative Eu anomaly, and La_n/Lu_n ratio is around 4.

The geochemical differences (lower content of Al_2O_3 , TiO_2 , Fe_2O_3 , K_2O , higher SiO_2 in the leucocratic orthogneisses) between the two orthogneiss groups of the Western Tatra Mountains are also reflected by the radiogenic isotope characteristics.

The $^{206}\text{Pb}/^{204}\text{Pb}$ ratios of the orthogneiss whole rock samples range from 17.99 to 18.89 and the $^{207}\text{Pb}/^{204}\text{Pb}$ ratios from 15.60 to 15.74 (Tab. 4), both of which are significantly higher than mantle or MORB values. The leucocratic sample UP 1012 shows lower Pb–Pb ratios than the porphyric orthogneisses. The $^{206}\text{Pb}/^{204}\text{Pb}$ vs. $^{207}\text{Pb}/^{204}\text{Pb}$ data of the Western Carpathian rocks (Fig. 4b) fit well within the continental crust fields of DOE and ZARTMAN (1979). Nevertheless, it is impossible to distinguish whether in addition to the continental crust component there is also a contribution from other sources such as recycled oceanic crust or an ocean island arc. Thus, there is evidently participation of continental-derived sediments and/or oceanic-pelagic sediments during the genesis of these rocks, as documented by the crust-dominated Pb–Pb ratios.




Fig. 4 $^{207}\text{Pb}/^{204}\text{Pb}$ vs. $^{206}\text{Pb}/^{204}\text{Pb}$ diagrams illustrate (a) evolution, (b) origin and (c) isotopic relationship of the Western Carpathian rocks towards typical Variscan areas. In figure 4a the Tatra rocks as well as reference fields for upper crust, enriched mantle II and MORB (after DOE and ZARTMAN, 1979) are shown with their present day position as well as recalculated to 350 Ma with a μ of 9. This illustration reflects the evolution of the Pb–Pb for the last 330 Ma. No essential change between the relative position of the Western Carpathians and the reference fields is documented. Therefore, in figures 4b and 4c the present day Pb–Pb values are used. In figure 4b the Tatra rocks are compared to several sources (after DOE and ZARTMAN, 1979; ZINDLER and HART, 1986). Figure 4c shows the good correspondence between the Tatra and several other Variscan orogenic areas (data from REISCHMANN and ANTHES, 1996; VIDAL and POSTAIRE, 1985). The reference line for upper crust refers to DOE and ZARTMAN (1979).

Sm–Nd whole-rock studies resulted in $\epsilon\text{Nd}_{(330)}$ values between -2.8 and -9.4 (Fig. 5). The $^{87}\text{Sr}/^{86}\text{Sr}_{(330)}$ ratios show wide variation between 0.7080 (UP 1025) and 0.7322 (UP 1012). Similarly, the $\epsilon\text{Sr}_{(330)}$ values range from 45 to 132 in the porphyritic types, whereas for the leucocratic orthogneisses the values scatter between 205 and 387. Also, the Nd model ages of the two orthogneisses are different. Whereas the leucocratic orthogneisses have $T_{(DM)}$ ages between 1650 and 1780 Ma, interpreted by a two-stage evolution, the porphyritic orthogneisses have mean crustal residence ages (one step model) ranging from 1220–1350 Ma.

GRANITOIDS AND MIGMATITES

Petrographically, all Tatra granites represent basically crustal-anatectic rocks with primary magmatic muscovite. Biotite is the dominant Fe–Mg mineral, whereas hornblende occurs only rarely in dioritic xenoliths. Prevalence of plagioclase (An_{20-45}) over K-feldspar is a typical attribute of the investigated granitoid rocks. The mafic and

Tab. 1 Coordinates of the sample locations from the Tatra Mountains.

| samples | northern latitude | eastern longitude |
|---------|-------------------|-------------------|
| UP 1002 | 49° 09'04" | 19° 42'35" |
| UP 1005 | 49° 08'40" | 19° 43'00" |
| UP 1012 | 49° 09'28" | 19° 46'45" |
| UP 1014 | 49° 08'24" | 19° 48'00" |
| UP 1015 | 49° 09'56" | 20° 09'25" |
| UP 1016 | 49° 09'56" | 20° 09'25" |
| UP 1017 | 49° 09'47" | 20° 09'17" |
| UP 1018 | 49° 09'47" | 20° 09'17" |
| UP 1023 | 49° 10'16" | 19° 45'02" |
| UP 1025 | 49° 10'32" | 19° 44'35" |
| UP 1026 | 48° 59'49" | 19° 16'31" |
| UP 1027 | 48° 59'51" | 19° 15'47" |
| UP 1028 | 48° 59'55" | 19° 15'39" |
| UP 1030 | 48° 59'28" | 19° 08'40" |
| UP 1031 | 49° 00'18" | 19° 09'58" |
| UP 1036 | 49° 10'45" | 19° 51'23" |
| UP 1038 | 49° 04'52" | 19° 10'26" |
| UP 1039 | 49° 11'04" | 19° 46'40" |
| UP 1040 | 49° 11'50" | 19° 46'45" |
| UP 1044 | 49° 09'16" | 19° 58'50" |
| UP 1049 | 49° 09'47" | 20° 09'17" |
| UP 1050 | 49° 09'47" | 20° 09'17" |
| UP 1051 | 48° 56'30" | 19° 34'20" |
| UP 1052 | 49° 08'00" | 20° 07'22" |
| UP 1053 | 49° 08'00" | 20° 07'22" |
| UP 1055 | 49° 16'04" | 19° 59'05" |
| UP 1057 | 49° 10'48" | 20° 03'08" |
| UP 1063 | 49° 10'30" | 19° 50'27" |
| UP 1088 | 48° 56'26" | 19° 34'35" |
| UP 1090 | 48° 54'00" | 19° 24'55" |

Tab. 2a Major element (in wt%), trace element, and REE (in ppm) data for the investigated rocks of Western and High Tatra Mountains.

| sample | leucocratic orthogneisses | | | porphyric orthogneisses | | | High Tatra diorites | | | High Tatra granitoids and migmatites | | | | | | | | | | |
|--------------------------------|---------------------------|------|------|-------------------------|------|-------|---------------------|------|------|--------------------------------------|------|------|------|------|------|------|------|------|------|------|
| | 1005 | 1012 | 1002 | 1014 | 1025 | 1015 | 1016 | 1023 | 1036 | 1040 | 1017 | 1018 | 1044 | 1049 | 1050 | 1052 | 1053 | 1055 | 1057 | 1063 |
| SiO ₂ | 78.6 | 75.4 | 70.4 | 63.6 | 67.8 | 58.9 | 57.6 | 73.8 | 70.9 | 73.1 | 71.1 | 69.4 | 69.9 | 69.9 | 67.6 | 73.4 | 68.7 | 66.6 | 69.8 | 67.8 |
| TiO ₂ | 0.1 | n.d. | 0.3 | 0.8 | 0.6 | 0.8 | 0.9 | 0.1 | 0.4 | 0.2 | 0.3 | 0.4 | 0.3 | 0.4 | 0.5 | 0.6 | 0.4 | 0.6 | 0.3 | 0.5 |
| Al ₂ O ₃ | 13.2 | 14.5 | 16.3 | 17.2 | 16.9 | 17.1 | 16.8 | 14.7 | 14.9 | 15.1 | 15.0 | 15.6 | 15.6 | 15.0 | 16.2 | 12.3 | 16.2 | 16.7 | 15.9 | 16.7 |
| Fe ₂ O ₃ | 0.9 | 0.5 | 2.6 | 5.5 | 3.4 | 6.8 | 7.4 | 0.7 | 2.8 | 1.8 | 2.2 | 2.9 | 2.2 | 2.4 | 3.1 | 4.4 | 3.0 | 3.4 | 2.3 | 3.9 |
| MgO | 0.4 | 0.1 | 1.1 | 2.4 | 1.5 | 3.8 | 4.4 | 0.2 | 1.0 | 0.5 | 0.8 | 0.9 | 0.7 | 0.8 | 1.0 | 0.8 | 1.0 | 1.5 | 0.8 | 1.3 |
| CaO | 0.5 | 0.6 | 1.7 | 2.9 | 2.4 | 5.4 | 5.7 | 3.2 | 2.9 | 1.1 | 2.0 | 4.2 | 3.7 | 3.4 | 4.2 | 2.3 | 4.7 | 4.5 | 4.6 | 2.0 |
| Na ₂ O | 4.8 | 4.6 | 6.0 | 4.0 | 4.5 | 3.1 | 3.2 | 0.5 | 5.0 | 4.4 | 3.6 | 2.7 | 1.4 | 2.0 | 2.6 | 3.2 | 2.1 | 2.4 | 2.2 | 4.8 |
| K ₂ O | 1.3 | 4.3 | 1.4 | 2.6 | 2.2 | 2.4 | 2.4 | 4.9 | 1.0 | 3.1 | 4.3 | 2.6 | 4.3 | 4.5 | 3.3 | 1.9 | 2.1 | 2.4 | 2.2 | 1.8 |
| P ₂ O ₅ | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.4 | 0.4 | 0.1 | 0.4 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | <0.1 |
| LOI | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 0.8 | 1.5 | 0.9 | 0.88 | 0.6 | 1.2 | 1.6 | 1.5 | 1.2 |
| Ba | 91 | 70 | 300 | 792 | 549 | 1317 | 1272 | 2481 | 307 | 826 | 1245 | 739 | 1183 | 1374 | 1129 | 286 | 545 | 578 | 879 | 324 |
| Ga | 12 | 15 | 15 | 22 | 20 | 21 | 22 | 12 | 16 | 18 | 17 | 20 | 19 | 18 | 20 | 18 | 22 | 21 | 21 | 18 |
| Nb | 6 | 5 | 5 | 16 | 10 | 10 | 11 | 3 | 7 | 7 | 8 | 12 | 8 | 9 | 10 | 9 | 10 | 9 | 7 | 8 |
| Ni | 5 | 2 | 9 | 28 | 14 | 10 | 12 | 6 | 5 | 2 | 5 | 3 | 3 | 5 | 5 | 3 | 3 | 7 | 4 | 19 |
| Pb | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 40 | n.d. | n.d. | n.d. | 35 | 20 | 47 | 41 | 17 | 26 | 18 | 24 | 28 |
| Rb | 37 | 136 | 57 | 86 | 75 | 97 | 90 | 75 | 28 | 71 | 102 | 80 | 99 | 110 | 95 | 66 | 76 | 79 | 53 | 66 |
| Sr | 109 | 52 | 300 | 429 | 515 | 1439 | 1394 | 312 | 357 | 310 | 553 | 556 | 352 | 554 | 620 | 189 | 412 | 568 | 562 | 281 |
| Th | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 2 | n.d. | n.d. | n.d. | 18 | 5 | 28 | 27 | 3 | 13 | 9 | 7 | 7 |
| U | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 1 | 1 | 2 | 1 | 4 | 2 | 1 | 1 | n.d. |
| Y | <3 | 6 | 6 | 13 | 10 | 24 | 26 | 6 | 27 | 12 | 8 | 10 | 12 | 11 | 11 | 29 | 11 | 11 | 11 | 22 |
| Zn | 5 | 5 | 35 | 131 | 88 | 95 | 95 | 32 | 50 | 39 | 40 | 54 | 29 | 46 | 55 | 61 | 68 | 69 | 45 | 75 |
| Zr | 18 | 23 | 28 | 187 | 145 | 125 | 130 | 39 | 97 | 97 | 134 | 178 | 124 | 135 | 181 | 123 | 174 | 181 | 150 | 112 |
| La | n.d. | 2.5 | 3.9 | 41.1 | 25.3 | 56.6 | n.d. | 13.4 | 45.3 | 16.3 | n.d. | 25.0 | 22.1 | 34.6 | 45.5 | 8.0 | 38.1 | 31.0 | 28.1 | 21.0 |
| Ce | n.d. | 5.0 | 7.8 | 79.4 | 50.9 | 107.7 | n.d. | 25.5 | 92.4 | 32.3 | n.d. | 48.0 | 31.0 | 68.3 | 90.5 | 18.1 | 75.5 | 59.9 | 53.1 | 43.2 |
| Pr | n.d. | 0.5 | 1.0 | 9.7 | 6.4 | 13.6 | n.d. | 3.1 | 11.9 | 3.8 | n.d. | n.d. | n.d. | 8.7 | 11.6 | n.d. | 9.8 | 7.5 | 6.6 | n.d. |
| Nd | n.d. | 1.9 | 3.9 | 36.1 | 24.1 | 51.0 | n.d. | 11.6 | 43.6 | 14.1 | n.d. | 23.4 | 20.1 | 32.0 | 14.0 | 10.3 | 36.7 | 28.6 | 24.1 | 17.1 |
| Sm | n.d. | 0.6 | 1.1 | 6.5 | 4.8 | 8.3 | n.d. | 2.4 | 9.8 | 2.6 | n.d. | 6.0 | 5.2 | 6.6 | 8.2 | 5.0 | 7.1 | 5.2 | 4.6 | n.d. |
| Eu | n.d. | 0.1 | 0.9 | 1.5 | 1.3 | 1.9 | n.d. | 1.9 | 1.9 | 0.6 | n.d. | n.d. | n.d. | 1.1 | 1.3 | n.d. | 1.3 | 1.0 | 1.0 | n.d. |
| Gd | n.d. | 0.5 | 1.2 | 4.9 | 3.6 | 6.0 | n.d. | 1.8 | 8.5 | 2.0 | n.d. | n.d. | n.d. | 4.9 | 5.8 | n.d. | 5.2 | 3.5 | 3.1 | n.d. |
| Tb | n.d. | 0.1 | 0.2 | 0.6 | 0.5 | 0.8 | n.d. | 0.2 | 1.2 | 0.3 | n.d. | n.d. | n.d. | 0.6 | 0.7 | n.d. | 0.6 | 0.4 | 0.4 | n.d. |
| Dy | n.d. | 0.6 | 1.1 | 2.9 | 2.3 | 4.4 | n.d. | 1.1 | 6.4 | 1.6 | n.d. | n.d. | n.d. | 2.8 | 3.1 | n.d. | 2.8 | 2.2 | 2.1 | n.d. |
| Ho | n.d. | 0.1 | 0.2 | 0.5 | 0.4 | 0.8 | n.d. | 0.2 | 1.1 | 0.3 | n.d. | n.d. | n.d. | 0.4 | 0.5 | n.d. | 0.4 | 0.4 | 0.4 | n.d. |
| Er | n.d. | 0.3 | 0.5 | 1.1 | 0.9 | 2.3 | n.d. | 0.4 | 2.3 | 0.8 | n.d. | n.d. | n.d. | 0.8 | 0.9 | n.d. | 0.9 | 1.0 | 1.0 | n.d. |
| Tm | n.d. | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | n.d. | 0.1 | 0.3 | 0.1 | n.d. | n.d. | n.d. | 0.1 | 0.1 | n.d. | 0.1 | 0.1 | 0.1 | n.d. |
| Yb | n.d. | 0.5 | 0.3 | 0.7 | 0.7 | 2.1 | n.d. | 0.4 | 1.3 | 0.7 | n.d. | n.d. | n.d. | 0.6 | 0.5 | n.d. | 0.6 | 0.9 | 0.8 | n.d. |
| Lu | n.d. | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | n.d. | 0.1 | 0.2 | 0.1 | n.d. | n.d. | n.d. | 0.1 | 0.6 | n.d. | 0.1 | 0.1 | 0.1 | n.d. |

n.d. = not determined

LOI = loss on ignition

Tab. 2b Major element (in wt%), trace element, and REE (in ppm) data for the investigated rocks of Fatra and Low Tatra.

| sample | Fatra granitoids | | | | | | Low Tatra granitoids | | |
|--------------------------------|------------------|------|------|------|------|------|----------------------|------|------|
| | 1026 | 1027 | 1028 | 1030 | 1031 | 1038 | 1051 | 1088 | 1090 |
| SiO ₂ | 73.8 | 65.6 | 67.1 | 70.4 | 69.0 | 76.0 | 70.4 | 71.3 | 68.4 |
| TiO ₂ | 0.1 | 0.8 | 0.7 | 0.4 | 0.6 | 0.1 | 0.4 | 0.4 | 0.6 |
| Al ₂ O ₃ | 15.0 | 16.8 | 16.3 | 15.6 | 16.0 | 14.1 | 14.4 | 14.6 | 17.1 |
| Fe ₂ O ₃ | 1.3 | 4.1 | 3.7 | 2.6 | 3.3 | 0.4 | 3.4 | 2.9 | 4.0 |
| MgO | 0.4 | 1.6 | 1.3 | 1.0 | 1.2 | 0.1 | 1.5 | 1.2 | 1.9 |
| CaO | 0.8 | 2.6 | 3.1 | 1.8 | 2.1 | 0.6 | 3.5 | 2.0 | 2.5 |
| Na ₂ O | 4.3 | 4.3 | 4.2 | 4.1 | 4.0 | 4.3 | 2.1 | 3.7 | 4.4 |
| K ₂ O | 3.3 | 2.7 | 2.4 | 2.9 | 3.1 | 3.4 | 2.9 | 3.6 | 2.9 |
| P ₂ O ₅ | 0.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | 0.3 |
| LOI | 0.9 | 1.1 | 0.8 | 1.1 | 0.8 | 0.7 | 0.9 | 0.8 | 1.7 |
| Ba | 479 | 786 | 788 | 864 | 906 | 399 | 362 | 567 | 1498 |
| Ga | 16 | 23 | 20 | 21 | 21 | 17 | 18 | 16 | 21 |
| Nb | 8 | 9 | 9 | 8 | 8 | 5 | 10 | 9 | 11 |
| Ni | 4 | 4 | 2 | 3 | 4 | 1 | 15 | 13 | 11 |
| Pb | 27 | 17 | 18 | 28 | 28 | 17 | 19 | 22 | 15 |
| Rb | 102 | 100 | 70 | 83 | 98 | 104 | 116 | 114 | 83 |
| Sr | 137 | 604 | 570 | 436 | 547 | 151 | 154 | 163 | 548 |
| Th | 6 | 7 | 5 | 8 | 9 | 1 | 11 | 8 | 9 |
| U | 3 | n.d. | n.d. | 2 | n.d. | n.d. | 3 | 2 | 1 |
| Y | 8 | 7 | 7 | 13 | 13 | 5 | 41 | 29 | 9 |
| Zn | 42 | 85 | 76 | 62 | 62 | 11 | 42 | 38 | 69 |
| Zr | 58 | 214 | 192 | 162 | 191 | 50 | 108 | 100 | 178 |
| La | 14.0 | 28.4 | 18.1 | 28.2 | 28.3 | 7.1 | 24.3 | 14.0 | 37.0 |
| Ce | 21.1 | 55.3 | 36 | 59.0 | 59.2 | n.d. | 47.8 | 34.1 | 71.2 |
| Pr | n.d. | 5.2 | n.d. | 3.1 | n.d. | n.d. | 6.0 | 6.2 | 11.0 |
| Nd | 10.0 | 24.0 | 18.3 | 27.0 | 25.1 | 5.0 | 22.7 | 20.4 | 36.3 |
| Sm | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 5.3 | 2.1 | 8.4 |
| Eu | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 0.8 | n.d. | n.d. |
| Gd | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 5.3 | n.d. | n.d. |
| Tb | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 0.9 | n.d. | n.d. |
| Dy | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 6.4 | n.d. | n.d. |
| Ho | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 1.4 | n.d. | n.d. |
| Er | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 0.4 | n.d. | n.d. |
| Tm | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 0.6 | n.d. | n.d. |
| Yb | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 3.6 | n.d. | n.d. |
| Lu | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | 0.5 | n.d. | n.d. |

n.d. = not determined

LOI = loss on ignition

the metamorphic xenoliths occur together, enclosed in rather homogenous granitoid rocks of the High Tatra. These granitoids have Sr contents above 300 ppm, and low ⁸⁷Rb/⁸⁶Sr ratios of 0.2 to 1.0. The FeO_{tot}/(FeO_{tot}+MgO) ratios of all investigated granitoids lie between 0.69 and 0.76.

Similar to the orthogneisses, the granitoids of the Tatra Mountains also have a peraluminous character as reflected in alumina saturation values of 1.05 to 1.3, whereas the diorites have ASI values of 0.92 to 0.97 and are therefore meta-aluminous. The granitoids represent medium- to high-potassium magmatic rocks. According to CHAPPELL and WHITE (1974), the granitoids can

be classified as S-type rocks: only the mafic dioritic xenoliths are clearly I-type derived (Fig. 2).

The chondrite normalised REE patterns (Fig. 3) of the granitoids exhibit only slight negative Eu anomalies and show uniform fractionation trends (La_n/Yb_n = 51–15).

Whole rock Pb–Pb analyses (Table 4) show a spread in ²⁰⁶Pb/²⁰⁴Pb of between 18.34 and 19.08 and in ²⁰⁷Pb/²⁰⁴Pb between 15.66 and 15.74. The granitoids occupy nearly the same position as the orthogneisses in figure 4; thus their crustal character is well constrained by the common Pb system.

Sm–Nd whole rock analyses resulted in εNd₍₃₃₀₎ values of –4 to –1, in accordance with prelimi-

nary data of KOHÚT et al. (1999). In combination with $\epsilon\text{Sr}_{(330)}$ values ranging from 10 to 60 (Tab. 3, Fig. 5), they reflect a strong crustal influence for the granitoids. The Nd model ages for the granitoids range from 1250–1450 Ma (Fig. 6) and compare well with those of the porphyric orthogneisses.

The migmatites (UP 1052 and UP 1063) show nearly the same geochemical composition as the surrounding granites, reflecting a close relationship to the associated granites. Nevertheless, their isotopic composition is different. With $\epsilon\text{Sr}_{(330)}$ values of 114 and 112, the migmatites show a stronger crustal influence than the surrounding granites. In contrast to the granites, the $\epsilon\text{Nd}_{(330)}$ value of the migmatite UP 1052 is positive and its Nd model age is much younger (about 900 Ma), probably reflecting juvenile input. The study of STILLE and SCHALTEGGER (1996) postulated a lithospheric mantle, enriched in LREE and character-

ised by elevated Sm/Nd ratios up to 0.168 for Central Europe 900–700 Ma ago (so-called Central European Enriched Mantle). The Sm/Nd ratios of the Tatra migmatites range at 0.118 and are, thus, rather “normal” (i.e. typical for continental crust), but the high $\epsilon\text{Nd}_{(330)}$ may indicate the involvement of such enriched lithospheric mantle material. Migmatization during Variscan time might have caused mobilisation of parts of this old lithospheric mantle and of continental crust. This resulted in fractionation of the REE and finally shifted the Sm/Nd ratios to “normal” crustal values. Despite this possible explanation for the high Nd isotope ratios, it is not possible to rule out that the anatexis of the migmatites was too fast for a complete equilibration of the Sr and Nd isotopes. As noted by DAVIES and TOMMASINI (2000), equilibration of the Nd isotopes during anatexis of granitoid rocks may require several tens of millions of years. For the Tatra Mountains a rather

Tab. 3 Sm–Nd and Rb–Sr isotope data from the Western Carpathians (Western Tatra, High Tatra, Low Tatra, Fatra).

| sample | Nd ppm | Sm ppm | $\frac{^{143}\text{Nd}}{^{144}\text{Nd}}$ | $\pm 2\sigma$ mean | $\frac{^{147}\text{Sm}}{^{144}\text{Nd}}$ | ϵNd (330) | Sr ppm | Rb ppm | $\frac{^{87}\text{Sr}}{^{86}\text{Sr}}$ | $\pm 2\sigma$ mean | $\frac{^{87}\text{Rb}}{^{86}\text{Sr}}$ | ϵSr (330) |
|---|-----------|-----------|---|-----------------------|---|------------------------------|-----------|-----------|---|-----------------------|---|------------------------------|
| W–Tatra Orthogneisses (porphyric) | | | | | | | | | | | | |
| UP 1002 | 2.06 | 0.64 | 0.512178 | 33 | 0.12053 | –5.7 | 236.0 | 14.8 | 0.715059 | 13 | 0.18049 | 132.1 |
| UP 1014 | 28.92 | 5.40 | 0.512216 | 13 | 0.11286 | –4.7 | 375.3 | 24.8 | 0.711120 | 41 | 0.19073 | 76.2 |
| UP 1025 | 15.66 | 3.19 | 0.512337 | 26 | 0.12319 | –2.8 | 484.6 | 53.4 | 0.709588 | 13 | 0.31982 | 45.2 |
| W–Tatra Orthogneisses (leucocratic) | | | | | | | | | | | | |
| UP 1005 | 1.80 | 0.46 | 0.512067 | 20 | 0.15448 | –9.3 | 104.9 | 37.3 | 0.724154 | 12 | 1.02587 | 204.8 |
| UP 1012 | 2.00 | 0.54 | 0.512184 | 21 | 0.16300 | –7.4 | 50.9 | 117.6 | 0.763669 | 14 | 6.70700 | 387.2 |
| High Tatra Diorites (Xenoliths) | | | | | | | | | | | | |
| UP 1015 | 17.36 | 3.31 | 0.512491 | 20 | 0.11540 | 0.5 | 459.5 | 29.52 | 0.705037 | 79 | 0.18452 | –9.7 |
| UP 1016 | 10.40 | 2.27 | 0.512594 | 24 | 0.13213 | 1.8 | 345.9 | 32.35 | 0.705285 | 63 | 0.26798 | –11.7 |
| Western and High Tatra Granitoids / Migmatite | | | | | | | | | | | | |
| UP 1023 | 9.88 | 2.00 | 0.512315 | 32 | 0.12203 | –3.2 | 313.8 | 71.6 | 0.712230 | 20 | 0.65707 | 59.1 |
| UP 1036 | 39.62 | 8.74 | 0.512369 | 23 | 0.13343 | –2.6 | 364.1 | 24.1 | 0.708453 | 155 | 0.19131 | 38.3 |
| UP 1040 | 13.00 | 2.51 | 0.512317 | 31 | 0.11694 | –2.9 | 304.1 | 59.0 | 0.708852 | 23 | 0.55615 | 20.0 |
| UP 1017 | 22.44 | 4.56 | 0.512393 | 44 | 0.12290 | –1.7 | 556.3 | 100.3 | 0.708066 | 34 | 0.52306 | 10.7 |
| UP 1018 | 35.66 | 6.74 | 0.512328 | 31 | 0.11431 | –2.6 | 557.1 | 79.7 | 0.707641 | 26 | 0.41519 | 11.9 |
| UP 1044 | 7.20 | 1.65 | 0.512366 | 37 | 0.13848 | –2.8 | 123.3 | 27.7 | 0.711144 | 27 | 0.64280 | 46.4 |
| UP 1049 | 13.08 | 2.93 | 0.512420 | 67 | 0.13548 | –1.7 | 218.3 | 34.4 | 0.708517 | 65 | 0.45302 | 21.8 |
| UP 1050 | 14.76 | 3.13 | 0.512312 | 50 | 0.12804 | –3.5 | 284.8 | 40.8 | 0.707800 | 43 | 0.41405 | 14.2 |
| UP 1052 | 32.81 | 6.44 | 0.512562 | 34 | 0.11889 | 1.8 | 230.2 | 82.9 | 0.717762 | 29 | 1.03287 | 114.3 |
| UP 1053 | 33.60 | 6.45 | 0.512331 | 42 | 0.11612 | –2.6 | 436.1 | 59.6 | 0.710662 | 36 | 0.39557 | 56.1 |
| UP 1055 | 23.98 | 4.19 | 0.512376 | 23 | 0.10569 | –1.3 | 491.2 | 46.7 | 0.707725 | 30 | 0.27256 | 22.6 |
| UP 1057 | 19.96 | 3.77 | 0.512303 | 26 | 0.11407 | –3.1 | 371.5 | 24.2 | 0.707698 | 20 | 0.18707 | 27.9 |
| Fatra Granitoids | | | | | | | | | | | | |
| UP 1026 | 11.43 | 2.51 | 0.512322 | 33 | 0.13268 | –3.5 | 130.4 | 35.9 | 0.719053 | 57 | 2.05193 | 64.7 |
| UP 1027 | 23.76 | 4.20 | 0.512301 | 53 | 0.10692 | –2.8 | 506.3 | 10.6 | 0.708409 | 22 | 0.15532 | 40.1 |
| UP 1028 | 21.36 | 3.96 | 0.512361 | 78 | 0.11202 | –1.8 | 704.1 | 33.1 | 0.707838 | 37 | 0.13492 | 33.3 |
| UP 1030 | 23.78 | 4.53 | 0.512333 | 36 | 0.11522 | –2.5 | 401.8 | 29.3 | 0.709339 | 20 | 0.54365 | 27.4 |
| UP 1031 | 26.51 | 4.93 | 0.512292 | 45 | 0.11245 | –3.2 | 477.2 | 61.6 | 0.708925 | 49 | 0.37189 | 33.0 |
| UP 1038 | 6.75 | 1.34 | 0.512301 | 44 | 0.11961 | –3.3 | 132.6 | 99.5 | 0.716430 | 28 | 2.15152 | 20.9 |
| Low Tatra Granitoids | | | | | | | | | | | | |
| UP 1051 | 21.69 | 4.94 | 0.512144 | 55 | 0.13759 | –7.1 | 194.2 | 135.1 | 0.723573 | 67 | 2.01800 | 131.1 |
| UP 1088 | 16.90 | 2.99 | 0.512115 | 30 | 0.10715 | –6.4 | 147.1 | 92.5 | 0.723157 | 19 | 1.80444 | 139.5 |
| UP 1090 | 31.88 | 7.04 | 0.512379 | 22 | 0.13347 | –2.4 | 447.4 | 36.4 | 0.707550 | 23 | 0.23283 | 43.3 |

2σ mean errors refer to mean of the blocks.

rapid anatexis is proposed on the basis of U–Pb zircon dating. The migmatisation took place 335 Ma ago, and the last collisional granites (not affected by the migmatisation) intruded only 20 Ma later (POLLER and TODT, 2001). Therefore, a complete equilibration of Sr and Nd isotopes in the migmatites in such a short time span is at least questionable.

Tab. 4 Common Pb data for the Western Carpathian rocks.

| sample | $\frac{^{206}\text{Pb}}{^{204}\text{Pb}}$ | $\frac{^{207}\text{Pb}}{^{204}\text{Pb}}$ | $\frac{^{208}\text{Pb}}{^{204}\text{Pb}}$ |
|--|---|---|---|
| W-Tatra Orthogneisses (porphyric) | | | |
| 1002-1 | 18.2554 | 15.6482 | 38.4473 |
| 1002-2 | 18.9062 | 15.6837 | 38.4718 |
| 1014-1 | 18.7979 | 15.7202 | 39.1558 |
| 1014-2 | 18.8969 | 15.6859 | 39.9310 |
| 1025-2 | 18.7486 | 15.7470 | 38.8883 |
| W-Tatra Orthogneisses (leucocratic) | | | |
| 1005-1 | 18.0958 | 15.6363 | 38.2198 |
| 1005-2 | 18.1410 | 15.6647 | 38.2970 |
| 1012-1 | 17.9995 | 15.6193 | 37.9985 |
| 1012-2 | 18.1031 | 15.6065 | 37.9630 |
| Western and High Tatra Granitoids | | | |
| 1023-1 | 18.5946 | 15.6997 | 38.5827 |
| 1023-2 | 18.5675 | 15.6741 | 38.4885 |
| 1036-1 | 18.7033 | 15.6993 | 39.2420 |
| 1040-1 | 18.5368 | 15.6703 | 38.4764 |
| 1044-1 | 18.4125 | 15.6990 | 38.5024 |
| 1049-1 | 18.3485 | 15.6678 | 37.3983 |
| 1050-1 | 18.8499 | 15.6871 | 39.1130 |
| 1053-1 | 18.6178 | 15.6644 | 38.8183 |
| 1054-1 | 18.7587 | 15.7262 | 38.5105 |
| 1054-2 | 19.0834 | 15.7433 | 39.0427 |
| 1055-1 | 18.7476 | 15.7113 | 38.6565 |
| 1055-2 | 18.5029 | 15.6768 | 38.3930 |
| 1057-1 | 18.6883 | 15.7062 | 39.0218 |
| 1057-2 | 18.6503 | 15.7158 | 37.7916 |
| 1063-1 | 18.7123 | 15.7039 | 38.7899 |
| Low Tatra Granitoids | | | |
| 1088-1 | 18.8072 | 15.6829 | 38.5531 |
| 1090-1 | 18.6096 | 15.6754 | 38.5588 |
| 1092-1 | 18.4679 | 15.6676 | 38.5429 |
| 1094-1 | 18.7333 | 15.6979 | 39.4536 |
| 1095-1 | 18.6890 | 15.6727 | 38.4324 |
| Tatra Galenas | | | |
| 1089 | 18.4892 | 15.6738 | 38.4759 |
| 1092 | 18.4679 | 15.6676 | 38.5429 |
| Tatra Feldspars | | | |
| 1044-F | 18.4125 | 15.6990 | 38.5024 |
| 1056-F | 18.4925 | 15.6591 | 38.3344 |

All data are corrected for fractionation. The 2σ external reproducibility is about 300 ppm.

Discussion

Two main questions can be addressed with the data presented: (1) What source material is incorporated in the Tatra rocks? (2) Is there a close isotopic relationship between the Western Carpathian basement and the Variscan orogen? Finally, the geotectonic environment and evolution of the Tatra can be constrained.

SOURCE ROCK OF THE TATRA GRANITOIDS

According to the granite classification of BARBARIN (1999) the granitoids have most characteristics of the MPGs; only the An content of the plagioclases (20–45%) seems to be too high for typical crustal derived granodiorites and syenogranites. This is possibly due to less differentiation and fractional crystallisation in comparison to the orthogneisses.

However, granites with intermediate Sr and Nd isotopic composition and crustal Pb–Pb isotopic ratios usually are envisaged to be the result of mixing with more primitive material through direct input or by a heat source from the underplating basaltic magma (e.g. PIN and DUTHOU, 1990; COCHERIE et al., 1994). We interpret this feature as being inherited from repeatedly melted and recycled crust. Additionally, the granitoids are assumed to originate not only from old continental




Fig. 5 $\epsilon\text{Nd}_{(330)}$ versus $^{87}\text{Sr}/^{86}\text{Sr}_{(330)}$ data of the different Western Carpathian areas (Tatra and Fatra) are compared to also recalculated (to 330Ma) data from Bohemian Massif, Moldanubian unit (JANOUSEK et al., 1995), Black Forest, Spessart, Odenwald (LIEW and HOFMANN, 1988), and Massif Central (BEN OTHMAN et al., 1984; PIN and DUTHOU, 1990).

sediments but also may have incorporated mafic crustal material such as reworked amphibolites.

Even a minor contribution from a primitive source (such as ocean-derived amphibolites or even small amounts of lithospheric mantle material) is possible, mainly because of the presence of infracrustal mixed dioritic magma, which reached the anatexis zone during the collisional processes and caused the high isotopic ratios (the most negative $\epsilon\text{Nd}_{(330\text{Ma})}$ is -3.7) of the Tatra Mountain granitoids.

The origin of the Tatra rocks can be characterised in more detail by the radiogenic isotopes. As documented by the Pb–Pb data, the orthogneisses of the Tatra Mountains were generated mainly from upper crustal material. The Nd and Sr isotopic results indicate that a more mafic component was also involved, e.g. lithospheric mantle or recycled oceanic crust.

In the $^{206}\text{Pb}/^{204}\text{Pb}$ vs. $^{207}\text{Pb}/^{204}\text{Pb}$ diagram (Fig. 4b) of DOE and ZARTMAN (1979), the present Pb–Pb data from the Western Carpathians overlap both with the upper and lower crustal field. Nevertheless, a small amount of mafic, but mainly crustal, material should be responsible for the present Pb composition of Tatra rocks. This is supported by the recalculated values and fields shown in figure 4a. The Pb data were calculated back in time to 330 Ma, using several μ -values. This is a key age for the Western Carpathians (this is the time of emplacement of the dioritic xenoliths and final remelting of the migmatites). In figure 4a the positions of present-day isotopic compositions (same as in Figs 4b and 4c) are shown, together with those of 330 Ma for $\mu = 9$. The result is clear in that, the position of the Western Car-

pathians relative to the reference fields of upper crust, mid ocean ridge basalts (MORB) or enriched mantle II (EM II = recycled oceanic crust) does not change significantly. Consequently, the present-day Pb isotope data as shown in figures 4b and 4c reflects the position of the Western Carpathian rocks.

The leucocratic orthogneisses show significant input from reworked crust, which seems to be reasonable considering the crustal residence ages (~ 1750 Ma) of these orthogneisses (Fig. 6). Because of the high Sm/Nd ratios (> 0.155), these model ages were calculated using a two-step evolution. The leucocratic orthogneisses seem to be the most evolved rocks analysed in the Western Tatra. Obviously, their Sm/Nd ratios were changed by fractional crystallisation during the later metamorphic overprinting and the following granite intrusion.

The upper intercept U–Pb zircon ages of orthogneisses and granites, because of inherited cores, also reflect the influence of old crust. Consequently, the generation of the Tatra granitoids and gneisses seems to be essentially crustal, probably involving continentally collision and crustal thickening.

RELATIONSHIPS BETWEEN THE WESTERN CARPATHIANS AND THE VARISCIDES

The samples from the Tatra Mountains and from the neighbouring granitoid plutons of Velká Fatra (KOHÚT, 1992) and the Lower Tatra Mountains show close similarities in their isotopic composition. The Velká Fatra samples show $\epsilon\text{Sr}_{(330)}$ values




Fig. 6 Nd model ages calculated as one-step evolution (only for leucocratic orthogneisses: two step evolution after LIEW and HOFMANN, 1988). The Tatra rocks show 3 different groups of mean crustal residence ages.























between 21 and 65, depending on the amount of juvenile components. The corresponding $\epsilon\text{Nd}_{(330)}$ values of the Fatra rocks range from -2 to -4 . Granodiorites and syenogranites from the Lower Tatra Mountains show a general overlap with the Tatra Mountains with regard to Sr and Nd, as well as to Pb-Pb isotopic composition.

Comparisons with the Bohemian Massif, the Moldanubian units (JANOUSEK et al., 1995) and some regions from the western Variscides such as Massif Central, Black Forest, Spessart and Odenwald (LIEW and HOFMANN, 1988), indicate a close

isotopic relationship to the Western Carpathians (Fig. 5; all Sr and Nd literature data also were recalculated to 330 Ma).

The metasediments from the Lower Unit of the Western Tatra Mountains (GURK, 1999) show good isotopic correspondence with Moldanubian paragneisses. The isotopic character of the granites and orthogneisses of the Tatra Mountains and the Velká Fatra is consistent with several Variscan granitoids of France and Germany.

Only the diorites of the Tatra Mountains with high ϵNd (330) values (above zero) have no

| Age Unit | Protolith ages & Model ages | | Intrusion, Migmatitisation, Metamorphism | | | |
|-----------------------------------|---|---|---|---|---|--|
| | Archean-Proterozoic > 1600 Ma | < 1600 Ma | Devonian 410-380 Ma | 380-355 Ma | Carboniferous 355-325 Ma | 325-290 Ma |
| Low-Tatra Granites | |  1870-1420 Ma | | |  330±10 Ma | |
| Fatra Granites |  2520 ± 40 Ma |  1500-1200 Ma | | | |  310 ± 8 Ma |
| porphyric Ortho- gneisses |  1980±40 Ma |  1350-1200 Ma |  405±4 Ma |  360±10 Ma | | |
| leucocratic Ortho- gneisses |  1950-2400 Ma | | | | | |
| West-Tatra Granites |  2530±400 Ma |  1400-1240 Ma | |  360 - 350 Ma | | |
| High-Tatra Granites |  2000±98 Ma |  1450-1200 Ma | | | |  314±4 Ma |
| High-Tatra Migmatites |  2960 ± 15 Ma |  1450-1200 Ma | |  356±7 Ma |  332±5 Ma | |
| High-Tatra Diorites | |  990-970 Ma | | |  341±5 Ma | |







| Methods: | Interpretation: |
|--|---|
|  U-Pb "upper intercept ages |  Intrusion |
|  concordant U-Pb single zircon ages |  Metamorphism |
|  Nd - model ages |  Migmatitisation |

Fig. 7 Summary diagram for several rock series of the Western Carpathians including Nd model ages, U-Pb upper intercept ages indicating the protolith ages, and several concordant U-Pb ages defining intrusion, metamorphism, and migmatitisation.

equivalent rock series with the same emplacement age. However, similar ϵNd values between +2 and -8 are reported by FINGER et al. (1997) for post-orogenic granitoids from the Saxothuringian (Reichenberger granite), the Moldanubian (Freistadt granite) or even from the Alps (Düssi diorite and Venediger metatonalite). Also, geochemically similar rocks have been reported from the Limousin, the Massif Central (SHAW et al., 1993) and Corsica (COCHERIE et al., 1994).

Nd model ages for most of the investigated Western Carpathian granitoid rocks are between 1250 and 1450 Ma. Together with an average $^{147}\text{Sm}/^{144}\text{Nd}$ ratio of 0.12 they represent a crustal composition that occurs elsewhere in the European Variscides. Several Variscan regions, such as the Pyrenees (BEN OTHMAN et al., 1984) or the Mid German Crystalline Rise (REISCHMANN and ANTHES, 1996), show the same features. The Pb-Pb isotopes also confirm the isotopic correspondence between the Western Carpathian rocks and granitoids from the Black Forest, the Massif Central, and the Mid German Crystalline Rise (Fig. 4b). An overview of all isotopic data including the U-Pb zircons ages of the Tatra Mountains is given in figure 7.

GEODYNAMIC IMPLICATIONS FOR THE WESTERN CARPATHIANS

Based on the presented isotopic and geochemical data and U-Pb zircon ages mentioned here, we propose the following model for the geotectonic evolution of the Tatra Mountains.

The intrusion of the orthogneiss precursor 405 Ma ago probably was triggered by the subduction of oceanic crust under a continental plate at the northern border of Gondwana (POLLER et al., 2000). The orthogneisses in the Tatra Mountains are closely associated with partially melted banded amphibolites containing eclogite relics as well as kyanite-bearing migmatites, which were affected by medium- to high-pressure (HP) and high-temperature (HT) metamorphism reaching ~12 kbar and 750 °C (JANÁK et al., 1996; JANÁK et al., 1999). Therefore, the overprinting of the orthogneisses seems to have occurred under medium- to high-pressure and HT conditions in deep crustal levels. The absence of significant amounts of juvenile material in the orthogneisses suggests that the medium- to high-pressure and HT conditions were the result of crustal thickening related to continental collision. It is assumed that two microplates at the northern border of Gondwana were involved in this process (POLLER et al., 2000). Such a scenario accounts for the observed

petrographic, geochemical, and isotopic results. Similar ages as for the intrusion of the orthogneiss precursor (405 ± 4 Ma) are reported from rocks of the Mid German Crystalline Rise which are generally considered to be related to subduction zones (ALTENBERGER et al., 1990; ALTENBERGER and BESCH, 1993; REISCHMANN and ANTHES, 1996).

High pressure metamorphism analogous to that of the Tatra Mountains between 360 Ma and 350 Ma is known e.g. from the Erzgebirge Crystalline Complex. SCHMÄDICKE et al. (1995) reported Sm-Nd ages between 360 ± 7 Ma and 333 ± 6 Ma for eclogitic rocks of the Erzgebirge. Synmetamorphic granite intrusion (SCHULMANN et al., 1991), followed by exhumation and extension, are reported for the Bohemian Massif and the Moldanubian zone. Therefore, similarities between the Western Carpathians and the Western Variscides are documented not only by the geochemical and isotopic composition, but in ages and metamorphic evolution as well.

Immediately after the crustal thickening, resulting from microplate collision which was responsible for the overprint of the orthogneiss, the Western Tatra granites intruded (350 Ma, POLLER et al., 2000). The geotectonic evolution of the Tatra Mountains then underwent extension for a short time. Finally, the last granitoids intruded 315 Ma ago in the High Tatra Mountains. Such an Early Carboniferous magmatism with intrusion of large amounts of granitoids is documented for the Bohemian Massif as well (FRIEDL et al., 1994).

Final cooling took place between 330 and 300 Ma as recorded by Ar-Ar data (MALUSKI et al., 1993; JANÁK and ONSTOTT, 1993; JANÁK 1994). A similar evolution with collision-related processes, involving tectonic exhumation of high-grade units, has been described in well-documented parts of the Variscan belt (BURG et al., 1984; MATTE, 1986; O'BRIEN and CARSWELL, 1993; BROWN and DALLMEYER, 1996; ESCUDER VIRUETE et al., 1997; VON RAUMER, 1998).

Acknowledgements

We are very grateful to G. Feyerherd and I. Bambach for the preparation of the drawings and I. Raczek for her help with Nd and Sr analyses as well as constructive discussions and remarks. Ú.P. thanks A. Hofmann for the possibility to work in his institute, and for useful comments and careful reading of an early version of this manuscript. We thank C. Goodrich for the correction of the style. Discussions with F. Finger, A. Kröner and P. Uher contributed to the final version. Comments of H.J. Förster improved a previous version of the manuscript. Reviews of M. Thöni and M. Engi are acknowledged. This work was supported by the Max-Planck-Gesellschaft and the DFG (PO608/1-1 and PO608/1-3).

References

- ALTENBERGER, U. and BESCH, T. (1993): The Böllstein Odenwald: evidence for pre- to Early Variscan plate convergence in the Central European Variscides. *Geol. Rundsch.* 82, 475–488.
- ALTENBERGER, U., BESCH, T., MOCEK, B., ZAIPENG, Y. and YONG, S. (1990): Geochemie und Geodynamik des Böllsteiner Odenwaldes. *Mainzer Geowiss. Mitt.* 19, 183–200.
- ANDRUSOV, D. (1968): Grundriss der Tektonik der Nördlichen Karpathen. Veda Publishing House, Bratislava.
- ARNDT, N. and TODT, W. (1994): Formation of 1.9 Ga-old Trans-Hudson continental crust: Pb isotopic data. *Chem. Geol.* 118, 9–26.
- BARBARIN, B. (1996): Genesis of the two main types of peraluminous granitoids. *Geology* 24, 295–298.
- BARBARIN, B. (1999): A review of the relationships between granitoid types, their origins and their geodynamic environments. *Lithos* 46, 605–626.
- BEN OTHMAN, D., FOURCADE, S. and ALLÈGRE, C.J. (1984): Recycling processes in granite-granodiorite complex genesis: the Querigut case studied by Nd–Sr isotope systematics. *Earth Planet. Sci. Lett.* 74, 235–250.
- BROWN, M. and DALLMEYER, R.D. (1996): Rapid Variscan exhumation and the role of magma in core complex formation: southern Brittany metamorphic belt, France. *J. Metamorphic Geol.* 14, 361–379.
- BURCHART, J. (1968): Rubidium-strontium isochron ages of the crystalline core of the Tatra mountains, Poland. *Am. J. Sci.* 266, 895–907.
- BURG, J.P., LEYRELOUP, A., MARCHAND, J. and MATTE, Ph. (1984): Inverted metamorphic zonation and large scale thrusting in the Variscan Belt: an example in the French Massif Central. In: HUTTON, D.H.W. and SANDERSON, D.J. (eds): *Variscan Tectonics of the North Atlantic Region*. *Geol. Soc. London, Spec. Publ.* 14, 47–61.
- CHAPPELL, B.W., BRYANT C.J., WYBORN, D., WHITE, A.J.R. and WILLIAMS, I.S. (1998): High- and Low Temperature I-type Granites. *Resource Geology* 48, 225–235.
- CHAPPELL, B.W. and WHITE, A.J.R. (1974): Two contrasting granite types. *Pacific Geol.* 8, 173–174.
- COCHERIE, A., ROSSI, Ph., FOUILLAC, A.M. and VIDAL, Ph. (1994): Crust and mantle contributions to granite genesis – an example from the Variscan batholith of Corsica, France, studied by trace element and Nd–Sr–O isotope systematics. *Chem. Geol.* 115, 173–211.
- DAVIES, G.R. and TOMMASINI, S. (2000): Isotopic disequilibrium during rapid crustal anatexis: implications for petrogenetic studies of magmatic processes. *Chem. Geol.* 162, 169–191.
- DOE, B.R. and ZARTMAN, R.E. (1979): Plumbotectonics, The Phanerozoic. In: BARNES, H.L. (ed.): *Geochemistry of hydrothermal ore deposits*, Wiley Interscience Publication, New York, 22–70.
- ESCUDER VIRUETE, J., INDARES, A. and ARENAS, R. (1997): P-T path determination in the Tormes Gneissic Dome, NW Iberian Massif, Spain. *J. Metamorphic Geol.* 15, 645–663.
- EVENSEN, M.N., HAMILTON, P.J. and O'NIONS, R.K. (1978): Rare earth abundances in chondritic meteorites. *Geochim. Cosmochim. Acta* 42, 1199–1212.
- FINGER, F., ROBERTS, M.P., HAUNSCHMID, B., SCHERMAIER, A. and STEYRER, H.P. (1997): Variscan granitoids of central Europe: their typology, potential sources and tectonothermal relations. *Mineralogy and Petrology*, 61, 67–96.
- FRANKE, W. (1989): Tectonostratigraphic units in the Variscan belt of central Europe. *Geol. Soc. America, Spec. Paper* 230, 67–90.
- FRIEDL, G. and FINGER, F. (1994): Zur Intrusionsfolge im südböhmischen Batholith: Neue Aspekte bezüglich der Stellung des Eisgarner Granits. *Mitt. Österr. Mineral. Ges.* 139, 298–299.
- FRITZ, H., NEUBAUER, F., JANÁK, M. and PUTIS, M. (1992): Variscan mid-crustal thrusting in the Carpathians II: Kinematics and fabric evolution of the Western Tatra basement. *Terra Abstracts, Supplement 2 to Terra Nova* 4, 24.
- GAWEDA, A. (1995): Geochemistry and Rb/Sr isochron age of pegmatites from the Western Tatra Mountains. *Geologica Carpathica* 46, 95–99.
- GURK, C. (1999): Petrographie, Geochemie und Geochronologie der Unteren Einheit, Tatricum, Westliche Tatra (Slowakei). Unpubl. Diploma thesis, Univ. Mainz, 124 pp.
- JACOBSEN, S.B. and WASSERBURG, G.J. (1980): Sm–Nd evolution of chondrites. *Earth Planet. Sci. Lett.* 50, 139–155.
- JANÁK, M. (1994): Variscan uplift of the crystalline basement, Tatra Mountains, Central Western Carpathians: evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ laser probe dating of biotite and P-T-t paths. *Geol. Carpath.* 45, 293–300.
- JANÁK, M., HURAI, V., LUDHOVÁ, L., O'BRIEN, P.J. and HORN, E.E. (1999): Dehydration melting and devolatilization of high-grade metapelites: the Tatra Mountains, Western Carpathians. *J. Metamorphic Geol.* 17, 379–396.
- JANÁK, M., O'BRIEN, P.J., HURAI, V. and REUTEL, C. (1996): Metamorphic evolution and fluid composition of garnet-clinopyroxene amphibolites from the Tatra Mountains, Western Carpathians. *Lithos* 39, 57–79.
- JANÁK, M. and ONSTOTT, T.C. (1993): Pre-Alpine tectono-thermal evolution of metamorphism in the Tatra Mountains, Western Carpathians: P-T paths and $^{40}\text{Ar}/^{39}\text{Ar}$ laser probe dating. *Terra Abstracts, supplement 1 to Terra Nova* 5, 238.
- JANOUSEK, V., ROGERS, G. and BOWES, D.R. (1995): Sr–Nd isotopic constraints on the petrogenesis of the Central Bohemian Pluton, Czech Republic. *Geol. Rundsch.* 84, 520–534.
- KAHAN, S. (1969): Eine neue Ansicht über den geologischen Aufbau des Kristallinikums der West Tatra. *Acta Geologica et Geographica Universitatis Comenianae* 12, 115–122.
- KOHÚT, M., 1992. The Vel'ká Fatra granitoid pluton – an example of a Variscan zoned body in the Western Carpathians. In: VOZÁR, J. (ed.): *The Paleozoic geodynamic domains of the western Carpathians, Eastern Alps and Dinarides*. *Spec. Vol IGCP Project 276*, Bratislava, 79–92.
- KOHÚT, M., KOVACH, V.P., SALNIKOVA, E.B. and SAVATENKOV, V.M. (1999): Sr and Nd isotope geochemistry of Hercynian granitic rocks from the Western Carpathians – implications for granite genesis and crustal evolution. *Geol. Carpath.* 50, 477–487.
- KOHÚT, M. and JANÁK, M. (1994): Granitoids of the Tatra Mountains, Western Carpathians: field relationships and petrogenetic implications. *Geol. Carpath.* 45, 301–311.
- KOVÁČ, M., KRÁL, J., MARTON, E., PLASIENKA, D. and UHER, P. (1994): Alpine uplift history of the Central Western Carpathians: geochronological, paleomagnetic, sedimentary and structural data. *Geol. Carpath.* 45, 83–96.
- KRIST, E., KORIKOVSKY, S.P., PUTIS, M., JANÁK, M. and FARYAD, S.W. (1992): *Geology and Petrology of*

- metamorphic rocks of the Western Carpathian Crystalline Complexes. Comenius University Press, Bratislava, 324 pp.
- LAMEYRE, J., AUTRAN, A., BARRIÈRE, M., BONIN, B., DIDIER, J., FLUCK, P., FOURCADE, S., GIRAUD, P., JONIN, M., ORSINI, J.B. and VIVIER, G. (1980): Les granitoïdes de France. In: AUTRAN, A. and DERCOURT, J. (eds): *Evolutions Géologiques de la France*. Vol. 107, Mem. B.R.G.M., 51–97.
- LIEW, T.C. and HOFMANN, A.W. (1988): Precambrian crustal components, plutonic associations, plate environment of the Hercynian Fold Belt of Central Europe: Indications from a Nd and Sr isotopic study. *Contrib. Mineral. Petrol.* 98, 129–138.
- LISTER, G.S. and SNOKE, A.W. (1984): S–C mylonites. *J. Struct. Geol.* 6, 617–638.
- MALUSKI, H., RAJLICH, P. and MATTE, Ph. (1993): ^{40}Ar – ^{39}Ar dating of the Inner Carpathians Variscan basement and Alpine mylonitic overprint. *Tectonophysics* 223, 313–337.
- MATTE, Ph. (1986): Tectonics and Plate Tectonics Model for the Variscan Belt of Europe. *Tectonophysics* 126, 329–374.
- NEUBAUER, F. and VON RAUMER, J. (1993): The Alpine Basement-Linkage between Variscides and East-Mediterranean Mountain Belts. In: VON RAUMER, J. and NEUBAUER, F. (eds): *Pre-Mesozoic Geology of the Alps*, Springer, Berlin, Heidelberg, 641–663.
- NEUGEBAUER, J. (1988): The Variscan plate tectonic evolution: an improved “Iapetus model”. *Schweiz. Mineral. Petrogr. Mitt.* 68, 313–333.
- O'BRIEN, P.J. and ENGLAND, P.C. (1993): Tectonometamorphic evolution of the Bohemian Massif: evidence from high-pressure metamorphic rocks. *Geol. Rundsch.* 82, 531–555.
- PATTERSON, S.R., VERNON, R.H. and TOBISCH, O.T. (1989): A review of criteria for the identification of magmatic and tectonic foliations in granitoids. *J. Struct. Geol.* 11, 349–363.
- PIN, C. and DUTHOU, J.L. (1990): Sources of Hercynian granitoids from the French Massif Central: inferences from Nd isotopes and consequences for crustal evolution. *Chem. Geol.* 83, 281–296.
- POLLER, U., JANÁK, M., KOHÚT, M. and TODT, W. (2000): Pre-Variscan collision related magmatism in the Western Carpathians. U–Pb zircon data from granitoids and orthogneisses of the Tatra Mountains (Slovakia). *Int. J. Earth Sciences* 89, 336–349.
- POLLER, U. and TODT, W. (2001): U–Pb single zircon data of granitoids from the High Tatra Mountains (Slovakia): implications for the geodynamic evolution. *Transactions of Geol. Soc. London*, Vol. 91, 1/2, 235–243.
- REISCHMANN, T. and ANTHES, G. (1996): Geochronology of the mid-German crystalline rise west of the River Rhine. *Geol. Rundsch.* 85, 761–774.
- SCHMÄDICKE, E., MEZGER, K., COSCA, M.A. and OKRUSCH, M. (1995): Variscan Sm–Nd and Ar–Ar ages of eclogite facies rocks from the Erzgebirge, Bohemian Massif. *J. Metamorphic Geol.* 13, 537–552.
- SCHULMANN, K., LEDRU, P., AUTRAN, A., MELKA, R., LARDEAUX, J.M., URBAN, M. and LOBKOWICZ, M. (1991): Evolution of nappes in the eastern margin of the Bohemian massif: a kinematic interpretation. *Geol. Rundsch.* 80, 73–92.
- SHAW, A., DOWNES, H. and THIRLWALL, M.F. (1993): The quartz-diorites of Limousin: elemental and isotopic evidence for Devonian–Carboniferous subduction in the Hercynian belt of French massif Central. *Chem. Geol.*, 107, 1–18.
- STACEY, J.S. and KRAMERS, J.D. (1975): Approximation of terrestrial lead isotope evolution by two-stage model. *Earth Planet. Sci. Lett.* 26, 207–221.
- STILLE, P. and SCHALTEGGER, U. (1996): The Lithospheric Mantle Beneath Central Europe: Nd Isotopic Constraints for its Late Proterozoic Enrichment and implications for Early Crustal Evolution. In: BASU, A. and HART, S. (eds): *Earth Processes: reading the isotopic code*. Geophysical Monograph Series 95, Am. Geophys. Union, Washington, 269–276.
- VIDAL, P. and POSTAIRE, B. (1985): Étude par la méthode Pb–Pb de roches de haut grade métamorphique impliquées dans la chaîne Hercynienne. *Chem. Geol.* 49, 429–449.
- VON RAUMER, J.F. (1998): The Palaeozoic evolution in the Alps: from Gondwana to Pangea. *Geol. Rundsch.* 87, 407–435.
- WHITE, W.M. and PATCHETT, J. (1984): Hf–Nd–Sr isotopes and incompatible element abundances in island arcs: implications for magma origins and crust-mantle evolution. *Earth Planet. Sci. Lett.* 67, 167–185.
- ZIEGLER, P.A. (1986): Geodynamic model for the Paleozoic crustal consolidation of Western Central Europe. *Tectonophysics* 126, 303–328.
- ZINDLER, A. and HART, S. (1986): Chemical Geodynamics. *Ann. Rev. Earth Planet. Sci.* 14, 493–571

Manuscript received March 13, 2001; revision accepted June 6, 2001.

Appendix 1

SAMPLE DESCRIPTIONS

The investigated *orthogneisses* are represented by five samples. Samples UP 1002 (Ziarska valley), UP 1014 (Jamnická valley), and UP1025 (Baranec) are coarse grained, with porphyric, augen-like K-feldspar or plagioclase of 2–3 cm size, exhibiting a mylonitic S–C fabric (LISTER and SNOKE, 1988). Feldspars are elongated and show dynamic recrystallisation with characteristic tails. K-feldspar show typical microcline twinning. Plagioclase compositions range from albite to andesine, locally, myrmekite developed. Micas form characteristic “mica-fish” porphyroblasts. Muscovite is slightly phengitic, biotite is Fe-rich and usually replaced by chlorite. Quartz shows undulatory extinction with elongated and recrystallised grains, forming aggregates and ribbons. Minor garnet corresponds to almandine-spessartine, with up to 15% of pyrope component. However, garnet is strongly retrogressed and partly replaced by chlorite.

Samples UP 1005 (Ziarska valley) and UP 1012 (Jamnická valley) are leucocratic, fine-grained rocks, composed of plagioclase (dominantly albite), K-feldspar (microcline), and quartz. Muscovite is more abundant than biotite. Minor garnet was mostly replaced by chlorite. All investigated orthogneisses were affected by a solid-state deformation at ductile to brittle conditions (PATTERSON et al., 1989) and experienced strong regression.

The *granites* and *migmatites* are represented by three granitic samples from the Western Tatra and nine from the High Tatra. Additionally, two dioritic xenoliths of the High Tatra Mountains

and two migmatites were sampled. The granites of the Tatra generally are divided into two groups: the so-called High Tatra granitoids and the so-called common Tatra granitoids, occurring mostly in the Western Tatra (KOHÚT and JANÁK, 1994).

The “High Tatra granites” are granodioritic to tonalitic in composition. They contain plagioclase, quartz, rare K-feldspar, biotite, and rarely muscovite. In the High Tatra near Gerlachovsky peak, contacts between the High Tatra granitoids and some tonalitic to dioritic enclaves as well as contacts to garnet-sillimanite-bearing metapelites were observed. These metapelites may be deformed and show transition to migmatitic rocks (samples UP 1052; UP 1063). These migmatites were studied recently under petrological aspects and the initial migmatitisation is suggested to have taken place above 730 °C and 11–12 kbar (JANÁK et al., 1999).

The “common Tatra granitoids” are more intermediate in composition and can be characterised as monzogranites. They surface not only in the High Tatra, but also in the Western Tatra. They are composed of plagioclase, K-feldspar, biotite, and muscovite. Only sample UP 1036 (Bystrá) is free of muscovite and plagioclase is more abundant than K-feldspar.

The fabric of all analysed granites is homogeneous. The texture is holocrystalline, euhedral to subhedral feldspars are mostly randomly oriented and partly replaced by sericite. Micas are locally deformed into kink-bands, and quartz exhibits undulatory extinction. In general, the granitoids show only brittle and no penetrative deformation.

Appendix 2

ANALYTICAL TECHNIQUES

For the geochemical and geochronological investigation 10 to 20 kg of fresh material were collected in the field. A list with the exact sample locations is given in table 1. The samples were crushed, and approximately 300 g of the resulting material were pulverised using an agate mill. For mineral separation, the crushed material was ground using a rotary mill to a grain size < 500 µm.

Zircons were separated using a Wilfley table. The heaviest fraction was processed with heavy liquids and a Frantz magnetic separator to obtain the zircon fraction. Suitable grains for U–Pb

measurements and cathodoluminescence (CL) documentation were selected by hand picking.

For Pb–Pb whole rock analyses, fresh splits of the rocks were taken and dissolved in Savilex beakers during 3 days on the hotplate with HF and HNO₃. For Sm–Nd and Rb–Sr analyses as well as for XRF and ICP-MS measurements, the fine grained powder (<50 µm) from the agate mill was used.

Isotopic measurements were carried out at the Max-Planck-Institut für Chemie in Mainz using a Finnigan MAT 261 mass spectrometer.

Pb, Rb–Sr and Sm–Nd were measured with multiple collectors operating in static mode. Errors on isotopic ratios are given as 2σ errors of the block mean (10–25 blocks and 10 Scans each block). Blanks for Nd, Sm, Sr, and Rb analyses were below 100 pg and thus not significant. All Nd and Sr measurements were done as IC and ID runs. The Rb–Sr chemistry was performed by standard ion exchange methods. For Sm–Nd chemistry HDHP columns were used following the chemical procedures described in WHITE and PATCHETT (1984). For the Rb–Sr measurements a mixed $^{84}\text{Sr} - ^{85}\text{Rb}$ spike was used. The Sm–Nd spike was enriched in ^{150}Nd and ^{149}Sm . For both spikes the concentrations are known up to 0.1 % precision. Therefore the $^{87}\text{Rb}/^{86}\text{Sr}$ and $^{147}\text{Sm}/^{144}\text{Nd}$ ratios are limited to 0.1 % precision also. The ϵNd and ϵSr values are calculated for 330 Ma in table 3; figure 5 uses also recalculated (330 Ma) values.

The Pb–Pb procedure was done by HBr-chemistry on microcolumns (see ARNDT and TODT, 1994).

For Pb measurements, two NBS 982 or NBS 981 were loaded on each turret and measured at the beginning and at the end of the sample analyses. The fractionation was $1.45 \pm 0.3\%$ per amu. The blanks were less than 50 pg Pb total and therefore insignificant.

Geochemical analyses were performed by XRF at the Institut für Mineralogie, Johannes-

Gutenberg Universität Mainz. The measurements were controlled with respect to international standards.

The REE were analysed in the Department of Earth Sciences, the Memorial University of Newfoundland, St. Johns (Canada). Both, the major and trace element as well as the REE data are given in table 2.

The Pb data refer to present-day (Tab. 4). Recalculations of initial values for reasonable ages (e.g. 330 Ma, intrusion of diorites; migmatization of migmatites) gave no essential changes (see Fig. 4a). These calculations were done with several μ -values (ranging from 7 to 15) which resulted in different changes of the distance between the present day and the 330 Ma position. In all situations, the relative position of the Western Carpathian field and the reference fields of upper crust, MORB and enriched mantle II (after DOE and ZARTMAN, 1979) did not change significantly. Therefore, all values and fields in the $^{207}\text{Pb}/^{204}\text{Pb}$ vs. $^{206}\text{Pb}/^{204}\text{Pb}$ plots of figures 4b and 4c refer to present-day.

Additionally, the Pb whole rock analyses were controlled by feldspar and galena Pb analyses (crosses in Figs 4b and 4c). These mineral analyses show good correspondence with the whole rock data of the Western Carpathians.