

Zeitschrift: Bulletin für angewandte Geologie
Herausgeber: Schweizerische Vereinigung der Petroleum-Geologen und –Ingenieure;
Schweizerische Fachgruppe für Ingenieur-Geologie
Band: 12 (2007)
Heft: 1

Artikel: The geothermal conditions in the Rhine Graben : a summary
Autor: Rybach, Ladislaus
DOI: <https://doi.org/10.5169/seals-226366>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.06.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The geothermal conditions in the Rhine Graben - a summary

Ladislaus Rybach¹

A data base of borehole measurements collected over decades indicate that generally elevated temperatures prevail in the Rhine Graben (Fig. 1). For example at 1 km depth, instead of having the expected 40° C, the temperature range extends up to 140° C. The presentation at the 73rd ASP Convention addressed the geothermal conditions and processes, which are responsible for this feature.

The Rhine Graben, being part of the European Mid-Continental Rift System (Cloetingh et al. 2005), is characterized by thinned crust (and therefore higher heat flow; see Fig. 2), respectively by mantle proximity (e.g. ³He anomalies; Clauser et al. 2002). The temperature field is characterized by generally elevated values in the topmost 2 kilometers, in an inhomogeneous pattern: relatively high and low temperatures can be located close to each other (Pribnow & Schellschmidt 2000). Recently there is increasing evidence for convective heat transfer, at least in parts of the Rhine Graben subsurface (see e.g. Aquilina et al. 2000). Convection cells develop to quite some extent within the crystalline basement; numerical model simulations show that the cells can extend over several kilometres (Bächler et al. 2003). Water circulation velocity in the cells attains values up to cm/yr (Fig. 3 from Kohl et al. 2000). Convective heat transfer results in highly non-linear temperature-depth profiles, like the one measured at the European EGS project site at Soultz-sous-

Forêts (France): relatively high temperatures at shallow depth, and anomalously low temperature gradients underneath (Fig. 4). It is still an open question in which Graben realms could such features be present (potentially also at the Basel Deep Heat Mining site?); the data density for depths > 3 km is very low.

The Rhine Graben has generally favourable conditions for geothermal energy development and utilization (FESA 2005): generally elevated subsurface temperatures, regional aquifers at suitable depth (e.g. Buntsandstein, Upper Muschelkalk), extensional tectonics (provides vertical permeability). The first successful application, the Riehen/BS district heating system, started to operate in




Fig. 1: Compilation of borehole temperature measurements in the Rhine Graben (from Münch et al 2005; modified). Most data plot above the average geothermal gradient line with 30° C/km.

¹ Managing Director GEOWATT AG Zürich,
rybach@geowatt.ch

Summary of a talk given at the VSP/ASP annual convention, Rheinfelden, Switzerland, June 2006.

1994 and was later extended into Lörrach in Germany (Oppermann 2001). Nowadays there is a large number of geothermal projects, especially in Germany, at various stages of planning and realisation (Table 1; a further 26 projects have been initiated). Hopefully they will be successful.




Fig. 2: The Rhine Graben heat flow data (see box) confirm the general rule of decrease with increasing crustal thickness [early data from Čermák & Rybach 1979; later confirmed by new measurements of Clauser & Villinger 1995].




Fig. 3: Convection cell with details of the temperature and fluid flow field in the realm of well GPK2 at Soultz-sous-Forêts, France; result of numerical model calculations to fit the borehole temperature measurements in GPK2 (see Fig. 4). The flow velocities in the central model domain have been scaled down by a factor of 20 in order to represent even smaller components. The maximum fluid (= water) velocities exceed several cm/year. From Kohl et al. 2000.




Fig. 4: Temperature logs in the well GPK2 at Soultz-sous-Forêts, France: Comparison of measured (solid lines) and modelled (circles) temperatures. Convection causes high near-surface gradients, low gradients below and normal values further down. Crystalline basement top is also indicated. From Kohl et al. (2000), modified.

Project	Horizons	Depth	Temp.	Flow	Power	Status
Bruchsal	Buntsandstone	2000 m	130° C	> 20 l/s	500 kW	Circulating Power plant installation
Speyer	Buntsandstone	2900 m	150° C	> 20 l/s/well	5.4 MW	9 wells planned 1 st well completed
Offenbach	Muschelkalk	2800 m	150° C	100 l/s	4.8 MW	1 st well completed
Bellheim	Muschelkalk	ca. 2900 m, drilling	?	80-100 l/s	ca. 5 MW	1 st well completed
Riedstadt	Rotliegendes	2300 m	140° C	70 l/s	3 MW	Early preparation phase
Karlsruhe	Muschelkalk	3100 m	150° C	75 l/s	2.8 MW	Early preparation phase
Landau	Muschelkalk Buntsandstone Granite	3000 m	150° C	70 l/s	2.5 MW	1 st well completed 2 nd well in progress
Kehl	?	2500-3000 m	130° C	20-30 l/s	?	Early preparation phase
Ettenheim	Crystalline	5000 m	170° C	?	3-5 MW	Feasibility study

Tab. 1: Some geothermal utilization projects in the Rhine Graben (status February 2006, from Baumgärtner 2006).

References

- Aquilina, L., Genter, A., Elsass, P., Pribnow, D. (2000): Evolution of fluid circulation in the Rhine Graben: Constraints from the chemistry of present fluids. In: I. Stober & K. Bucher (eds.): *Hydrogeology of Crystalline Rocks*. Kluwer Academic Publishers, Dordrecht, 284 p.
- Baumgärtner, J. (2006): Geothermal exploration in the Upper Rhine Valley, Germany. EU Project ENGINE Launching Conference, <http://conferences-engine.brgm.fr/conf>.
- Bächler, D., Kohl, T., Rybach, L. (2003): Impact of graben-parallel faults on hydrothermal convection - the Rhine Graben case study. *Physics and Chemistry of the Earth* 28, 431-441.
- Čermák, V., Rybach, L. (eds.) (1979): *Terrestrial Heat Flow in Europe*. Springer Verlag, Heidelberg-Berlin-New York, 328 p.
- Clauser, C., Villinger, H. (1990): Analysis of conductive and convective heat transfer in a sedimentary basin, demonstrated for the Rhine Graben. *Geophys. J. Int.* 100, 393-414.
- Clauser, C., Griesshaber, E., Neugebauer, H.J. (2002): Decoupled thermal and mantle helium anomalies: Implications for the transport regime in continental rift zones. *J. Geophys. Res.* 107/B11, 2669, doi:10.1029/2001JB000675.
- Cloetingh, A., Ziegler, P.A., Beekman, F., Andriesen, P.A.M., Matenco, L., Bada, G., Garcia-Castellanos, D., Hardebol, N., Dèzes, P., Sokoutis, D. (2005): Lithospheric memory, state of stress and rheology: neotectonic controls on Europe's intraplate continental topography. *Quaternary Sci. Rev.* 24, 241-304.
- FESA (2005): *Geothermie am Oberrhein - Leitfaden und Marktführer für eine zukunftsfähige Energieform*. Förderverein Energie- und Solaragentur Regio Freiburg e.V. Freiburg, 72 p.
- Kohl, T., Bächler, D., Rybach, L. (2000): Steps towards a comprehensive thermo-hydraulic analysis of the HDR test site Soultz-sous-Forêts. In: *Proc. World Geothermal Congress 2000, Kyushu-Tohoku, Japan*, 2671-2676.
- Münch, W., Sisténich, H.P., Büker, Ch., Blanke, Th. (2005): Möglichkeiten der geothermischen Stromerzeugung im Oberrheingraben. - Eine Analyse der geologischen Bedingungen, der Bohr- und Fördertechnik sowie der Kraftwerkstechnik und Wirtschaftlichkeit. *VGB Power Tech* 10/2005, 1-8.
- Oppermann, G. (2001): Nachhaltigkeit über Landesgrenzen: Landesgrenzen überschreitende Nutzung der Geothermie in der Schweiz und in Deutschland. *Journal «mailing»* 1.01, 11-13.
- Pribnow, D., Schellschmidt, R. (2000): Thermal tracking of Upper Crustal fluid flow in the Rhine Graben. *Geophys. Res. Lett.* 27, 1957-1960.